

Appendix table A-1: Characteristics of included vs excluded participants from the cluster analysis.

	Overall N = 2,082	Excluded N = 701	Included N = 1,381	p-value
Age (years)	66.1 (7)	66.2 (6.6)	66 (7.1)	0.54
Sex				0.35
Women	970 (47%)	316 (45%)	654 (47%)	
Men	1,112 (53%)	385 (55%)	727 (53%)	
BMI (kg/m²)	27.7 (4.7)	29 (4.9)	27 (4.5)	<0.001
Waist circumference (cm)	96.7 (13.4)	99.4 (13.8)	95.3 (13)	<0.001
Visceral fat (cm²)	8.1 (2.8)	8.9 (3)	7.9 (2.7)	<0.001
Subcutaneous fat (cm ²)	2.5 (1.1)	2.6 (1.3)	2.5 (1.1)	0.053
HbA1c (%)	5.8 (0.5)	6 (0.7)	5.7 (0.4)	<0.001
HbA1c (mmol/mol)	39.8 (5.7)	41.7 (7.1)	38.8 (4.6)	<0.001
Smoking status				0.25
Current smoker	1,318 (64%)	455 (66%)	863 (63%)	
Never smoked	746 (36%)	238 (34%)	508 (37%)	
Physical activity	52.5 (24.9)	53.4 (26.5)	52 (24)	0.25

n (%); Mean (SD). Bold estimates indicate statistically significant differences between included and excluded participants (p < 0.05).