

Appendix table A-2: Characteristics of included and excluded participants from the regression analysis.

	Overall N = 1,381	Excluded N = 175	Included N = 1,206	p-value
Age (years)	66 (7.1)	66.4 (7.3)	66 (7.1)	0.50
Sex				0.82
Women	654 (47%)	81 (46%)	573 (48%)	
Men	727 (53%)	94 (54%)	633 (52%)	
BMI (kg/m²)	27 (4.5)	28.5 (5.2)	26.8 (4.4)	<0.001
Waist circumference (cm)	95.3 (13)	97.9 (13.2)	95 (13)	0.006
HbA1c (%)	5.7 (0.4)	5.8 (0.4)	5.7 (0.4)	<0.001
HbA1c (mmol/mol)	38.8 (4.6)	40 (4.9)	38.6 (4.5)	<0.001
AUC glucose	934 (925; 943)	1,003 (976; 1,030)	925 (915; 934)	<0.001
AUC insulin	23,410 (22,708; 24,133)	26,615 (24,128; 29,359)	22,978 (22,261; 23,717)	0.001
AUC GLP-1	2,733 (2,654; 2,815)	2,560 (2,326; 2,819)	2,759 (2,676; 2,845)	0.79
AUC GIP	5,050 (4,946; 5,155)	4,938 (4,650; 5,243)	5,066 (4,956; 5,178)	0.51
Glucagon suppression _{0-120 min}	33.9 (53)	35 (34.7)	33.7 (55.2)	0.67
Glycemic status				<0.001
NGT	734 (53%)	66 (38%)	668 (55%)	
iIFG	249 (18%)	29 (17%)	220 (18%)	
iiGT	122 (9%)	24 (14%)	98 (8%)	
IFG+IGT	116 (8%)	23 (13%)	93 (8%)	
Type 2 diabetes	160 (12%)	33 (19%)	127 (11%)	

n (%); Mean (SD); Geometric mean (95% CI) for AUC glucose, AUC insulin, AUC GLP-1 and AUC GIP.
 Bold estimates indicate statistically significant differences between included and excluded participants (p < 0.05).