

Appendix table A-4: Sensitivity analysis excluding individuals with screen-detected type 2 diabetes: Associations between cluster and standardized measures of visceral fat (VAT), subcutaneous fat (SAT), VAT/SAT ratio and body fat percentage.

Adjusted for age, sex, smoking status and physical activity	
	Estimate (95%CI)
Visceral fat (cm)	
Cluster 1	-0.02 (-0.14; 0.10)
Cluster 3	0.56 (0.43; 0.70)
Subcutaneous fat (cm)	
Cluster 1	0.24 (0.11; 0.37)
Cluster 3	0.29 (0.14; 0.44)
VAT/SAT ratio	
Cluster 1	-0.21 (-0.33; -0.09)
Cluster 3	0.10 (-0.04; 0.24)
Body fat (%)	
Cluster 1	0.13 (0.03; 0.24)
Cluster 3	0.55 (0.44; 0.67)

Reference: Cluster 2. Data are adjusted mean differences (95% CI) in standardized outcomes (SD units). VAT, SAT, and VAT/SAT ratio were log-transformed due to skewed distributions and standardized before regression analyses; body fat percentage was standardized on the original scale. Estimates are from the fully adjusted model excluding individuals with screen-detected type 2 diabetes. Bold estimates indicate statistically significant associations ($p < 0.05$).