

Supplementary table 3: Associations between cluster and non-standardized measures of VAT, SAT, VAT/SAT ratio and body fat percentage.

	Adjusted for age and sex	Further adjusted for smoking status and physical activity
	Estimate (95% CI)	Estimate (95% CI)
VAT (cm)		
Cluster 1	-1.0 (-4.8; 3.0)	-0.5 (-4.4; 3.6)
Cluster 3	26.0 (20.9; 31.3)	26.1 (21.0; 31.4)
SAT (cm)		
Cluster 1	12.3 (5.8; 19.2)	11.9 (5.4; 18.8)
Cluster 3	13.8 (6.9; 21.1)	13.7 (6.9; 21.0)
VAT/SAT ratio		
Cluster 1	-11.8 (-17.5; -5.7)	-11.1 (-16.8; -4.9)
Cluster 3	10.7 (3.3; 18.7)	10.8 (3.4; 18.8)
Body fat (%)		
Cluster 1	1.2 (0.4; 2.0)	1.2 (0.3; 2.0)
Cluster 3	5.1 (4.3; 6.0)	5.1 (4.2; 5.9)

Reference: Cluster 2. Data are expressed as percentage differences (95% CI) for VAT, SAT and VAT/SAT ratio, and as differences (95% CI) in percentage-points for body fat percentage. Models are adjusted for age and sex, with further adjustment for smoking status and physical activity. Bold estimates indicate statistically significant associations ($p < 0.05$).