

Table 2: Associations between cluster and standardized measures of visceral fat (VAT), subcutaneous fat (SAT), VAT/SAT ratio and body fat percentage.

	Adjusted for age and sex	Further adjusted for smoking status and physical activity
	Estimate (95% CI)	Estimate (95% CI)
Visceral fat (cm)		
Cluster 1	-0.03 (-0.15; 0.09)	-0.01 (-0.13; 0.10)
Cluster 3	0.68 (0.56; 0.81)	0.69 (0.56; 0.81)
Subcutaneous fat (cm)		
Cluster 1	0.25 (0.12; 0.38)	0.25 (0.11; 0.38)
Cluster 3	0.28 (0.15; 0.42)	0.28 (0.14; 0.42)
VAT/SAT ratio		
Cluster 1	-0.23 (-0.34; -0.11)	-0.21 (-0.33; -0.09)
Cluster 3	0.18 (0.06; 0.31)	0.18 (0.06; 0.31)
Body fat (%)		
Cluster 1	0.15 (0.05; 0.25)	0.14 (0.04; 0.24)
Cluster 3	0.63 (0.52; 0.73)	0.62 (0.51; 0.72)

Reference: Cluster 2. Data are adjusted mean differences (95% CI) in standardized outcomes (SD units). VAT, SAT, and VAT/SAT ratio were log-transformed due to skewed distributions and standardized before regression analyses and body fat percentage was standardized on the original scale. Models are adjusted for age and sex, with further adjustment for smoking status and physical activity. Bold estimates indicate statistically significant associations ($p < 0.05$).