


TECHNIS UNIVERS

Covid-19 Mutation Dynamics



- **Reproduction period** τ : average time interval between two infection generations
- Actively infected #persons x in the 'E' or 'I' states
- ▶ Base reproduction numbers R_0^{wild} and R_0^{mut} of the wild type and the mutated strain (e.g., B.1.1.7), respectively
- Penetration rate p of the mutated strain

$$\begin{split} x(t+\tau) &= R_0 x \stackrel{!}{=} (1-p) R_0^{\text{wild}} x + p R_0^{\text{mut}} x \\ \Rightarrow R_0 &= (1-p) R_0^{\text{wild}} + p R_0^{\text{mut}} \end{split}$$



- **Reproduction period** au: average time interval between two infection generations
- ► Actively infected #persons *x* in the 'E' or 'I' states
- **Base reproduction numbers** R_0^{wild} and R_0^{mut} of the wild type and the mutated strain (e.g., B.1.1.7), respectively
- Penetration rate p of the mutated strain

$$x(t+\tau) = R_0 x \stackrel{!}{=} (1-p) R_0^{\text{wild}} x + p R_0^{\text{mut}} x$$
$$\Rightarrow R_0 = (1-p) R_0^{\text{wild}} + p R_0^{\text{mut}}$$



- **Reproduction period** τ : average time interval between two infection generations
- ► Actively infected #persons *x* in the 'E' or 'I' states
- **Base reproduction numbers** R_0^{wild} and R_0^{mut} of the wild type and the mutated strain (e.g., B.1.1.7), respectively
- Penetration rate p of the mutated strain

$$x(t+\tau) = R_0 x \stackrel{!}{=} (1-p) R_0^{\mathsf{wild}} x + p R_0^{\mathsf{mut}} x$$
$$\Rightarrow R_0 = (1-p) R_0^{\mathsf{wild}} + p R_0^{\mathsf{mut}}$$



- **Reproduction period** au: average time interval between two infection generations
- Actively infected #persons x in the 'E' or 'I' states
- **Base reproduction numbers** R_0^{wild} and R_0^{mut} of the wild type and the mutated strain (e.g., B.1.1.7), respectively
- Penetration rate p of the mutated strain

$$x(t+\tau) = R_0 x \stackrel{!}{=} (1-p) R_0^{\mathsf{wild}} x + p R_0^{\mathsf{mut}} x$$
$$\Rightarrow R_0 = (1-p) R_0^{\mathsf{wild}} + p R_0^{\mathsf{mut}}$$



- **Reproduction period** au: average time interval between two infection generations
- ightharpoonup Actively infected #persons x in the 'E' or 'I' states
- **Base reproduction numbers** R_0^{wild} and R_0^{mut} of the wild type and the mutated strain (e.g., B.1.1.7), respectively
- Penetration rate p of the mutated strain

$$x(t+\tau) = R_0 x \stackrel{!}{=} (1-p) R_0^{\mathsf{wild}} x + p R_0^{\mathsf{mut}} x$$
$$\Rightarrow R_0 = (1-p) R_0^{\mathsf{wild}} + p R_0^{\mathsf{mut}}$$

Dynamics of the **percentage** p of the new strain:

$$p(t+\tau) = \frac{x^{\mathsf{mut}}(t+\tau)}{x^{\mathsf{wild}}(t+\tau)} = \frac{pR_0^{\mathsf{mut}}}{(1-p)R_0^{\mathsf{wild}}}$$

Dynamics of the **odds ratio** y = p/(1-p)

$$y(t+\tau) = \frac{R_0^{\text{mut}}}{R_0^{\text{wild}}} \ y(t)$$

Formulation as a differential equation:

$$\frac{\mathrm{d}y}{\mathrm{d}t} \approx \frac{y(t+\tau) - y(t)}{\tau} = \frac{1}{\tau} \left(\frac{R_0^{\mathsf{mut}}}{R_0^{\mathsf{wild}}} - 1 \right) y \stackrel{!}{=} r_y y \tag{1}$$

$$p = \frac{y}{1+y}, \quad y(t) = y_0 e^{r_y t}, \quad p(t) = \frac{y_0 e^{r_y t}}{1+y_0 e^{r_y t}} = \frac{p_0 e^{r_y t}}{1+p_0 \left(e^{r_y t}-1\right)}$$



Dynamics of the **percentage** p of the new strain:

$$p(t+\tau) = \frac{x^{\mathsf{mut}}(t+\tau)}{x^{\mathsf{wild}}(t+\tau)} = \frac{pR_0^{\mathsf{mut}}}{(1-p)R_0^{\mathsf{wild}}}$$

Dynamics of the **odds ratio** y = p/(1-p):

$$y(t+\tau) = \frac{R_0^{\text{mut}}}{R_0^{\text{wild}}} \ y(t)$$

Formulation as a differential equation:

$$\frac{\mathrm{d}y}{\mathrm{d}t} \approx \frac{y(t+\tau) - y(t)}{\tau} = \frac{1}{\tau} \left(\frac{R_0^{\mathsf{mut}}}{R_0^{\mathsf{wild}}} - 1 \right) y \stackrel{!}{=} r_y y \tag{1}$$

$$p = \frac{y}{1+y}, \quad y(t) = y_0 e^{r_y t}, \quad p(t) = \frac{y_0 e^{r_y t}}{1+y_0 e^{r_y t}} = \frac{p_0 e^{r_y t}}{1+p_0 (e^{r_y t} - 1)}$$



Dynamics of the **percentage** p of the new strain:

$$p(t+\tau) = \frac{x^{\mathsf{mut}}(t+\tau)}{x^{\mathsf{wild}}(t+\tau)} = \frac{pR_0^{\mathsf{mut}}}{(1-p)R_0^{\mathsf{wild}}}$$

Dynamics of the **odds ratio** y = p/(1-p):

$$y(t+\tau) = \frac{R_0^{\text{mut}}}{R_0^{\text{wild}}} \ y(t)$$

Formulation as a differential equation:

$$\frac{\mathrm{d}y}{\mathrm{d}t} \approx \frac{y(t+\tau) - y(t)}{\tau} = \frac{1}{\tau} \left(\frac{R_0^{\mathsf{mut}}}{R_0^{\mathsf{wild}}} - 1 \right) y \stackrel{!}{=} r_y y \tag{1}$$

$$p = \frac{y}{1+y}, \quad y(t) = y_0 e^{r_y t}, \quad p(t) = \frac{y_0 e^{r_y t}}{1+y_0 e^{r_y t}} = \frac{p_0 e^{r_y t}}{1+p_0 (e^{r_y t} - 1)}$$



Dynamics of the **percentage** p of the new strain:

$$p(t+\tau) = \frac{x^{\mathsf{mut}}(t+\tau)}{x^{\mathsf{wild}}(t+\tau)} = \frac{pR_0^{\mathsf{mut}}}{(1-p)R_0^{\mathsf{wild}}}$$

Dynamics of the **odds ratio** y = p/(1-p):

$$y(t+\tau) = \frac{R_0^{\text{mut}}}{R_0^{\text{wild}}} \ y(t)$$

Formulation as a differential equation:

$$\frac{\mathrm{d}y}{\mathrm{d}t} \approx \frac{y(t+\tau) - y(t)}{\tau} = \frac{1}{\tau} \left(\frac{R_0^{\mathsf{mut}}}{R_0^{\mathsf{wild}}} - 1 \right) y \stackrel{!}{=} r_y y \tag{1}$$

$$p = \frac{y}{1+y}$$
, $y(t) = y_0 e^{r_y t}$, $p(t) = \frac{y_0 e^{r_y t}}{1+y_0 e^{r_y t}} = \frac{p_0 e^{r_y t}}{1+p_0 (e^{r_y t} - 1)}$



Given:

- lacktriangle Observed strain fractions p_1 and p_2 at the times t_1 and t_2
- lacktriangle Observed or simulated base reproduction number R_0 at time t

Wanted: dynamic parameters r_y , R_0^{wild} and R_0^{mut} , and all the rest

Solution

• use the relation $y(t) = y_0 e^{r_y(t-t_0)}$ to estimate the growth rate r_y :

$$r_y \approx (\ln y(t_2) - \ln y(t_1))/(t_2 - t_1)$$

Use Relation (1) to estimate the reproduction number ratio

$$R_0^{\mathsf{mut}}/R_0^{\mathsf{wild}} pprox au r_y + 1$$

 \blacktriangleright Use the observed/simulated total base reproduction number to estimate the R_0 's individually:

Given:

- lacktriangle Observed strain fractions p_1 and p_2 at the times t_1 and t_2
- lacktriangle Observed or simulated base reproduction number R_0 at time t

Wanted: dynamic parameters r_y , R_0^{wild} and R_0^{mut} , and all the rest

Solution

• use the relation $y(t) = y_0 e^{r_y(t-t_0)}$ to estimate the growth rate r_y :

$$r_y \approx (\ln y(t_2) - \ln y(t_1))/(t_2 - t_1)$$

Use Relation (1) to estimate the reproduction number ratio

$$R_0^{\mathsf{mut}}/R_0^{\mathsf{wild}} pprox au r_y + 1$$

 \blacktriangleright Use the observed/simulated total base reproduction number to estimate the R_0 's individually:

$$R_0^{\mathrm{wild}} = \frac{R_0(t)}{1 + p(t)\tau r_y}, \quad R_0^{\mathrm{mut}} = (\tau r_y + 1)R_0^{\mathrm{wild}}$$

Given:

- lacktriangle Observed strain fractions p_1 and p_2 at the times t_1 and t_2
- lacktriangle Observed or simulated base reproduction number R_0 at time t

Wanted: dynamic parameters r_y , R_0^{wild} and R_0^{mut} , and all the rest

Solution:

• use the relation $y(t) = y_0 e^{r_y(t-t_0)}$ to estimate the growth rate r_y :

$$r_y \approx (\ln y(t_2) - \ln y(t_1))/(t_2 - t_1)$$

Use Relation (1) to estimate the reproduction number ratio:

$$R_0^{\mathsf{mut}}/R_0^{\mathsf{wild}} pprox au r_y + 1$$

 \blacktriangleright Use the observed/simulated total base reproduction number to estimate the R_0 's individually:

$$R_0^{\mathrm{wild}} = \frac{R_0(t)}{1 + p(t)\tau r_u}, \quad R_0^{\mathrm{mut}} = (\tau r_y + 1)R_0^{\mathrm{wild}}$$

Given:

- $lackbox{ Observed strain fractions } p_1 \text{ and } p_2 \text{ at the times } t_1 \text{ and } t_2$
- lacktriangle Observed or simulated base reproduction number R_0 at time t

Wanted: dynamic parameters r_y , R_0^{wild} and R_0^{mut} , and all the rest

Solution:

• use the relation $y(t) = y_0 e^{r_y(t-t_0)}$ to estimate the growth rate r_y :

$$r_y \approx (\ln y(t_2) - \ln y(t_1))/(t_2 - t_1)$$

▶ Use Relation (1) to estimate the reproduction number ratio:

$$R_0^{\mathrm{mut}}/R_0^{\mathrm{wild}} pprox au r_y + 1$$

ightharpoonup Use the observed/simulated total base reproduction number to estimate the R_0 's individually:

$$R_0^{\mathrm{wild}} = \frac{R_0(t)}{1 + p(t)\tau r_y}, \quad R_0^{\mathrm{mut}} = (\tau r_y + 1)R_0^{\mathrm{wild}}$$

Given:

- lacktriangle Observed strain fractions p_1 and p_2 at the times t_1 and t_2
- lacktriangle Observed or simulated base reproduction number R_0 at time t

Wanted: dynamic parameters r_y , R_0^{wild} and R_0^{mut} , and all the rest

Solution:

• use the relation $y(t) = y_0 e^{r_y(t-t_0)}$ to estimate the growth rate r_y :

$$r_y \approx (\ln y(t_2) - \ln y(t_1))/(t_2 - t_1)$$


▶ Use Relation (1) to estimate the reproduction number ratio:

$$R_0^{\mathrm{mut}}/R_0^{\mathrm{wild}} pprox au r_y + 1$$


lackbox Use the observed/simulated total base reproduction number to estimate the R_0 's individually:

$$R_0^{\mathrm{wild}} = \frac{R_0(t)}{1 + p(t)\tau r_y}, \quad R_0^{\mathrm{mut}} = (\tau r_y + 1)R_0^{\mathrm{wild}}$$










Mutation





Example Germany, 2021-03-12

