


Scenario projections of the Covid-19 pandemic using a data-driven macroscopic model

Martin Treiber, TU Dresden


R_0 wild=1.65
 R_0 mut=2.51
 p mut=100%
 R_0 =2.51



Scenario projections of the Covid-19 pandemic using a data-driven macroscopic model

Martin Treiber, TU Dresden

$R_0 \text{ wild} = 1.65$
 $R_0 \text{ mut} = 2.51$
 $p_{\text{mut}} = 100\%$
 $R_0 = 2.51$




- ▶ 1. The SEIR Model with Delays and Test Dynamics
 - ▶ 1.1 Modelling the mutation dynamics
 - ▶ 1.2 Modelling the age-specific IFR
- ▶ 2. Scenario Projections

Scenario projections of the Covid-19 pandemic using a data-driven macroscopic model

Martin Treiber, TU Dresden

R_0 wild=1.65
 R_0 mut=2.51
 p mut=100%
 R_0 =2.51




- ▶ 1. The SEIR Model with Delays and Test Dynamics
 - ▶ 1.1 Modelling the mutation dynamics
 - ▶ 1.2 Modelling the age-specific IFR
- ▶ 2. Scenario Projections

Scenario projections of the Covid-19 pandemic using a data-driven macroscopic model

Martin Treiber, TU Dresden

R_0 wild=1.65
 R_0 mut=2.51
 p mut=100%
 $R_0=2.51$



- ▶ 1. The SEIR Model with Delays and Test Dynamics
 - ▶ 1.1 Modelling the mutation dynamics
 - ▶ 1.2 Modelling the age-specific IFR
- ▶ 2. Scenario Projections


1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments

Susceptible (S)

1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics

- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1. Macroscopic SEIR model with explicit delays and test dynamics


- ▶ Different infection *phases* (compartments) and their *transitions*
- ▶ Macroscopic dynamic variables: fractions of the population in the compartments




1.1 Modelling the mutation dynamics



1.1 Modelling the mutation dynamics




1.1 Modelling the mutation dynamics



Odds ratio

$$y = \frac{p}{1 - p}$$

1.1 Modelling the mutation dynamics




Odds ratio

$$y = \frac{p}{1-p}$$

$$y = y_0 e^{r_y t}$$

1.1 Modelling the mutation dynamics



Odds ratio


$$y = \frac{p}{1 - p}$$

$$y = y_0 e^{r_y t}$$

Estimation growth rate

$$r_y \approx \frac{\ln y(t_2) - \ln y(t_1)}{t_2 - t_1}$$

1.1 Modelling the mutation dynamics



Odds ratio

$$y = \frac{p}{1 - p}$$

$$y = y_0 e^{r_y t}$$

Estimation growth rate


$$r_y \approx \frac{\ln y(t_2) - \ln y(t_1)}{t_2 - t_1}$$

Estimation B.1.1.7 share

$$p = \frac{y}{1 + y}$$

$$= \frac{p_0 e^{r_y t}}{1 + p_0 (e^{r_y t} - 1)}$$

1.1 Modelling the mutation dynamics



Odds ratio

$$y = \frac{p}{1 - p}$$

$$y = y_0 e^{r_y t}$$

Estimation growth rate


$$r_y \approx \frac{\ln y(t_2) - \ln y(t_1)}{t_2 - t_1}$$

Estimation B.1.1.7 share

$$p = \frac{y}{1 + y}$$

$$= \frac{p_0 e^{r_y t}}{1 + p_0 (e^{r_y t} - 1)}$$


1.1 Modelling the mutation dynamics



τ : generation time

$$R_0^{\text{mut}} / R_0^{\text{wild}} = \tau r_y + 1$$

1.1 Modelling the mutation dynamics




τ : generation time

$$R_0^{\text{mut}} / R_0^{\text{wild}} = \tau r_y + 1$$

Effective R_0 :

$$R_0 = (1 - p)R_0^{\text{wild}} + pR_0^{\text{mut}}$$

1.1 Modelling the mutation dynamics



τ : generation time

$$R_0^{mut}/R_0^{wild} = \tau r_y + 1$$


Effective R_0 :

$$R_0 = (1 - p)R_0^{wild} + pR_0^{mut}$$

$$\Rightarrow R_0^{wild}, R_0^{mut} \text{ known}$$


$$\Rightarrow \text{forecast } R_0(t)$$

1.2 Modelling the age-specific infection fatality rate (IFR)




- ▶ The IFR exhibits a nearly exponential growth with the age
- ▶ The same is true for the case fatality rate (CFR) and for the total one-year fatality
- ▶ The gender difference is weak (corresponds to ≈ 5 years)
- ▶ The infection probability exhibits a much weaker age dependency
- ▶ \Rightarrow leave infection spread model unchanged (enables interactive just-in-time calibration) and calculate the IFR in age strata afterwards

1.2 Modelling the age-specific infection fatality rate (IFR)




- ▶ The IFR exhibits a nearly exponential growth with the age
- ▶ The same is true for the case fatality rate (CFR) and for the total one-year fatality
- ▶ The gender difference is weak (corresponds to ≈ 5 years)
- ▶ The infection probability exhibits a much weaker age dependency
- ▶ \Rightarrow leave infection spread model unchanged (enables interactive just-in-time calibration) and calculate the IFR in age strata afterwards


Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}




Calculating the global IFR as multiple of IFR_{65}



Calculating the global IFR as multiple of IFR_{65}



Calculating the global IFR as multiple of IFR₆₅



Interactive simulations

