

# Cardiac Output (CO) & Cardiac reserve

---

Physiology department merit university

# ILOS

- Describe cardiac output and state components controlling it.
- Define various related terms, including stroke volume, cardiac index, end diastolic and end systolic volumes.
- Explain the importance of cardiac reserve, mechanisms and limitations, and heart failure (pathophysiology, causes and consequences).



# Cardiac output

- **Cardiac output (COP):** Is the volume of blood pumped by each ventricle per minute.

About 5L\min during rest.

$$\text{COP} = \text{SV} \times \text{HR} = 70 \times 70 = 4900 \text{ ml}\backslash\text{min}$$

- The CO depends on:

1- stroke volume (SV)

2- heart rate (HR)

## definitions

- **Stroke volume:** is the volume of blood pumped by each ventricle per beat.

About 70-90 ml/beat during rest

- **Cardiac output (COP):** Is the volume of blood pumped by each ventricle per minute.

About 5L/min during rest.



## definitions

- **Cardiac Index (CI):** Volume of blood pumped by each ventricle per minute **per** square meter body surface area.

Normally = 3.2 L/min/m<sup>2</sup>

- **End Diastolic Volume:** is the volume of blood at the ventricle at the end of the diastole. About (110-130 ml).
- **End systolic ventricle:** is the volume of blood at the ventricle at the end of the systole. About (40-60 ml).
- $SV = EDV - ESV = 130 - 60 = 70ML$

# definitions

## Ejection Fraction (EF):

Definition: EF is the fraction of the EDV that is ejected with each beat

## Calculation:

$$EF = \frac{\text{Stroke volume}}{\text{End diastolic volume}} \times 100$$

Normally: Ejection fraction is 55-75%

Importance : Assessment of Contractility.



## **Variations of cardiac output (CO):**

**UNCHANGED**

**1- SLEEP**

**2- MODERATE CHANGES IN EXTERNAL TEMPRATURE.**

## ***Variations of cardiac output (CO):***

### INCREASED IN

- 1- **E**XIETMENT 50-100%
- 2- **E**XERCISE 300%
- 3- **E**XPOSURE TO HIGH TEMPRATURE
- 4- **E**EATING
- 5- **E**PINEPHRINE
- 6- **E**ND OF PREGNANCY
- 8- INSPIRATION
- 9- SHIFTING FROM STANDING TO RECOMBINANT POSITION. (Gravity → decreases VR )



# **Variations of cardiac output (CO):**

## **DECREASED IN**

- 1- SITTING OR STANDING FROM LYING POSITION (gravity)**
- 2- ARRYTHMEA**
- 3- HEART FAILURE**

# **Control of the cardiac output**

1. HEART RATE REGULATION.
2. CHANGES IN PRELOAD.
3. EFFECTS OF AFTERLOAD.
4. EFFECTS OF CONTRACTILITY.

NB: 2,3,4 = factor regulating Stroke volume(preload, afterload, contractility).



# 1-Heart Rate(HR)

- An increase of HR up to 200/min. (eg during exercise) leads to an increase of CO (????)→ during exercise both HR & SV( increased VR) increase→ COP increase.
- Moderate changes in heart rate(50-100 beat\min) : no change in COP  
?????? The increase or decrease in HR is corrected by SV, **EXAMPLE:**

When heart rate increased to 100 beat \min→ shortening of the diastole→ SV decreases to 50 ml so  $cop = SV \times HR = 100 \times 50 = 5000 \text{ ml}\backslash\text{min}$

## 1-Heart Rate(HR)

- Pathological or excessive change in HR → decrease COP ???

In paroxysmal tachycardia → heart rate is increased but stroke volume is decreased → shortened diastole & decreased contractility (over use of energy).



## 2-Change of Preload

- ❑ **Preload determines muscle length before start of contraction.**
- ❑ **Ventricular preload is determined by EDV**
- ❑ **EDV depends on venous return**
- ❑ **According to Starling's Law:**
  - The more the end diastolic volume the more is the force of contractions; within limits.
  - An increase of EDV (preload) leads to increase SV with consequent increase of CO

## 2-Change of Preload

- Preload is affected mainly by venous return
- **Factors that increase venous return**: Means increase filling of ventricles increased EDV → increased myocardial contractility → increased SV → increased cardiac output



## 2-Change of Preload:

### - Factors that affect ventricular preload:

1. **Venous Return (VR):** Increased venous return will increase preload
2. **Ventricular compliance:** Decreased ventricular compliance will decrease preload eg myocardial infarction
3. **Heart rate:** Increased rate more than 200/min leads to a decrease of preload due to shortening of diastole decreased ventricular filling

### 3- Effect of afterload on SV

- After load is the load against which the ventricle contracts and eject blood
- The main after load for left ventricle is the **aortic pressure**.
- An **increase** of after load leads to a **decrease** of SV → decrease CO.
- **Decreased** after load leads to **increase** of SV → increase CO.



## 4- Effect of contractility (inotropic state) on SV

- Positive inotropic → increase contractility of the ventricle → ejecting more blood → increase SV → increase CO.

# *Cardiac Reserve "CR"*

- **Definition:** It is difference between cardiac output pumped by the heart under ordinary circumstances & its maximum capacity for pumping blood

**Assessed by:**

Low-dose dobutamine stress  
echocardiography



# Mechanisms of Cardiac Reserve CR

## 1- +ve chronotropic effects

**(Heart rate reserve)**

**During** HR = 70

**rest:**

**During** HR = 200 (so, COP can increase 3 times)

**exercise:**

**Limitation:** Diastolic period is shortened

# Mechanisms of Cardiac Reserve CR

## 2- factors that increase the preload

- During rest: = 135 ml (EDV reserve)
- During exercise: **Venous return increase** led to **increase** ventricular filling (increase in the initial length) → **increase** EDV → lead to **increase** force of contraction
- **Limitation:** over stretch as in heart failure → decrease contractility

## 3- factors that decrease the afterload

- Factors that decrease the peripheral resistance ( will be discussed later)



# Mechanisms of Cardiac Reserve CR

## 4- +ve inotropic conditions:

During rest the ESV is 50-60 ml.

*During exercise* sympathetic stimulation increase the force of contraction → the ESV decrease → increase of SV → increase in COP.

**Limitation:** too much decrease in ESV → injury to endocardium.

# Mechanisms of Cardiac Reserve CR

5-Cardiac hypertrophy: physiological (athletes) or pathological (hypertension)

hypertrophy → increase contractility → increase SV → increase COP.

**Limitation:** relative ischemia as ventricular capillaries do not proliferate



FINALLY;

*If there is no matching between COP & body needs: so,  
this Heart is Failed (HF)*

# Heart Failure


## ILOs

1. Describe cardiac output and state components controlling it.
2. Define various related terms, including stroke volume, cardiac index, end diastolic and end systolic volumes.
3. Explain the importance of cardiac reserve, mechanisms and limitations, and heart failure (pathophysiology, causes and consequences).



# Heart Failure

- Definition: It is a clinical syndrome in which the heart is unable to pump an adequate amount of blood to satisfy metabolic requirements of the body
- Etiology:
  1. Ischemic heart diseases (most common)
  2. Systemic hypertension
  3. Valvular disease
  4. Myocardial disease as cardiomyopathy.



# Clinical presentation of HF

Right-Sided Heart Failure	Left-Sided Heart Failure
<b>❶ Backward manifestations</b> : eg: <b>(Systemic congestion)</b> <ul style="list-style-type: none"> <li>- Lower Limb edema</li> <li>- Ascites</li> <li>- Congested neck veins (↑↑ JVP)</li> <li>- Congested liver</li> </ul>	<b>❶ Backward manifestations</b> : eg <b>(Pulmonary congestion)</b> <ul style="list-style-type: none"> <li>- Dyspnea</li> <li>- Cough</li> <li>- Hemoptysis</li> <li>- Recurrent chest infection</li> </ul>
<b>❷ Forward manifestations</b> : eg syncope, fatigue <b>(Low COP manifestations)</b>	<b>❷ Forward manifestations</b> : eg syncope, fatigue <b>(Low COP manifestations)</b>
<b>❸ Cardiovascular manifestations</b> <u>S<sub>3</sub> &amp; gallop</u>	<b>❸ Cardiovascular manifestations</b> <u>S<sub>3</sub> &amp; gallop</u>



## LEFT SIDED ♥ FAILURE

- Paroxysmal Nocturnal Dyspnea
  - Elevated Pulmonary Capillary Wedge Pressure
  - Pulmonary Congestion
    - Cough
    - Crackles
    - Wheezes
    - Blood-Tinged Sputum
    - Tachypnea
  - Orthopnea
  - Tachycardia
  - Exertional Dyspnea
  - Fatigue
  - Cyanosis
  - Endocrine
  - Confusion
- 

## RIGHT SIDED ♥ FAILURE (Cor Pulmonale)

- Fatigue
  - ↑ Peripheral Venous Pressure
  - Ascites
  - Enlarged Liver & Spleen
  - Dependent Edema
  - May be secondary to chronic pulmonary problems
  - Distended Jugular Veins
  - Anorexia & Complaints of GI Distress
  - Weight Gain
- 

Med Surg Nursing - PowerPoint Web site:  
<http://www.cengage.com/medsurgnursing>  
 Downloaded May 6, 2009. Image: "Left and right sided 'Cor Failure'"

# INVESTIGATIONS

1. Chest Xray → cardiac size
2. echocardiography: → assesses ventricular function → ejection fraction (55-70% normal)



# Treatment

## ① Medical:

### ① ↓ Preload:

① Rest: Semi-sitting or sitting

② Diet: Salt restriction (< 2 gm/day)

③ Diuretics

④ VenoDilators: eg Nitrates “Nitroglycerin”

② ↓ Afterload: Arteriodilators: eg Ca channel blocker (CCB) : Diltiazem

③ ↑ Contractility: Inotropic drugs: eg:

1. Digitalis

1. Dobutamine

# Treatment

## ② Interventional:

① ↓ Preload:

Mechanical removal of fluids: By Dialysis

② ↓ Afterload:

Mechanical assistance of circulation: By  
IABP “Intra-Aortic Balloon Pump”

③ ↑ Contractility: CRT (Cardiac Resynchronization Therapy)

## ③ Surgical:

Cardiac transplantation



**Thank you**

Dr\dina monir

**KEEP  
WORKING  
HARD.  
LET THE  
RESULTS  
SPEAK FOR  
THEMSELVES.**