

chemical potential calculation

Jacob Jeffries

February 12, 2024

1 Insertions

We first perform insertions onto a lattice starting for an initial configuration. This initial configuration has type $t(\sigma)$ at each site σ . The solution has atom types $\alpha \in \mathcal{A}$ and the atomic fraction of α in the solution is x_α .

Then, to calculate chemical potentials, we first note that, given a system with free energy G° , we can add an extra atom of type α , yielding a free energy:

$$G_\beta = G^\circ + \mu_\beta \quad (1)$$

So:

$$G_\beta - G_\alpha = \mu_\beta - \mu_\alpha \quad (2)$$

which then measures the free energy difference from swapping an α atom to an β atom. Then, approximating (at 0 K and 0 bar) the free energy as internal energy yields:

$$\langle E_\sigma^{(\beta)} - E_\sigma^{(\alpha)} \rangle = \mu_\beta - \mu_\alpha \quad (3)$$

where $E_\sigma^{(\alpha)}$ is the energy of the lattice with an atom of type α at site σ . Additionally, we can form an Euler equation for the reference configuration:

$$\frac{1}{N} E_\sigma^{(t(\sigma))} = \sum_\alpha \mu_\alpha x_\alpha \quad (4)$$

where N is the number of sites in the lattice. For an alloy with k elements, we can form $\frac{1}{2}k(k-1)$ equations for each α - β pair. So, including the Euler equation below, we can form a system of $\frac{1}{2}k(k-1) + 1$ equations, which forms either a fully determined system for $k = 2$, or an overdetermined system for $k > 2$. Both cases can be solved using least squares.

2 Fitting

We have multiple data points, i.e. a set of chemical potentials for each composition, by creating multiple solutions with specified compositions. Then, we can fit the chemical potentials as a function of concentration:

$$\mu_\alpha = \mu_\alpha^\circ + \sum_{\beta} A_{\alpha\beta}(x_\beta - x_\beta^\circ) \quad (5)$$

where μ_α° and $A_{\alpha\beta}$ are fitting parameters and x_α° is some reference concentration. Then, without loss of generality, we can always set $x_\beta^\circ = 0$, since:

$$\mu_\alpha = \mu_\alpha^\circ - \sum_{\beta} A_{\alpha\beta}x_\beta^\circ + \sum_{\beta} A_{\alpha\beta}x_\beta \quad (6)$$

where $\mu_\alpha^\circ - \sum_{\beta} A_{\alpha\beta}x_\beta^\circ$ is a constant. So, setting $x_\beta^\circ = 0$, we fit:

$$\mu_\alpha = \mu_\alpha^\circ + \sum_{\beta} A_{\alpha\beta}x_\beta \quad (7)$$

This can be achieved with multivariate multiple linear regression. Note that, though, there are only $k - 1$ independent variables since $\sum_{\alpha} x_\alpha = 1$. So, one element γ must be excluded from the fit for unique $A_{\alpha\beta}$'s. This can be achieved by setting any row of A to 0. The function `lm` in `R` automatically excludes columns to enforce linear independence between independent variables.