

## **SAC 301 SOIL FERTILITY , FERTILISERS AND MANURES 2+1**

**Course teacher : Dr. V.V. Krishnamurthi**

“Soil fertility is the ability of the soil to supply essential plant nutrients during growth period of the plants, without toxic concentration of any nutrients”. i.e. **“the capacity of soil to supply nutrient in available form to crop”**.

### **1) Soil productivity**

“Soil productivity is ability of soil to produce a particular crop or sequence of crops under a specified mgt system” i.e the crop producing of soil”.

All the productive soils are fertile but many of the fertile soils are unproductive, because they are subjected to drought or other unsatisfactory growth factors or management practices.

### **Soil Management**

1. Excess elements may be toxic eg. Mo, Cu, Fe.
2. Excess of one element may cause deficiency of other elements.
  - a. Excess of N may cause K deficiency
  - b. Excess of K may cause Mn deficiency
  - c. Excess of P may cause Zn deficiency
  - d. Excess of Ca may cause Fe deficiency

Imbalanced fertigation will not help in increases the yield but it increases only the cost of production. This may lead to **Luxury consumption**.

### **2) LUXURY CONSUMPTION**

“Luxury consumption is defined as the nutrient concentration range in which added nutrient will not increases yield but can increases nutrient concentration”.

### **History of development of soil fertility**

**Francis Bacon** (1591- 1624) suggested that the principle nourishment of plants was water and the main purpose of the soil was to keep plants erect and to protect from heat and cold.

Jan Baptiste **Van Helmant** (1577 – 1644) was reported that water was sole nutrient of plants.

**Rober Boyle** (1627 – 1691) an England scientist confirmed the findings of Van Helmant and proved that plant synthesis salts, spirits and oil etc from H<sub>2</sub>O.

**Anthur young** (1741 – 1820) an English agriculturist conducted pot experiment using Barley as a test crop under sand culture condition. He added charcoal, train oil, poultry dung, spirits of wine, oster shells and numerous other materials and he conduced that some of the materials were produced higher plant growth.

**Priestly** (1800) established the essentiality of O<sub>2</sub> for the plant growth.

J.B. Boussingault (1802-1882) French chemist conducted field experiment and maintained balance sheet. He was first scientist to conduct field experiment. He is considered as **father of field experiments**.

**Justus Van Liebig (1835)** suggested that

- a. Most of the carbon in plants comes from the  $\text{CO}_2$  of the atmosphere.
- b. Hydrogen and  $\text{O}_2$  comes from  $\text{H}_2\text{O}$ .
- c. Alkaline metals are needed for neutralization of acids formed by plants as a result of their metabolic activities.
- d. Phosphorus is necessary for seed formation.
- e. Plant absorb every thing from the soil but excrete from their roots those materials that are not essential.

The field may contain some nutrient in excess, some in optimum and some in least, but the limiting factor for growth is the least available nutrient. **The law of Mn**, stated by **Liebig in 1862**, is a simple but logical guide for predicting crop response to fertilization. This law states that, “the level of plant production cannot be grater than that allowed by the most limiting of the essential plant growth factors”. The contributions made by Liebig to the advancement of agriculture were monumental and he is recognized as the **father of Agricultural chemistry**.

J.B. Lawes and J. H. Gilbert (1843) established **permanent manurial experiment** at Rothemsted Agricultural experiment station at England. They conducted field experiments for twelve years and their findings were

- a. Crop requires both P and K, but the composition of the plant ash is no measure of the amounts of these constituents required by the plant.
- b. No legume crop require N. without this element, no growth will be obtained regardless of the quantities of P and K present. The amount of ammonium contributed by the atmosphere is insufficient for the needs of the crop.
- c. Soil fertility can be maintained for some years by chemical fertilizers.
- d. The beneficial effect of fallow lies in the increases in the available N compounds in the soil.

**Robert Warrington England** showed that the nitrification could be supported by carbon disulphide and chloroform and that it would be stopped by adding a small amount of unsterilized soil. He demonstrated that the reaction was two step phenomenon. First  $\text{NH}_3$  being converted to nitrites and the nitrites to nitrites.

#### **Essentiality of the elements established by**

Carbon	: Priestly (1800)
Nitrogen	: Theodore De saussure (1804)
Ca, Mg, K, S	: Carl sprengel (1839)
Phosphorus	: Von Liebig (1844)
Iron (Fe)	: E. Greiss (1844)
Mn	: J.S. Hargue (1922)
Zn	: Sommer and Lipman (1926)
Cu	: Sommer, Lipman and Mc Kenny (1931)
Mo	: Arnon and Stout (1939)
Na	: Brownell and wood (1957)
Co	: Ahamed and Evans (1959)
B	: Warring ton (1923)
Cl	: Broyer (1954)

### **Essential elements**

“A mineral element is considered to plant growth and development if the element is involved in plant metabolic functions and the plant cannot complete its life cycle without the element”.

### **Major Nutrients or Macro elements**

The plant/ plants require comparatively larger quantity of nutrients for complete the life cycle they are

**C, H, O, N, P, K, Ca, Mg and S.**

### **Minor elements of Micronutrients**

The plants require comparatively small or very trace quantity of nutrients for complete the life cycle.

**Fe, Zn, Cu, Mn, B, Mo, Cl, Na, Va, Si, Co and Ni**

### **Criteria of essentiality – Arnon and stout 1939.**

- a. The plant must be unable to grow normally or complete its life cycle in the absence of the element.
- b. The element is specific and cannot be replaced by other.
- c. The element plays a direct role in the metabolism of the plant.

Until the middle of 19<sup>th</sup> century, the scientist did not have experimental technique to distinguish essential and non essential elements. With the development of **“Solution Culture”** technique the scientist established the essentiality of nutrients.

## **SOIL FERTILITY EVALUATION**

Optimum productivity of any cropping systems depends on an adequate supply of nutrients. Although one or more nutrients are commonly applied to most crops, the quantity of nutrient removed in the harvested crop is much grater than the quantity added.

The proper rate of plant nutrient is determined by knowing the nutrients requirement of the crop supply is power of the soil.

**Diagnostic techniques**, including identification of deficiency symptoms are helpful in determining specific nutrient stress and the quantity of nutrients needed to eliminate the stress.

**Several techniques are commonly employed to asses the fertility status of a soil.**

1. Nutrient deficiency symptoms of a plant.
2. Analysis of tissue from the plants growing in the soil.
3. Biological tests in which the growth of either higher plants or certain micro organism is used as measure of soil fertility.
4. Soil analysis.

### **Nutrient deficiency symptoms**

It can be classified as

- a. Complete crop failure at the seedling stage.
- b. Severe **stunting** of plants.
- c. Specific leaf symptoms appearing at varying times during the season.

- d. Internal abnormalities such as clogged conductive tissues.
- e. Delayed abnormal maturity.
- f. Obvious yield differences.
- g. Poor quality of crops including difference in protein, oil or starch.

### **Hidden hunger**

“It refers to a situation in which a crop needs more of a given nutrient yet has shown no deficiency symptoms”. The nutrient content is above the deficiency symptom zone but still considerably below that needed for optimum crop production.

Hidden Hunger is term used to described a plant that shows no obvious symptoms, yet the nutrient content is not sufficient to give the top profitable yield.

### **Seasonal effects**

Nutrient shortages in the soil may be caused by abnormal weather conditions. Nutrient may be present in sufficient quantities when conditions are ideal, but in drought, excessive moisture. Unusual temperature conditions plant may not be able to obtain an adequate supply.

### **Nutrient uptake is generally reduced by**

1. Mass flow of nutrients is reduced by decreases growth rate and transpiration.
2. Nutrient diffusion rate decreases with declining temperature and lower concentration gradient.
3. Mineralisation of nutrients complexed with organic matter is reduced.

### **Plant analysis**

Two types of plant analysis

1. Tissue test on fresh tissue in the field.
2. Tissue analysis performed in the laboratory.

Plant analysis are based on the premise that the amt of a given nutrient in a plant is directly related to the availability of the nutrient in soil. Since shortage of nutrient will affect the growth others may accumulate in the cell sap regardless of their supply.

### **Tissue test and plant analysis are made for**

1. To aid in determining the nutrient supplying power of the soil.
2. To help identify deficiency symptoms and to determine nutrient shortages before they appear as symptoms.
3. To aid in determining the effect of fertility treatment on the nutrient supply in the plant.
4. To study the relationship between the nutrient status of the plant and crop performance.

### **Tissue test**

The plant root absorbs the nutrient from the soil and these nutrients are transported to other port of the plant where they are needed. The concentration of the nutrients in the cell sap is usually a good indication of how the plant is supply at the time of testing. It is possible to anticipate certain problems of deficiency of nutrients.

### **1. In general the tissue test methods employed are**

1. The intensity of the colour developed is compared with **standards** and used as a measure of the supply of nutrient.
2. In plant sap, the colour developing reagents were added, and the resulting colour intensity is compared with standard chart, that indicates very low, low medium and high nutrient content.

### **2. Plant parts to be tested**

It is essential to test the part of the plant that will give the best indication of the nutritional status. In general, the recently matured leaf is used for testing and the immature leaf in the top of the plants are avoided.

### **3. Time of testing**

In general, most critical stages for tissue testing is a bloom to the early fruiting stage.

### **Total analysis**

Total analysis is performed on the plants or on plant parts. After sampling, the plant material is dried, ground and the nutrient content determined by digesting or ashing of the plant materials. With total analysis, the content of all descents, essential or non essential can be determined.

### **Critical nutrient range**

Critical nutrient range is that range of nutrient concentration at a specified growth stage above which the crop is amply supplied and below which the crop is deficient.

### **Diagnosis and recommendations integrated system (DRIS)**

DRIS is a system that identifies all the nutrimental factors limiting crop production and thus increases the chance of obtaining high crop yields by improving fertilizer recommendations.

### **Index value**

Index value that measure how far particular nutrients in the leaf or plant are from the optimum are used in the calibration of classify yield factors.

### **Biological tests**

It involves in the field test, strip tests and in the laboratory green house tests. The nutrients was assessed by using higher plants and micro organisms.

#### **1. Mitscherlich pot culture method.**

Mitscherlich predicted the plant nutrient reserved (native fertility) as the percentage of increases in the yield expected from the addition of a given quantity of fertilizers.

#### **2. Jenny pot culture tests**

He grouped the plants into 3 categories viz., (a) definite deficiency (b) probable deficiency and uncertain deficiency based on the percentage of yield.

#### **3. Neubauer seedling method**

It is based on the uptake of nutrient by a large no. of plants grown on a small amount of soil. The roots thoroughly penetrate the soil, exhausting the available nutrient supply within a short

time. The total nutrients removed are quantified and tables are established to give the Mn value of macro and micro nutrients available for satisfactory yields of various crops.

#### **4. Sunflower pot culture techniques for boron**

Deficiency is classified as marked deficiency Moderate deficiency and not or little deficiency based upon the no of days after which boron deficiency is observed.

#### **5. A value technique**

Fried and Dean (1952) calculated the avail nutrient in the soil be using Radio active isotopes.

A value is defined as that amt of nutrients in soil, which behaves in a similar way as the applied fertilizers nutrient doses.

$$A = B \left( \frac{1 - y}{y} \right)$$

where,      A      =      Avail soil nutrient  
                 B      =      Amount of fertility nutrient applied.  
                 Y      =      Fraction of the nutrients derived from the fertilizers  
   contained in the plant.

#### **Soil testing**

Analysis of soil is essential for determining the nutrient requirement of a crop and nutrient supplying power of the soil.

#### **Objective of soil testing**

1. To provide an index of nutrient available or supply in a given soil.
2. To predict the probability of obtaining a profitable response to lime and fertilizer.
3. To provide a basis for recommendations on the amount of lime and fertilizer to apply.
4. To evaluate the fertility status of soils on a country soil area, state wise basis by the use of soil test summaries.
5. It helps to predict the amt of nutrients needed to supplement the supply in the soil.

#### **Soil analysis**

Soil sampling includes.

- a. Areas of sample
- b. No. of sites to sample composite
- c. Depth of sampling.
- d. Time of sampling.

#### **Laboratory characteristics**

Determination of  $p^H$ , soluble salts, calcium carbonate, organic carbon, available nutrients were done.

Grouping of soils in different classes based upon the soil test values. The different classes were very low, low medium, high and very high.

## 1) Nitrogen

The cheap source of N is the crop residues in temperate region the soil N occurs at 2.6 tonnes / ha. In acid tropical soils, the N content is 0.03-0.1 %. Rainfall is also source of N at 4.6 kg of N / ha is received  $\text{ha}^{-1} \text{yr}^{-1}$ . It is converted to  $\text{NO}_3$  during lightning addition of organic matter and fertilizer is other major sources.

### Functions of N :

- a] N is an essential constituent of proteins and is present in many other compounds of great physiological importance in plant metabolism
- b] N is an integral part of chlorophyll, which is primary observer of light energy needed for photosynthesis.
- c] N also imparts vigorous vegetative growth and dark green colour to plants.
- d] It produces early green and delay in maturity to plants
- e] It governs the utilization of K, P and other elements

### Deficiency of N:

1. Plants are stunted and yellow in appearance.
2. The loss of protein N from chloroplasts. In older leaves produces the yellowing or chlorosis. It appears first on the lower leaves, the upper leaves remain green, while under severe N deficiency lower leaves will turn brown and die.
3. The necrosis begins at the leaf tip and progress along the midrib until the leaf is dead.
  - In cauliflower - young leaves turn pale yellow and old leaves become orange.
  - In coffee - Veins becomes yellow and new leaves are very small.
  - Tomato - Stem become purple and hard. Flower buds become yellow and flower dropping rate also increases

### C. Excess of Nitrogen (Toxicity off nitrogen)

- a. Causes excess vegetative growth, dark green leaves, lodging, maturity is delayed with increases susceptibility to pest and disease.
- b. In cotton, weak fibre are resulted. In rice, lodging is common.
- c. Lengthening of crop duration and narrow leaf.
- d. Slender shoot, profuse vegetation, thick peel and skin will be rough and leathery in the case of citrus.
- e. Excess N in coffee plant, interferes the K uptake causing imbalance between N and K.

### Forms of soil nitrogen

Inorganic forms

1. Ammonium  $\text{NH}_4^+$
2. Nitrite  $\text{NO}_2^-$
3. Nitrate  $\text{NO}_3^-$
4. Elemental N (No)

Organic forms of soil N

Amide form ( $\text{NH}_2$ )

Plant absorbs N as both  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ .

### Losses of Nitrogen

1. Crop removal
2. Leaching (or) drainage (11-18% loss)
3. Gaseous losses as  $\text{NH}_4$  or elemental  $\text{N}_2$
4. Volatilization.
5. Erosion ( $8 - 15 \text{ kg ha}^{-1}\text{yr}^{-1}$ ).
6. Ammonia fixation by clays
7. Immobilization in organic materials.

### N transformations in soils

- a. N Mineralisation
- b. Aminisation
- c. Conversation of urea
- d. N Immobilization.
- e. N factor
- f. Ammonification.
- g. Nitrification
- h. Denitrification
- i. Organic fixation
- j. Elemental N loss.
- k. Nitrogen cycle.

### Phosphorus

Phosphorus occurs in most plants in concentrations between 0.1 and 0.4%. Plants absorb either  $\text{H}_2\text{PO}_4^-$  or  $\text{H} + \text{PO}_4^{2-}$  ortho  $\text{PO}_4$  ions. Absorption of  $\text{H}_2\text{PO}_4^-$  is greatest at low pH values, where as uptake of  $\text{HPO}_4^{2-}$  is greater at higher values of soil pH, plant uptake of  $\text{HPO}_4^-$  is much slower than  $\text{H}_2\text{PO}_4^-$ .

### Functions of P

1. It has a greater role in energy storage and transfer.
2. It is a constituent of nucleic acid, phytin and phospholipids
3. It is essential for cell division and development
4. P compounds act as energy currency within plants. The most common P energy currency is that found in ADP and ATP. Transfer of the energy rich  $\text{PO}_4$  molecules from ATP to energy requiring substances in the plant is known as “**Phosphorylation**”
5. It stimulates early root development and growth and there by helps to establish seedlings quickly.
6. It gives rapid and vigorous start to plants strengthen's straw and decreases lodging tendency.
7. It is essential for seed formation because larger quantities of P is found in seed and fruit-phytic acid is the principle storage form of phosphorus in seeds.
8. It increases the activity of Rhizobia and increases the formation of root nodules.

### Deficiency symptoms

P is mobile in plants and when a deficiency occurs it is translocated from older tissues to the active meristematic regions.

1. It arrests metabolism resulting in reduction of total N of plants.
2. Reduced sugar content.



3. Premature leaf fall.
4. Develops necrotic area on the leaf petiole and in the fruit
5. Leaves will show characteristic bluish green colour.

### Toxicity of phosphorus

- a. Profuse root growth i.e. lateral and fibrous root lets.
- b. It develops normal growth having green leaf colour.
- c. It may cause in some cases trace elements deficiencies i.e. **Zinc and Iron.**

### Soil P its origin and Nature

Soil P exists in many primary and secondary compounds. The **apatite** group of primary mineral is the original source i.e. 55% of soil P.

#### a) Aluminium and Iron Phosphates

In the initial stages of acid weathering  $\text{PO}_4$  becomes increasingly bound to  $\text{Fe}^{3+}$  and  $\text{Al}^{3+}$  ions released from silicate minerals by replacing  $\text{OH}^-$  from **hydroxyl minerals** or oxygen from **oxide minerals**.  $\text{FePO}_4$  and  $\text{ZnPO}_4$  has been found in  $\text{H}_2\text{O}$  logged or poorly drained soils, sand and silt fraction of some soils.

### Calcium phosphates

The group of compounds form an important category in the young soils and the matured soils of **neutral** to **alkaline** pH ranges. The  $\text{CaPO}_4$  found either in **stable** or **metastable** state in the soil fertilizer reaction zones.

<b>Eg.</b> Fluorapatite	-	$\text{Ca}_{10}(\text{PO}_4)_3 \text{Fe}$
Carbonate apatite	-	$\text{Ca}_{10}(\text{PO}_4)_3 \text{CO}_3$
Dicalcium $\text{PO}_4$	-	$\text{Ca HPO}_4 \cdot 2\text{H}_2\text{O}$
Monocalcium $\text{PO}_4$	-	$\text{CaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ .

### Organic phosphates

It is derived secondarily by the addition of organic matter to the soil through the growth of plants and the deposition of plant residues. The soil micro organic synthesize organic  $\text{PO}_4$  compounds and accumulate in the soil mixed with derived from plant tissues. It constitute 20 – 30% of total soil P. The major classes of organic compounds in soils are

- a. Phospholipids (0.6 – 0.9%)
- b. Phospho proteins of nucleic acid ( 0.6 – 2.4%)
- c. Phosphorylated sugar
- d. Phytin and Inositol  $\text{PO}_4$

The ideal C : N : P : S ratio in soils are

- a. Calcarious            113 : 10 : 1.3 : 1.3
- b. Non calcarious      147 : 10 : 2.5 : 1.4
- c. Indian soils          144 : 10 : 1.4 : 1.8

### Potassium

The potassium ion ( $\text{K}^+$ ) is actively taken up soil solution by plant roots. The concentration of  $\text{K}^+$  in vegetative tissue ranges from 1 to 4% on dry matter basis.

### Functions of potassium

1. Essential for photosynthesis, development of chlorophyll.
2. It improves vigour of the plants to enable to withstand adverse climatic conditions.
3. Reduces lodging in cereal crops.
4. It regulates stomata opening and closing.
5. It regulates the movement of ions within the plants and hence it is called **traffic policeman** of the plant.
6. Activation of enzymes, enzyme synthesis, peptide bonds synthesis.
7. Regulates H<sub>2</sub>O imbalance within the plant.

### Deficiency symptoms

1. Plant becomes stunted in growth with shortening of internodes and bushy in appearance.
2. K deficiency in plants shows reduced rate of photosynthesis.
3. Chlorosis, yellowing of leaves and leaf scorch in case of fruit trees.

**Rice :** The leaf tips will turn dark brown in colour and blades will be blueish green, chlorotic and necrotic are seen.

**Banana :** Deficiency is seen in the margin and bottom of leaves.

**Grapes :** Leaves are yellow with brown spots which are necrotic, brittle with uneven ripening.

### Sources of K


The micas and feldspars constitute the major K bearing minerals which on weathering slowly release K to the soil.

(Muscovite and biotite) (Orthoclase and microcline)

### Forms of potassium in soils

- a. H<sub>2</sub>O soluble K.
- b. Readily exchangeable K.
- c. Fixed K.
- d. Lattice or increased reserve K.

The different forms are in **dynamic equilibrium** with one another and represented as follows.



### H<sub>2</sub>O soluble K

The Concentration of H<sub>2</sub>O soluble K in soil depends on (a) type of clay (b) H<sub>2</sub>O content (c) Intensity of leaching (d) Amount of exchangeable K (e) Kind and concentration of other ions.

The dilution of the soil, increases the concentration of H<sub>2</sub>O soluble K and drying decreases it. The uptake of K is influenced by the presence of other cations, i.e. Ca<sup>2+</sup> and Mg<sup>2+</sup> of Al<sup>3+</sup> (acidic soils) and Na<sup>+</sup> in salt affected soils.

$$\text{The activity ratio (A}^{\text{K}}\text{)} = \frac{\text{Activity of K}^+}{\sqrt{\text{activity of Ca}^{2+} \text{ and Mg}^{2+}}} \quad (\text{or})$$

$$= \frac{a_K}{\sqrt{a_{\text{Ca-Mg}}}}$$

## 2. Exchange K

The K adsorbed on soil clay complex and replaceable with neutral salts in relatively short time. The Cl reacts with soil Ca and form  $\text{CaCl}_2$  which is leached by high rainfall. Hence **K is called as Decalcifier**

## 3. Fixed K

Added K which is firmly bound by the soil and not immediately replaceable with neutral salts.

## 4. Lattice K or Mincreasesal K

Muscovite, biotite and K feldspar. The capacity of soils to release lattice K by weathering depends on the content of K mincreasesals and soil texture.

### Potassium fixation

The important of K fixation is to regulate the supply of the soil a for the plants and protects it against loss through leaching.

In the dynamics of soil Potassium, the phenomenon of fixation of exchange K and the liberation of non exchange K play an important role. K ions are relatively small to enter the silica sheets where they are held firmly by **electrostatic forces**. The presence of  $\text{K}^+$  ions can block the release of fixed  $\text{NH}_4^+$  and vice visa.

### Calcium

Calcium is absorbed by plants as  $\text{Ca}^{2+}$  and its concentration ranges from 0.2 to 1.0% and it is supplied through mass flow method.

### Functions of calcium

1. It is immobile in plants and hence the deficiency is observed in younger leaves.
2. It is a constituent of cell wall and increases in stiffness of plants.
3. Promotes root development and growth of plants, root elongation and cell division.
4. Helps to translocate the sugar in the plants.
5. It involves chromosome stability and that it is a constituent of chromosome structure.
6. Affects translocation of CHO in plants.
7. Encourages seed production.
8. Activates enzyme phosphate and kinease.
9. Accumulated protein during respiration by mitochondria and it increases their protein content.
10. It binds DNA to protein molecules.

### Deficiency of calcium

- a. Young leaves of terminal buds dieback at the tip and margins.
- b. Normal growth in affected.
- c. Root may become short, stubby and brown.
- d. Causes acidity of soil.
- e. Cell may become rigid and brittle.
- f. Young leaves of cereals remain folded.

### Sources of soil calcium

Earth crust contains about 3.64%. The important source of calcium is **anorthite** ( $\text{Ca Al}_2\text{Si}_2\text{O}_7$ ). Generally arid region soils contain high amount of Ca regard less of texture, low rainfall and little leaching.

In arid and semiarid regions    Calcite ( $\text{CaCO}_3$ )  
   Dolomite ( $\text{Ca, Mg, (CO}_3)_2$ )  
   Gypsum ( $\text{CaSO}_4, 2\text{H}_2\text{O}$ )

In humid regions, even the soils formed from lime stone are frequently acid in the surface layers because of the removal of Ca and other cations by leaching.

### Factors affecting $\text{Ca}^{2+}$ availability

1. Total Ca supply : Sandy acidic soils with low CEC having less Ca.
2. Soil pH : In acid soils Ca is not readily available to plants at low cane.
3. CEC : Soils having low CEC might will supply more  $\text{Ca}^{2+}$ .
4. % Ca saturation : High  $\text{Ca}^{2+}$  saturation indicates favourable pH for pt growth and microbial activity.
5. Type of soil colloid : 2 : 1 type require higher Ca saturation than 1 : 1 type.
6. Ratio of  $\text{Ca}^{2+}$  to other catians : Increaseeasing the  $\text{Al}^{3+}$  cane in soil solution reduces Ca uptake in pt.

### Magnesium

Magnesium is absorbed as  $\text{Mg}^{2+}$  and the concentration in crop varies between 0.1 and 0.6%. It was taken by plant by **Mass flow and diffusion**.

### Functions of Mg

1. Primary constituent of chlorophyll
2. Imports dark green colour in leaves.
3. Serves as a structural components in ribosomes and stabilizing the ribosome configuration for protein synthesis.
4. Involves numbers of physiological and biochemical function.
5. Activates phosphorylating enzymes in CHO metabolism.
6. Act as a cofactor for certain enzymes other than  $\text{Po}_4$  transfer enzyme.
7. Increases in the oil content of oil seed crops.
8. Regulates the uptake of other nutrients.

### Deficiency of Magnesium

1. Interveinal chlorosis of the leaf in which only the leave veins remain green.
2. Stiff brittle, twisted leaves, wrinkled and distortion of leaves.
3. cotton –lower leaves may develop a reddish purple finally necrotic (Redding of leaves)
4. In brassica, Chlorosis with interveinal mottling uniformly distributed in older leaves while the other vascular tissues remain green. This condition is called “**Puckering**”.

### Sources of soil Magnesium

It constitutes 1.93% of earth crust

Primarily minceasesals (a) Biotite (b) Dolomite (c) Hornblende (d) olivine  
(e) serpentine.

Secondary minerals (a) Chlorite (b) Illite (c) Montmorillonite.

In arid region substantial amount of Mg present as **Epsomite** ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ )

### Functions of Mg in the soil

1. It occurs predominately as exchange and solution Mg.
2. Coarse textured soil exhibits the greatest potential for Mg deficiencies.
3. Competition between  $\text{NH}_4^+$  and  $\text{Mg}^{2+}$  also lowers the  $\text{Mg}^{2+}$  availability to crops.

### Losses of Mg

It is leached by soils and it depends on the Mg content, rate of weathering, intensity of weathering and uptake of plants.

### Sulphur

Sulphur is absorbed by plant roots as  $\text{SO}_4^{2-}$  ions. Concentration of S in plants range between 0.1 and 0.4%.

### Functions of sulphur

1. Essential for synthesis of sulphur containing amino acids cystine, cysteine and methionine.
2. Essential for synthesis of other metabolites including Co-enzyme A, Biotin, Thiamine of vitamin B and Glutathione.
3. Synthesis of chlorophyll.
4. It is a vital part of ferredoxins i.e Fe – S – protein occurring in the chloroplasts.
5. Responsible for the characteristic smell or odor and taste of mustard, onion and Garlics. (Pungent smell)
6. Enhances the oil formation in crops (Soybean)
7. Increases root growth.
8. Stimulate seed formation.
9. Promote nodule formation – Leguminous species.

### Deficiency of sulphur

- a. Stunted growth pale green to yellow colour.
- b. Immobile in plants and plant symptoms start first at younger leaves.
- c. Poor seed set in rapeseed.
- d. Tea – Tea yellows.

### Sources of sulphur

1. Amount has  $<0.05$  ppm in the form of  $\text{SO}_2$ .
2. Earth crust contains 0.06 to 0.10%.
3. Sulphur bearing minerals Gypsum -  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ .  
Epsomite -  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$   
Mirabilite -  $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ .  
Pyrite -  $\text{FeS}_2$ .
4. Silicate minerals contain  $<0.01\%$  S.
5. Igneous rocks 0.02 to 0.07%  
Sedimentary rocks 0.02 to 0.22%

### Forms of sulphur in soil

Present both organic (90%) and Inorganic forms.  
The inorganic forms are

1. Solution  $\text{SO}_4^{2-}$ .
2. Adsorbed  $\text{SO}_4^{2-}$  Readily available fraction.
3. Insoluble  $\text{SO}_4^{2-}$
4. Reduced inorganic compounds.

### Iron

Fe is absorbed by plants roots as  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$  and chelated forms. Sufficiency range of Fe in plant tissue is 50-250 ppm.

### Functions of Iron

1. It helps in the synthesis of chlorophyll.
2. Structural component of porphyrin molecules like cytochromes, hematin, hemes, ferrichrome and hemoglobin. These substances are involved in oxidation-reduction reactions in respiration and photosynthesis.
3. Structural component of non heme compounds like ferredoxins.
4. Constituent of enzyme systems E.g. cytochrome oxidase, catalase, nitrogenase reaction in plants.
5. Component of flavoprotein like  
 $\text{FMN}$  = Flavin mono nucleotides  
 $\text{FAD}$  = Flavin Adenosine Dinucleotide.

### Deficiency symptoms

1. Deficiency symptoms occur in younger leaves since Fe is immobile element within plant.
2. It occurs in soils of calcareous or alkaline soils and poorly drained  $\text{H}_2\text{O}$  logged soils.
3. Younger leaves develop interveinal chlorosis which progresses rapidly over the entire leaf. Severe cases entire leaf turns yellow colour.

### Sources of iron

Earth crust contains about 5%

- Primary and secondary minerals such as 1. Olivine, 2. Pyrite  
 3. Hematite, 4. Goethite, 5. Magnetite and limestone


### Forms of iron (Fe cycle)

Fe occurs in Four major forms in soil.

1. Primary and secondary minerals Fe
2. Adsorbed Fe
3. Organic Fe and
4. Solution Fe.

### Soil solution Fe

It occurs primarily as  $\text{Fe}(\text{OH})^+_{2-}$ ,  $\text{Fe}^{3+}$  concentration in solution is very low. In well drained, oxidized soils, the solution  $\text{Fe}^{2+}$  concentration is less than that of the dominant  $\text{Fe}^{3+}$  in solution. The pH dependent relationship for  $\text{Fe}^{3+}$  is described as



For every increase in pH,  $\text{Fe}^{3+}$  concentration decreases 1000 fold. Oxidation - reduction reactions, the result of change in  $\text{O}_2$  partial pressure, exert considerable influence on the amount of soluble Fe in the soil solution. The insoluble  $\text{Fe}^{3+}$  form predominates in well drained soils, while levels of soluble  $\text{Fe}^{2+}$  increase significantly when soils become  $\text{H}_2\text{O}$  logged. In general, lowering Redox increases  $\text{Fe}^{2+}$  solubility, 10 fold for each unit decrease in pH.

Over the normal pH range in soils, total solution Fe is not sufficient to meet plant requirements for Fe even in acid soils, where Fe deficiency Occur less frequently than in high pH and calcareous soils.

“Is a term derived from a Greek and meaning “Claw”.

Chelates are soluble organic compounds that bond with metals such as Fe, Zn, Cu and Mn increasing their solubility and their supply to plant roots.

“Natural organic chelates in soils are products of microbial activity and degradation of soil organic matter and plant residues. Root exudates are capable complexing nutrients substantial quantities of organic completed. Fe can be cycled through crop residue and compounds of citric acid and oxalic acids have chelating properties.

## **MANGANESE**

Mn concentration in plant ranges from 20 to 500 ppm

### **Functions of Mn**

1. Helps in chlorophyll formation
2. Involves in photosynthesis, particularly in evolution of O<sub>2</sub>.
3. Involves in oxidation - reduction - process in decarboxylation and hydrolysis reactions.
4. Involves in enzyme systems and various enzyme reactions in the citric acid cycle.
5. It is a substitute for Mg<sup>2+</sup> in many of the phosphorylating and group transfer reactions.

### **Deficiency of Mn**

1. Immobile in plant and def. starts in the younger leaves.
2. Interveinal chlorosis occur
  - Oats - Gray specks / streaks
  - Peas - Marsh spot
  - Sugarbeet - speckled yellow
  - Sugar cane - Pahala blight - midrib pale green and white.
3. Deficiency increases aspartic acid and decreases glutamine
4. Increases respiration
5. Accumulation N compounds mainly as amines.

### **Sources of Mn**

Earth crust contains 1000 ppm. Various oxides and hydroxides.  
Manganite MnO (OH)  
Braunite Mn<sub>2</sub>O<sub>3</sub>

### **Factors affecting Mn availability**

#### **1. Soil pH and carbonates**

Liming to acid soils decreases solution and exch. Mn<sup>2+</sup> due to precipitation Mn<sup>2+</sup> as MnO<sub>2</sub>.

#### **2. Excessive H<sub>2</sub>O and poor aeration**

H<sub>2</sub>O logging will reduce O<sub>2</sub> and lower redox potential will increases soluble Mn<sup>2+</sup>. Poor aeration increases Mn availability.

#### **3. Organic matter**

Increases solution and exchangeable Mn.

#### 4. Climatic factor

Increases soil temp during the growing season improves Mn uptake, because of greater plant growth and root activity.

#### 5. Soil micro org

Deficiency caused by soil organisms oxidizing  $Mn^{2+}$  to  $Mn^{4+}$ .

#### Forms of Mn

- i. Solution  $Mn^{2+}$
- ii. Exchange  $Mn^{2+}$
- iii. Organic bound Mn
- iv. Mn Mineral

#### ZINC

Normal concentration in plant 25 to 150 ppm

Deficiency level is < 20 ppm

Toxic level is > 400 ppm

Zn is present in all parts of the plants. In general root contain more zinc than fruits.

Cereals	: 27.8 ppm
Pulses	: 34.8 ppm
Vegetables	: 28.2 ppm
Fruits	: 36.6 ppm

#### Functions of Zn

- 1. Essential too formation of growth hormones
- 2. Helps in reproduction of certain plants.
- 3. Stabilise ribosomal fractions
- 4. Influence the activity of dehydrogenase enzymes
- 5. Involves in auxin metabolism like tryptophan synthetase.

#### Deficiency of Zn

- 1. Light yellow or white areas bet the veins of leaves particularly older leaves.
- 2. Death of tissue, discolored
- 3. Mal formation of fruits
- 4. Reduced growth hormone production.

Cotton	: White bud (or) little leaf
Citrus	: Mottle leaf
Potato	: Fern leaf
Fruit trees	: Rosette (Upnormal growth)
Paddy	: Khaira
Foliar spray	: 0.2 to 0.5% $ZnSO_4$ .

#### Sources of zinc

Soil	: 10-300 ppm
Igneous rock	: >10 ppm
Sedimentary	: > 95 ppm



**Forms of soil Zn**

1. Solution  $\text{Zn}^{2+}$
2. Adsorbed  $\text{Zn}^{2+}$
3. Organically completed  $\text{Zn}^{2+}$

**Factors affecting Zn availability**

1. Soil pH : Avail of Zn decrease with increase soil pH Zn def occur in neutral and calcareous soil.  
At soil pH, Zn precipitates as insoluble amorphous soil Zn, which reduces  $\text{Zn}^{2+}$  in soil.  
Al and Fe oxides, OM and  $\text{CaCO}_3$  increase with increasing pH.
2. Organic matter
3. Interaction with other nutrients  
 $\text{SO}_4$  and N can increase the Zn uptake in plants.  
Higher the soil pH, poor aeration greater the Zn deficiency

**COPPER**

Copper is absorbed by plants as cupric ion  $\text{Cu}^{2+}$   
Normal concentration in plants 5-20 ppm.

**Functions of copper**

1. Essential for the synthesis of vitamin A
2. Act as a catalyst in respiration
3. Act as a “electron carrier” in enzyme which bring about oxidation reduction reactions in plants.
4. Constituent of chlorophyll.

**Deficiency of copper**

1. Chlorosis, withering and distortion of terminal buds.
2. Dead tissue appears along the tips and edges of leaves.
3. Multiple bud formation in the leaf axil and mal formation of leaves.
4. Guava - cracking of fruits and terminal bud die back.
5. Heavy liming, excessive application of N and P - induces Cu deficiency

**Sources of copper**

1. Igneous rock 10-100 ppm of Cu.
2. Sedimentary rock 4-45 ppm
3. Primary minerals contains
  1. Chalcopyrite
  2. Chalcocite
  3. Bornite
4. Sedimentary minerals  
Oxides, Carbonates, Silicates, Sulphates and Chlorides.

**Forms of soil copper**

1. Soil solution - ionic and completed
2. Cation exchange sites of clay and organic matter
3. Occluded and co-precipitated in soil oxide material.
4. Specific adsorption sites
5. Inorganic matter and living organisms

**Factors affecting Cu availability**

1. Soil texture
2. pH
3. CEE
4. Org matter content
5. Hydrous oxides

**Soil texture**

Sandy soils contain lower amounts of Cu than silt and clay soils.

**pH**

The concentration of Cu in soil solution decreases with increase in pH.

**Interaction with other nutrients**

Application of NPK fertilizer induce Cu deficiency

Increase N supply to crops reduce mobility of Cu in plants

High concentration of Zn, Fe and P in soil solution also can depress Cu absorption by plant roots.

**BORON**

B concentration in mono cotyledons and dicotyledons (20-60 ppm) varies between 6 and 18 ppm.

It is absorbed by plants as undissociated boric acid ( $\text{H}_3\text{BO}_3$ ).

**Functions of boron**

1. New cell development in meristematic tissue.
2. Pollination, fruit / seed set.
3. Translocation of sugars, starches, N and P
4. Synthesis of proteins and amino acids.
5. Nodule formation in legumes
6. Regulation of CHO metabolism.

**Deficiencies of Boron**

1. Since it is immobile, deficiency Symptoms occurs in terminal bud growth.
2. Flowering and fruit development are restricted.
3. Sterility and mal formation of reproductive organs.
4. Thickened and curled leaves.
5. Discoloration, cracking or rotting of fruit, tubers or roots

Apple - Internal cracking.

Break down of internal tissue in root crops given rise to darkened areas referred to Brown heart / black heart : cotton - weeping disease.

**Sources of boron**

1. Non metal among the micronutrient
2. Low concentration in earth crust igneous rocks (<10 ppm)
3. Tourmaline and borosilicate contains B.

**Forms of boron (Boron cycle)**

B exists in 4 forms in soil.

1. Rocks and minerals.

2. Adsorbed on clay surface, Fe and Al oxides combined with O.M.
3. Boric acid ( $\text{H}_3\text{BO}_3$ ) and  $4 \text{ B}(\text{OH})_4^-$  in soil solution.

### **MOLYBDENUM**

Non metal anion absorbed as molybdate ( $\text{MoO}_4$ ). It is weak acid and form complex poly anions such as phosphomolybdate. Plant contains <1 ppm Mo.

#### **Functions of Mo**

1. Essential component of enzyme  $\text{NO}_3$  reductase, which catalyses  $\text{NO}_3^-$  to  $\text{NO}_2^-$ .
2. Structural component of nitrogenase enzyme - involved in N fixation.
3. Essential for absorption and translocation of Fe in plant.

#### **Deficiency of Mo**

1. Inhibits flower formation
  2. Imbalances various Amino Acids in plants.
  3. Reduce activity of symbiotic and non symbiotic N fixation.
- Cauliflower - Whip tail.

#### **Sources of Mo**

Earth crust 2 ppm : and range from 0.2 to 5 ppm.

#### **Forms of Mo**

1. Non exchangeable Mo in primary and secondary minerals.
2. Exchangeable Mo held by Fe and Al oxides.
3. Mo in soil solution
4. Organically bound Mo.

Mo in solution occurs mainly as  $\text{MoO}_4^{=}$ ,  $\text{HMoO}_4^-$ ,  $\text{H}_2\text{MoO}_4^0$ .

#### **Factors affecting Mo availability**

1. Soil pH and liming

Mo availability increases with increasing pH. Liming to correct acidity will increase Mo avail. Mo avail decrease with application of acid forming fertilizer  $(\text{NH}_4)_2\text{SO}_4$ .

#### **2. Reaction with Fe and Al**

Strongly absorbed by Fe and Al oxides.

#### **3. Interaction with other nutrients**

Mg and P enhances Mo absorption by plants.

High level  $\text{SO}_4$  decrease Mo absorption by plants.

Cu and Mn decrease Mo uptake by plants.

$\text{NO}_3^-$  encourages Mo uptake

$\text{NH}_4^+$  reduces Mo uptake

The beneficial effect of  $\text{NO}_3^-$  nutrition is related to the release of  $\text{OH}^-$  ions.

## CHLORINE

Normal concentration in plant is about 0.2-2.0%. Absorbed by plants as  $\text{Cl}^-$  through roots and aerial parts.

### Functions of chloride

1. Essential for biochemical reactions Osmotic cation neutralization reactors.
2. Act as a counter ion during rapid K fluxes.
3. Involves in the evaluation of  $\text{O}_2$  in photosynthesis.
4. Creates disease resistant by increase osmotic pressure in cell sap.

### Deficiency of Cl

1. Partial wilting and loss of turgidity.
2. Necrosis, leaf bronzing and reduction in growth.

### Sources of Cl

- i. Igneous and metamorphic rocks
- ii. Soluble salts such as  $\text{NaCl}$ ,  $\text{CaCl}_2$  and  $\text{MgCl}_2$ .
- iii. Earth crust 0.02-0.05%.

It is mobile with in the plant it can be rapidly recycled through soil systems.

## COBALT

Normal concentration of Co in drymatter - 0.02 - 0.5 ppm.

### Functions of cobalt

1. Essential for M.O. fixing atmosphere. N. It forms vit. B12 during growth and development of symbiotic M.O.
2. Improves growth, transpiration and photosynthesis.
3. Take parts in leg hemoglobin metabolism and inibonucliotide reductase.

### Deficiency

1. Acidic, highly leached, sandy soils with total Co low.
2. Calcareous and peaty soils.

### Factors affecting : Co avail

Soil pH : Avail increase with increase in soil acidity,  $\text{H}_2\text{O}$  logging conditions.

Liming and drainage practices - reduce co avail.

Application  $\text{CoSO}_4$  rectify the deficiency.

## VANADIUM

Low concentration of Vanadium is beneficial for growth of M.O.

Vanadium substitute for Mo in N fixation by Rhizobia.

Involves in Biological - oxidation reduction reactions.

### Functions of vanadium


## NICKEL

Nickel content in pt 0.1 - 1.0 ppm dry matter basis.

Taken by plant as  $\text{Ni}^{2+}$

High levels of Ni may induce Zn or Fe deficiency

1. Ni - metal component of urease that catalyse reaction.



2. Essential for N metabolism.
3. Stimulates nodule weight and yield of crops.

## **SODIUM**

### **Functions**

1. Essential for maintaining turgor and growth of Plants.
2. Helps oxalic acid accumulation in Plants.
3. Helps in stomatal opening and regulate  $\text{NO}_3^-$  reductase activity.

### **Sources of Na**

Earth crust 2.8% soils  
Minerals.

### **Forms of Na in soil**

Solution, exchangeable Na and in silicate minerals.  
In arid and semi arid soils Na exist in silicates, NaCl,  $\text{Na}_2\text{SO}_4$   
Na salts accumulating in poorly drained soils of wild and semi arid regions and causes soil salinity and sodicity.

### **Effect of Na on soil properties**

Dispersing action of  $\text{Na}^+$  on clay and organic matter reduces soil aggregation, permeability to air and  $\text{H}_2\text{O}$ , germination and root growth.

### **Sodium fertilizers**

Sod.  $\text{NO}_3$  : 25% Na  
Rhenonia  $\text{PO}_4$  12% NA

Abundant element. in lithosphere and absorbed as monosilicic acid.  
Cereals and grass contain 0.2 - 2.0%.

### **Functions of silica**

1. Contributes structure of cell wall.
2. Contributes drought tolerance to crops.
3. Regulates photosynthesis.
4. De activates invertase enzyme activity in sugarcane resulting grater sucrose production.  
Larger amounts of Si are accumulated in intracellular deposits known as 'plant opals'.
5. Increased available P.
6. Responsible for plant disease resistance.

### **Sources of silicon**

Earth crust : 27.6% Most abundant element in earth crust.  
Soils 23-35%  
Primary and secondary mineral and quartz - major source.  
Quartz is the most common mineral in soil, comprising 90-95% of all sand and silt tractions.

**Factors affecting Si availability**

1. High H<sub>2</sub>O content encourages Si uptake
2. Heavy application of N decreases Si concentration
3. Liming decreases Si uptake in plants.
4. Acidification increases Si uptake
5. Fe and Al oxides, influence Uptake of Si by plants.

**Si fertilizers**

1. Calcium silicate slag ( $\text{CaAl}_2\text{Si}_2\text{O}_4$ ) - 18.21% Si
2. Calcium silicate ( $\text{CaSiO}_3$ ) - 31% Si.
3. Sodium metasilicate  $\text{NaSiO}_3$  - 23% Si.

**Terminology****Straight fertilizer**

Synthetically manufactured inorganic fertilizer containing one major plant nutrient exemption urea.

Eg.  $(\text{NH}_4)_2\text{SO}_4$ ,  $\text{NH}_4\text{NO}_3$ , super, KCl,  $\text{K}_2\text{SO}_4$

Synthetically manufactured inorganic fertilizer containing two or more plant major nutrients. Each granule contain specified proportion.

Eg. Ammonphos, DAP, MAP,  $\text{KN0}_3$ .

### **Mixed fertilizer**

One with contains two or more plant nutrients obtained by **manual** mixing of two or **more straight** fertilizer. Each granule will not contain specified proportion. Eg. Cotton mix, sugarcane mix, groundnut mix.

### **Complete Fertilizer**

One which would supply all the 3 major plant nutrient element in required. proportions.

### **Complex Fertilizer**

One which usually carries two or more of major nutrient and it is obtained by **mechanical blending** of 2 or more straight fertilizer. Each granule may contain specified proportion of nutrient.

### **Fertilizer Grade**

Refer to Minimum guarantee of the plant nutrient in terms of total N,  $\text{P}_2\text{O}_5$ ,  $\text{K}_2\text{O}$  12: 6: 6.

### **Fertilizer ratio**

Refers to relative % on N :  $\text{P}_2\text{O}_5$  ;  $\text{K}_2\text{O}$  2: 1:1.

### **Fertilizer formula**

Refers to expression of quantity and analysis of the material in a mixed fertilizer

### **Acid forming fertilizer**

Acid forming fertilizer is one capable of increasing acidity of the soil which is derived principally from the nitrification of ammonium salts by soil bacteria.

### **Base Fertilizer**

Base fertilizers is one capable decreasing the acidity of the soil

### **Neutral fertilizer**

Neutral fertilizer is one that is guranted to leave neither an acidic nor basic residue in the soil.

### **Clear liquid fertilizer**

Clear liquid fertilizer is one in which the NPK and other materials are completely dissolved.

Production of small quantity of Nitrogenous fertilizer commenced in India (1930)  $(\text{NH}_4)_2\text{SO}_4$  as a by Product from Coke oven gas of the steel plants of Bengal and Bihar.

<b>MFCL</b>	Production of synthetic ammonia was started in India by Mysore chemicals and fertilizers limited near Mysore (1941). Using electrolysis of H <sub>2</sub> O.
<b>FACT</b>	Fertilizers and chemicals travancore Ltd. (FACT) at Alwaye using wood as Raw material for the Production of ammonia.
<b>HCFL</b>	Hindustan chemicals and fertilizers Ltd. (HCFL) was floated to manage Sindri and Nangal plants. In 1951- 61 two more fertilizer factories were establish one at Nangal – Govt. of India.
<b>NCJM</b>	Another at Varanasi - New central Jute mills (NCJM) now Hari fertilizers.
<b>FCI</b>	Sindari and Nangal plants were taken by Govt. of India by fertilizer corporation of India during 1961. (FCO) Between 1961 – 71, there was a considerable growth in fertilizer Industry in the country. Five New plants in the private sector.
<b>EID</b>	Namely EID –party at Ennore – Madras.
<b>East India Distilleries</b>	
<b>GSFC</b>	Gujarat State Fertilizer Co ltd. (GSFC) – Baroda.
<b>CFL</b>	Coramandal Fertilizer ltd. (CFL) – Visakapatnam.
<b>IEL</b>	India explosives Ltd. (IEL) – Kanpur
<b>SCI</b>	Shriram Chemical Industries (SCi) – Kota. The <b>Coal based</b> plant at <b>Talcher</b> and Ramagungam started production of Urea in November 1980.
<b>TACF</b>	During 1982- 83 Tuticorin Alkali chemicals and fertilizers were commenced.

### Phosphatic fertilizers

In India the first unit being EID – Parry – Ranipet in 1906.

### Consumption and production of fertilizers during the period. 1971 – 72 to 1983 – 84

Year	Nutrient	Consumption Jan. – Feb.	Production Apr. – Mar.	Gap. 1000 tonns
1980 – 81	N	3678	2224	1274
	P	1151	763	388
	K	606	-	606
				<b>2510</b>
1981 – 82	N	4069	3143	926
	P	1322	950	372
	K	676	-	676
				<b>1974</b>
1982 – 83	N	4224	3430	794
	P	1436	984	452
	K	727	--	727
				<b>1973</b>
1983 - 84	N	5336	3485	1751
	P	1757	1055	702
	K	799	-	799
				<b>3252</b>

### Technology / Production of Fertilizer Products Ammonia



It is an intermediate product in the MFE of nitrogenous fertilizers. Besides ammonia finds application in the production of nitric acid, soda ash, cleaning agents. Leather tanning petroleum refining, pulp and paper industry, textiles, refrigeration rubber and synthetic resin industries, explosives, food and beverage industries.

### Raw materials

(a) Atmospheric air (b) H<sub>2</sub>O, Natural gas, naphtha fuel oil or heavy petroleum fraction, coal and coke oven gas.

### Methods of Manufacture


Ammonia is a compound of N and H in the ratio of 3:1. It involves 4 steps namely.

(1) Gasification (2) Conversion of Co (3) Gas Purification and (4) Synthesis

Feed stocks for synthesis of Gas production		
	Feed stock	Process
1.	Water	Electrolyzing
2.	Natural gas	Steam refining / Partial oxidation
3.	Naphtha	Steam refining / Partial oxidation
4.	Fuel Oil	Partial oxidation
5.	Coal	Partial oxidation
6.	Coke oven gas	Partial oxidation / reforming low temperature separation.


### Electrolysis

Chemical components of H<sub>2</sub>O is Hydrogen and oxygen. Hydrogen is mixed with required nitrogen from the air and mixed. MFL Nangal has a 300 tpd ammonia plant based electrolysis.



### NH<sub>3</sub> synthesis

The purified mixture containing N<sub>2</sub> and H<sub>2</sub> in the ratio of 3:1 is reacted at a elevated temperature and pressure of the order of 500° C and 270 – 350 atm. Over a activated **iron oxide** catalyst which activates the reaction.



### Schematic reaction of manufacture of NH<sub>3</sub> from fuel oil

Air fractionating

		Fuel oil	
		Low sulphur Heavy stock	
		Heavy sulphur Heavy stock	
N <sub>2</sub>	Gasification chamber	carbon – eliminated out	
Flash gases Comes out	N <sub>2</sub> is added	Desulphurification champer	Unwanted gases are eliminated Eg.HCN elemental S is taken out.

Carbon monoxide      Unwanted gases are eliminated  $\text{CO}_2$  –  
 Conversion chamber      98.8% pure form can be utilized for the  
 manufacture.

Flash gases       $\text{N}_2$  is added      Decarbonation chamber

Liquid N wash      Fuel gas are also eliminated out.

Compressor

$\text{NH}_3$  synthesis      Hydrated liquid  $\text{NH}_3$ .

### **Proportion of ammonia**

Colour less, 99.5% purity containing 82% N sp. gr. 0.617 at  $15.6^\circ\text{C}$ , Boil point  $33.4^\circ\text{C}$

### **Specification for anhydrous ammonia**

- ammonia % by mass Minimum 99.0
- Moisture and residue on evaporation % by mass 1.0
- Oil content - Nil

### **MFE of Nitric Acid**

The raw materials required are  $\text{NH}_3$  and air





$\text{HNO}_3$  is produced in      NFL      -      Nangal  
    HSL      -      Rourala  
    RCF      -      Trombay

### **$(\text{NH}_4)_2\text{SO}_4$**

It is produced by (a) Recovery from coke ovens (b) Direct neutralization process (c) Gypsum process.

### **(a) Recovery from coke ovens**

Ammonia gas is absorbed in H<sub>2</sub>O and then converted into ammonium carbonate by absorbing CO<sub>2</sub>. NH<sub>4</sub> CO<sub>3</sub> is reacted with gypsum to produce Ammonium So<sub>4</sub> and calcium CO<sub>3</sub>.



(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> is evaporated, crystallised, centrifuged and dried.

### Properties

White crystalline compound having 21% N. Sp. gravity : 1.769 and Bulk density : 720 – 1040 kgm<sup>3</sup>.

### Specification given by FCO

Moisture, % wt. max = 1.0



NH<sub>4</sub> Cal N% by wt min = 20.6

Frequently % by wt. min = 0.025

Application of 100 kg of (NH<sub>4</sub>) produces acidity which requires 110 kg of CaCO<sub>3</sub> by neutralized.

### NH<sub>4</sub>NO<sub>3</sub> – 35% N

It is produce by (a) Prilling process (b) Crystallisation process. Ammonia vapour is reacted with acquires HNO<sub>3</sub> in a stainless steel reactor by agitation.



### Calcium NH<sub>4</sub>NO<sub>3</sub>

Produced by granulating concentration NH<sub>4</sub>NO<sub>3</sub> solution with limestone or dolomite in a granulator.

### FCO Specification


Moisture, % by wt. max = 1.0

Total N % by wt. min 25 or 26.0



NH<sub>4</sub> Cal N% by wt min = 12.5 or 13.0

Cal NO<sub>3</sub> % by wt. max = 0.5 or 0.5

### NH<sub>4</sub>Cl 26% N



Produced by reacting liquid NH<sub>3</sub> with hydrochloric acid (or) modified solvery process 2NH<sub>3</sub> + H<sub>2</sub>O + CO<sub>2</sub> → (NH<sub>4</sub>) CO<sub>3</sub>



### Sodium Nitrate

It is obtained as a natural product in chile and manufacture from synthetic ammonia. The important impurities are common salt (Nacl), Glaubers salt (Na<sub>2</sub>SO<sub>4</sub>), Gypsum, Iodates, Borates.

In the process of MFE of  $\text{NaNO}_3$  synthetically. Gaseous nitrogen obtained from coal gas is combined with it to form  $\text{NH}_3$ . Thus formed ammonia is oxidized to nitrous oxides in the presence of catalyst. The nitrous oxides are then observed in a solution of  $\text{Na}_2\text{CO}_3$ . The resulting solution is  $\text{NaNO}_3$ . It is evaporated and the solid material is crystallized.

#### **General characteristics**

1. Readily soluble in  $\text{H}_2\text{O}$ .
2. It is good for top and side dressing during the growth period.
3. It contains 27% sodium, it has a favourable influence of potassium economy.
4. Field crops benefited by Na application are sugar feet, cotton.
5. It may damage the structure of the soil by reducing flocculation.
6. It is a basic fertilizer 100 kg of  $\text{NaNO}_3$  reduces the acidity equivalent to 29 Kg of  $\text{CaCO}_3$ .
7. Besides N, Natural  $\text{NaNO}_3$  contains traces of micro nutrient such as Mg, Mn, B, Cu, Zn.

#### **Other nitrogenous fertilizers**

a. Calcium cyanamide $\text{CaCN}_2$	=	2% N
b. Anhydrous ammonia	=	82% N
c. Ammonia solution	=	20 – 25% N
d. Ammonium chloride and $\text{CaCO}_3$	=	15% N
e. Ammonium Nitrate – Ammonium solution	=	37 – 4% N.
f. Calcium Nitrate	=	15.5%N
g. Calcium Nitrate Urea	=	34% N
i. Urea formalde hyde	=	38%N
j. Oxamide	=	34% N

## Complex fertilisers

Are those which usually carries two and more of major plant nutrients and it is obtained by **mechanical** blending of 2 or more straight fertilizer it self.

Each granule may contain specified proportion of major nutrient elements.

### Ammonium sulphate nitrate 26% N.

It is a double salt of Ammonium  $\text{SO}_4$  and Ammonium  $\text{NO}_3$ . It is also known as **Leuna Salt petre** or **Sulpho nitrate** in Europe. It is a combination of 62.5%  $(\text{NH}_4)_2 \text{SO}_4$  and 37.5%  $\text{NH}_3\text{NO}_3$  and has a nitrogen content of 26%.

### Specifications of ASN

Moisture % by wt. maximum	=	1.0
Total nitrogen % by wt. minimum	=	26.0
$\text{NH}_4$ Cal N% by wt. minimum	=	19.25
Free acidity % by wt. maximum	=	0.15

It is acidic in nature. Application of 100 kg of ASN makes the soil acidity requires 93kg of  $\text{CaCO}_3$  to neutralise

### 2. Ammonium Phosphate sulphate : $\text{NH}_3\text{N} - 16$ , Total P = 20, $\text{H}_2\text{O P} = 19.5$

It contains mainly  $(\text{NH}_4)_2 \text{SO}_4$  (60%) and ammonium  $\text{PO}_4$ (40%) with a nitrogen content of 16% and  $\text{P}_2\text{O}_5$  content of 20% in the 16: 20 : 0 grade, 20: 20 : 0 grade – some urea is added to increases the nitrogen content.

### Raw materials

$\text{NH}_3$ ,  $\text{H}_3 \text{PO}_4$  acid and  $\text{H}_2\text{SO}_4$  acid. It is prepared by **Direct Neutralisation Process**  
 $\text{H}_2\text{SO}_4 + \text{H}_3 \text{PO}_4 + \text{NH}_3 \rightarrow (\text{NH}_4)_2 \text{PO}_4 \text{SO}_4$

### FCO specification

	16: 20 : 0	20: 20 : 0
Moisture max.	1	1
$\text{NH}_4$ Cal N Min	16	18
Total $\text{PO}_4$ min.	20	20
$\text{H}_2\text{O}$ soluble $\text{P}_2\text{O}_5$ min.	19.5	17

### Ammonium Phosphates

**Mono Ammoniums Phosphate  $_4$ (MAP)  $\text{P}_2\text{O}_5 = 55\%$ , Nitrogen = 11 – 12%**

### Raw materials :

$\text{H}_3\text{PO}_4$  acid and  $\text{NH}_3$ . vapourized  $\text{NH}_3$  is injected into a reactor containing wet process  $\text{H}_3\text{PO}_4$  acid of 45 – 52% concentration. The  $\text{NH}_3$  :  $\text{H}_3\text{PO}_4$  ratio is maintained at 1:1.

	$\text{NH}_3 + \text{H}_3\text{PO}_4 \rightarrow \text{NH}_4 \text{H}_2 \text{PO}_4$	
<b>Process:</b>	White crystal	
	pH of 10% solution =	3.5 – 4.0
	Bulk density kg per Cupic metre calcium	= 348.98
	Free flowing. Non hygroscopic material	

$(\text{NH}_4)_2\text{H}_2\text{PO}_4$  **Diammanium Phosphate DAP 18 :46 : 0.** High analysis fertilizer containing 18% by weight of  $\text{NH}_4 - \text{N}$  and 46% by weight of  $\text{P}_2\text{O}_5$  ( $\text{H}_2\text{O}$  soluble  $\text{P}_2\text{O}_5$ ).


Five units mfe DAP.

GSFC	-	Baroda
FACT	-	Cochin II
MFL	-	Manali
IFFCO	-	Kandla
Spic	-	Tuticorin

### Raw materials

1.  $\text{H}_3\text{PO}_4$  acid of 40 – 54%  $\text{P}_2\text{O}_5$
2.  $\text{NH}_3$

It is produced by reacting 2 moles of ammonia and one mole of  $\text{H}_3\text{PO}_4$ .



### FCO Specification

Nitrogen	%	=	18
$\text{P}_2\text{O}_5$	%	=	46
Bulk density Kg Per Calcium max		=	993
pH of 10% solution		=	7.8
Moist % by weight max		=	1
Total N by % weight min.		=	18
Total $\text{P}_2\text{O}_5$ by % weight min.		=	46
$\text{H}_2\text{O}$ soluble $\text{P}_2\text{O}_5$ by % weight min		=	41.

### Ammonium phosphate sulphate

(Ammophos ) 20 : 20: 0. It is MFE by treating the mix of  $\text{H}_3\text{PO}_4$  and  $\text{H}_3\text{PO}_4$  and  $\text{H}_2\text{SO}_4$  with  $\text{NH}_3$ . It contains principally a mixture of amount  $\text{PO}_4$  and Ammonium  $\text{SO}_4$ . FACT manufactures this fertilizer.

### Ammonium phosphate Nitrate **26 :26: 0**

It is prepared by wet process of  $\text{H}_3\text{PO}_4$  8:16: 0  
 (54%  $\text{P}_2\text{O}_5$ ) and  $\text{HNO}_3$  (60%) is neutralised 15 : 15: 0  
 with  $\text{NH}_3$  to produce APN.

### Urea ammonium Phosphates (UAP)

MFE by using urea,  $\text{NH}_4$  and  $\text{H}_3\text{PO}_4$  are called UAP urea and  $\text{H}_3\text{PO}_4$  in the required proportions are reacted in the pre neutralizes. The resulting ammonium  $\text{PO}_4$  slurry is pumped into the granulator. N content is increased by adding more  $\text{NH}_3$  and solid urea.

	24 : 24	or	28: 28
Total N	24		28
$\text{NH}_4 \text{ N}$	7.5		9
Citrate solution $\text{P}_2\text{O}_5$	24		28
$\text{H}_2\text{O}$ Solution $\text{P}_2\text{O}_5$	20.4		25.2

Ammonium sulphate is manufactured by

1. Gypsum process.
2. By product of coal and steel industries

### 1. Gypsum process

The raw materials required are (1) Ammonia (2) Pulverised  $\text{CaSO}_4$   
(3)  $\text{CO}_2$  (4)  $\text{H}_2\text{O}$

The process consists of

1. Ammonia by the synthesis of nitrogen and Hydrogen.
2. Ammonia  $\text{CO}_3$  by reacting Ammonia with  $\text{CO}_2$ .
3. Gypsum is reacted with a solution of  $\text{NH}_4 \text{CO}_3$  to produce  $(\text{NH}_4)_2 \text{SO}_4$ .

### Reactions

1.  $3 \text{H}_2 + \text{N}_2 \rightarrow 2 \text{NH}_3$
2.  $2\text{NH}_3 + 2 \text{H}_2\text{O} + \text{CO}_2 \rightarrow (\text{NH}_4)_2 \text{CO}_3 + \text{H}_2\text{O}$
3.  $(\text{NH}_4)_2 \text{CO}_3 + \text{CaSO}_4 \rightarrow (\text{NH}_4)_2 \text{SO}_4 + \text{CaCO}_3$ .

By this process  $(\text{NH}_4)_2 \text{SO}_4$  is MFE in India on a large scale at Sindri (Bihar) and Always (Kerala) the production of 1 ton of  $\text{NH}_4\text{SO}_4$  and required 1.7 four of  $\text{CaSO}_4$ , 0.6 tone of Coke.

### 2. By product of coal and steel industries

- a.  $\text{Coal} \rightarrow \text{NH}_3 \text{ (gas)} \rightarrow \text{with } \text{H}_2\text{O} \rightarrow (\text{NH}_4)_2 \text{CO}_3$ .  
(destructive + distillation)  $\text{CO}_2$  (wash  $\text{H}_2\text{O}$ )
- b.  $(\text{NH}_4)_2 \text{CO}_3 + \text{Ca}(\text{OH})_2 \rightarrow 2\text{NH}_3 + 2\text{H}_2\text{O} + \text{CaCO}_3$   
(or) Wash  $\text{H}_2\text{O}$  (Slaked lime)
- c.  $2\text{NH}_3 + \text{H}_2 \text{SO}_4 \rightarrow (\text{NH}_4)_2 \text{SO}_4$

Coal contains, on an average 1% N. When coal is distilled in closed evens or reports. A portion of the nitrogen is volatilized as ammonia and ammonium salt retained the 'Wash water' Slaked lime is added to this to called ammonia liquid which is distilled, the ammonia produced in absorbed in  $\text{H}_2\text{SO}_4$ . Ammonia and  $\text{H}_2\text{SO}_4$  combine to give  $(\text{NH}_4)_2\text{SO}_4$ .

It is quick acting fertilizer. It is absorbed in the soil colloids, clay and humus, replacing calcium. Resistant to leaching  $(\text{NH}_4)_2 \text{SO}_4$  Nitrifying  $\text{NO}_3$

Bacteria  $\longrightarrow$

### Urea ( $\text{Co}(\text{NH}_2)_2$ )

This organic fertilizer is cheaper per kg of nitrogen than any other solid nitrogenous fertilizers sold in India. Urea is manufactured by reacting an hydrous ammonia and  $\text{CO}_2$  gas under very high pressure in the presence of a suitable catalyst.


1.  $2\text{NH}_3 + \text{CO}_2$                        $\text{NH}_2 \text{COO NH}_4$   
(Ammonia)                                  an unstable inter mediate product.
2.  $\text{NH}_2 \text{COO NH}_4$                        $\text{CO}(\text{NH}_2)_2 + \text{H}_2\text{O}$   
urea

Urea is a white crystalline salt containing 44 to 46% Nitrogen. It readily absorbs moisture from air.

Specifications of urea as prescribed in the FCO are

- |                                 |   |      |
|---------------------------------|---|------|
| a. Moisture % by weight maximum | = | 1.0  |
| b. Total N % by weight Mn       | = | 46.0 |
| c. Biuret % by weight maximum   | = | 1.5% |

Urea is completely soluble in the soil solution, but it is not utilized in amide form by most of the field crops. Under field condition, it undergoes hydrolysis in the presence of **enzymes** by micro organisms and is converted into Ammonia carbonate and then microbial oxidation into ammonia and nitrate, and in these forms is absorbed by plants.



The conversion of urea into ammoniacal and nitrate forms is complete in about a week.

Urea is less **acidic** compared to  $(\text{NH}_4)_2\text{SO}_4$  since application of 100 kg of urea leaves acidity which requires 80 Kg of  $\text{CaCO}_3$  to neutralize.

### Sodium Nitrate ( $\text{NaNO}_3$ )

It is obtained as a natural product in Chile and MFE from synthetic ammonia. The important impurities are common salt ( $\text{NaCl}$ ), Glauber's salt ( $\text{Na}_2\text{SO}_4$ ), Gypsum, Iodates, Borates.

In the process MFE  $\text{NaNO}_3$  synthetically, Gaseous nitrogen obtained from coal gas is combined with hydrogen to form ammonia. Thus formed ammonia is oxidized to nitrogen oxides in the presence of a suitable catalyst. The nitrogen oxides are then absorbed in a solution of  $\text{Na}_2\text{CO}_3$ . The resulting solution of  $\text{NaNO}_3$  is evaporated and the solid material is crystallized.


### General characteristics

1. It is readily soluble in  $\text{H}_2\text{O}$ .
2. It is good for top and side dressing during the growth period.
3. It contains 27% Na. Na has a favourable influence on potassium economy and other phases of plant nutrition.
4. The field crops benefited by Na application are **sugar beet** and **cotton**.
5. It may damage the structure of the soil by reducing flocculation.
6. It is a basic fertilizer 100 kg of  $\text{NaNO}_3$  reduces the acidity equivalent to 29 kg of  $\text{CaCO}_3$ .
7. Besides Nitrogen, Natural  $\text{NaNO}_3$  contains traces of micro nutrients such as Mg, Mn, P, Cu and Zn.



### **NH<sub>4</sub>Cl Ammonium chloride**

It is produced by treating NaCl with NH<sub>3</sub> and CO<sub>2</sub> to form NaHCO<sub>3</sub> and NH<sub>4</sub>Cl. The resulting NHCl is then separated from the solution.



1. It contains 26% N.
2. It is resistant to reacting loss.
3. It is more acidic compare to (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 100 kg of Nh<sub>4</sub>Cl produces acidity which requires 128 kg of CaCO<sub>3</sub> of neutralise.
4. It contains Chlorine, as such it acts as decalcifying fertilizer.

### **Calcium Ammonium Nitrate (CAN)**

H<sub>2</sub>O is electrolysed into **hydrogen and O<sub>2</sub>**. Air is liquefied into by **Nitrogen and Oxygen**. The nitrogen and hydrogen are then composed and passed over Ammonia catalyst. A part of the Nh<sub>4</sub> is oxidized with oxygen enriched air in the nitric acid plant and absorbed in H<sub>2</sub>O to give HNO<sub>3</sub>.

The NH<sub>3</sub> is reacted with the HNO<sub>3</sub> to give NH<sub>4</sub> NO<sub>3</sub>. liquor. It is mixed with limestone to form **Nitro limestone** or CAN graceless. It is dried, cooled and coated with suitable coating material to give the final fertilizer material.

### **Specification by FCO**

1. Moist % by weight maximum : 1.0
2. Total Nitrogen % by weight maximum : 25.0
3. Ammonium % by weight maximum : 12.5

CaCO<sub>3</sub> combined with Ammonium improves the effeteness of the fertilizer.

1. The explosiveness and hygroscopicity will be reduced.
2. It is neutral in reaction.
3. It forms bigger granules capability easy application.
4. It contains 8.1% calcium (CAN).

### **Ammonium sulphate Nitrate (ASN)**

It is a double Salt of ammonium SO<sub>4</sub> and ammonium NO<sub>3</sub>. It is also known as **Leuna** salt petre or sulphonitrate in Europe. It is a combination of 62.5% ammonium SO<sub>4</sub> and 37.5% ammonium NO<sub>3</sub> and has a nitrogen content of 26%.


Specifications of ASN are prescribed FCO 1957 are

- |  |         |
|--|---------|
| 1. Moist % by weight, maximum                              | = 1.0   |
| 2. Total ammonia and nitrate nitrogen% by weight, Mn       | = 26.0  |
| 3. Amount N% by weight Mn                                  | = 19.25 |
| 4. Free acidity (as NHo <sub>3</sub> ) % by weight maximum | = 0.15  |

It is **acidic** in nature. Application of **100 kg of** ASN makes the soil acidity requires 93 kg ofCaCO<sub>3</sub> to neutralize.

### Ammonium Nitrate (NH<sub>4</sub> NO<sub>3</sub>)

It is MFE by passing NH<sub>3</sub> and oxygen gas with air over a platinum catalyst to form Nitric acid. This acid is neutralized by ammonia to form Ammonium Nitrate



It is highly hygroscopic and shows a marked tendency to absorb moisture from air to cake.

**The following Recommendations have made to minimise the explosive hazards of the salt.**

1. Do not store Ammonium NO<sub>3</sub> any longer than is absolute necessary.
2. Keep all the bags tightly sealed until ready to use.
3. Keep all the bags indoors in a dry place.
4. Pile the bags not higher than 6 bags.
5. Change position of the bags once in a week.

Ammonium NO<sub>3</sub> is 75% and NO<sub>3</sub>- N 25%.

It is readily soluble in H<sub>2</sub>O. NO<sub>3</sub> N is readily available to plants for rapid growth and NH<sub>4</sub> – N. resists leaching losses.

**It is acidic in their residual effect. Application of 100 kgs of ammonium NO<sub>3</sub> produces acidity which requires 60 kg of CaCO<sub>3</sub> to neutralize.**

### Amide Fertilizer

The amide form of nitrogen for fertilizer use is available as urea (CO(NH<sub>2</sub>)

Calcium cyanamide (CaCN<sub>2</sub>)

These materials are carbon compounds and are technically classified as organic compounds.

These are readily soluble in H<sub>2</sub>O and easily decomposed by micro organism in the soil. They are quickly changed into ammonia cal nitrogen and then to NO<sub>3</sub> form.

### Manufacturing of Ammonium sulphate

It contains 20 – 21% N. It is crystalline salt, soluble and soluble in H<sub>2</sub>O.



Specification of (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub> as prescribed by FCO 1957 are

1. Moisture content by weight maximum = 4.0
2. Ammonium Nitrogen % by weight Min = 20.6



### Nitrophosphates

**Raw materials** : Rock PO<sub>4</sub>, HNO<sub>3</sub> and Ammonia

It is produced by acidulation of Rock PO<sub>4</sub> with HNO<sub>3</sub> acid in a series of reactions. This reactant mass contains CaNO<sub>3</sub> and H<sub>3</sub>PO<sub>4</sub>. The mixture can be converted into solid granulated fertilizers by 4 methods.



### Sulphate recycle process



<b><u>FCO specification</u></b>	<b><u>20 : 20: 0</u></b>	<b><u>15 : 15 : 15</u></b>
Moist by weight max	1.5	1.5
Total N min	20	15
Citrate soluble min	20	15
H <sub>2</sub> O Soluble P <sub>2</sub> O <sub>5</sub> min	5.4	4.0

**The following recommendations have to minimize explosive hazards of the salt.**

- Do not store NH<sub>4</sub>NO<sub>3</sub> for longer period.
- Keep all the bags tidely sealed until ready to use.
- Keep all the bags indoor in a dry place.
- Pile the bags not higher than 6 bags.
- Change position of the bags once in a week.

NH<sub>4</sub> NO<sub>3</sub> on oxidation produces 100 kg acidity equivalent 6 kg of Ca CO<sub>3</sub>

### Slow release fertilizers

#### I. Nitrogen inhibitor

#### II. Chemically modified fertilizer

#### III. Coated fertilizers.

### I. Nitrogen inhibitor.

“The materials that are used to inhibit the activities of micro organism that are responsible for denitrification process”.

NH<sub>4</sub>Cal and amide form of N gets oxidized to NO<sub>3</sub> form by biochemical oxidation. The NO<sub>3</sub> can get denitrified into N<sub>2</sub> and N<sub>2</sub>O due to low redox potential and get lost into atmosphere.

One of the measures taken, therefore, to avoid denitrification is to control the production of NO<sub>3</sub>'s. This can be achieved by using nitrogen inhibitors. **Some of the inhibitors are**

#### a. N serve – 2chloro 6 pyridine

It is toxic to nitrosomonas which oxidizes NH<sub>4</sub> ion to nitrate. It is applied @ 0.2 – 2% of N (0.5 Kg/ha) effective for 22 days only.

#### b. 2 Amino – 4chloro – 6 methyl pyridine (AM)

It is toxic to nitrosomonas bacteria when applied @ 4 – 10 ppm by mixing with the fertilizer. It is also used as coating material for solid fertilizers using oil as a binder. The effect is only **40 days**.

#### c. Dicyano Diamide

It is applied @ 5 – 24% on N basis to inhibit nitrification. In winter its effectiveness may go upto 5 months. Higher conc is toxic to plants.

#### d. Thiourca

It inhibits the nitrification by lengthening the lag period prior to exponential growth of bacteria.

## II. Chemically modified fertilizers

The nature of the fertilizer is changed by treating with certain chemicals to reduce its solubility.

- a. Urea formaldehyde (38 – 42% N)
- $$\begin{array}{ccc} & \text{NH}_2 & \text{NH}_2 \\ & | & | \\ \text{do} & + \text{HCHO} & \text{CO} \\ & | & | \\ & \text{NH}_2 & \text{NH}_2 \end{array}$$
- b. Iso butyri denediurea (IBDU) – condensation product of urea and Iso butylenes in the ratio of 2:1.
- c. Urea acetaldehyde = 31.1 % N (Urea + acetaldehyde)
- d. Erotonylidene Urea (CDU) = 32.5 % N
- e. Glycoluril = 39.4% N (Urea + Glyoxal in the presence of HCl.
- f. Difurfuryledene tri Urea = 25% (Urea + Furfural)
- g. Guanyl Urea = 7% N.
- h. Metal am.  $\text{Po}_4$  = 7 – 8 % N.
- i. Traiazines = 32 – 66% N

## III. Coated fertilizers

Fertilizer granules when coated with any suitable material will release N at slow rate.

### a. Sulphur coated Urea

Urea granules are uniformly coated with a layer of sulphur. About 5% coal tar oil is mixed with urea to kill any organism that might break down the coating.

Composition of sca = Urea 80 – 85%  
Sulphur 13 – 16%  
Wax 2%.

**Advantages** It releases N at slow rate.  
It supplied S which is one of the essential

### Release of N depends on

- Thickness of coating
- Placement of fertilizer
- Use of microbicide
- Temperature
- Time of Incubation and
- Redox potential.

### b. Lac coated Urea (LUC)

Shallac is well known as surface coating material. The resin required for coating is prepared by mixing **one part** of matter Shallac with **2 part** of resin at  $260^\circ\text{C}$  and allowed for 30 min to complete reaction. This is cooled, solidified and powdered. Urea granules are uniformly smeared with linseed oil at  $50 \pm 2^\circ\text{C}$  followed by the addition of resin powders. The resin melts and gives coating over urea granules coal tar and soap stone are also added finally the granules are coated with paraffin wax to seal the fine cracks.

<b>Composition</b>	Urea 73.7 %	Resin 16.2%
	Linseed oil 3.3%	Soapstone 2.9
	Wax 3.6%	Coal tar 0.3

### 3. Neem coated Urea

100 kgs of urea is mixed with solution of 1 kg of coal tar in 2 litres of kerosene, 20 kgs of finally powdered neem cake is added in a drum and the contents were thoroughly mixed.

### 4. Urea super Granules (USG)

For controlling the release of N is to increase the size of the granule to such an extent that it affects its solubility. It is called as “Urea briquettes or super granules”.

### 5. Gypsum coated urea

100kgs of urea + 20Kgs of gypsum. Urea is coated with gypsum to ensure slow release prop. and prevent losses with the added secondary nutrients calcium, sulphur.

### Properties of compound fertilizers.

1. H<sub>2</sub>O soluble
2. N and P are easily available to the plants.
3. Acid forming fertilizer.
4. Because of highly soluble in H<sub>2</sub>O, P is converted into Fe and Al in the acid soils.
5. Key often it is used for top dressing.

		N	K <sub>2</sub> O	P <sub>2</sub> O <sub>5</sub>
Eg.	KNO <sub>3</sub>	13.5%	44%	-
	DAP	18%	-	46
	MAP	16%	-	48
	Ammophos	16%	-	20

### Complex fertilizers

#### Characteristics of complex fertilizers

- a. High analysis fertilizer
- b. Uniform grain size is ensured.
- c. Good physical condition for application.
- d. Easy for handling.
- e. Nutrients are H<sub>2</sub>O soluble form
- f. Per unit basis it is comparatively cheaper.
- g. Application should be done at a single stroke.
- h. Non caking and non hygroscopic fertilizers.

### Mixed fertilizer

#### Raw materials.

Urea  
H<sub>5</sub> PO<sub>4</sub> + Condition + filler  
KCl

**4 ingredients** are used for preparing mixed or complex fertilizer.

**a. Nutrient element** It should be straight of compound fertilizer.

**b. conditions** Some fertilizer are hygroscopic and caking properties. To overcome this conditioners are used.

**c. Neutralizing material** It is added to neutralize the acidity / basically eg. Lime stone, CaCO<sub>3</sub>, talc.

**d. Filler material** is a weight make up materials added to fertilizer materials to produce a mixture of desired grade eg. Clay , sand , saw dust.

## Phosphorus

It is present as Apatites of

1. Chlor apatite	$\text{Ca}_{10}(\text{PO}_4)_6\text{Cl}_2$
2. Fluor apatite	$\text{Ca}_{10}(\text{PO}_4)_6\text{F}_2$
3. Carbonate apatite	$\text{Ca}_{10}(\text{PO}_4)_6\text{CO}_3$
4. Hydroxy apatite	$\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$

**Based on solubility** the phosphate fertilizer are grouped into

- |   |   |                            |          |
|---|---|----------------------------|----------|
| 1. $\text{H}_2\text{O}$ soluble $\text{P}_2\text{O}_5$                  | : | SSP                        | 16%      |
|   | : | TSP                        | 45%      |
| 2. Citrate soluble $\text{P}_2\text{O}_5$                               | : | DCP                        | 34%      |
|   | : | TCP                        | 24%      |
| 3. Acid soluble $\text{P}_2\text{O}_5$                                  | : | Anhydrous R. $\text{PO}_4$ | 16 – 18% |
| 4. Both $\text{H}_2\text{O}$ and citrate soluble $\text{P}_2\text{O}_5$ | : | Kottka $\text{PO}_4$       | 25%      |

It is proposed by acidulating Rock  $\text{PO}_4$  with  $\text{H}_2\text{SO}_4$ .

## Interaction of phosphatic fertilizers with soil

Fertilizer P		Plant uptake P
Dissolves in soil	Very little P	Pool of plant avail
	Stays in soil solution	$\text{P H}_2\text{PO}_4$
Fast reaction		


Reacts with oxides of Fe, Al, $\text{CO}_3$ 's etc.	Contribute towards
--	--------------------

Diffuse away

Reaction products	Plant avail P in soil
ppt out differ in availability	

## Phosphatic fertilizer – single super $\text{PO}_4$

Ground rock  $\text{PO}_4$  is mixed with  $\text{H}_2\text{SO}_4$  in a specially designed mixer, which discharge the product on a belt conveyer where the reaction is completed.



The reacted mass is sent to a curing shed where the products is stored for 3 -4 weeks for curing and drying.


**Properties** It contains 16%  $\text{H}_2\text{O}$  soluble  $\text{P}_2\text{O}_5$   
It contains 4 % free  $\text{H}_3\text{PO}_4$

<b>FCO specification</b>		<b>SSP</b>	<b>TSP</b>
Moist % by weight max	:	12.0	12.0

Free H <sub>3</sub> PO <sub>4</sub> % by weight max	:	4.0	3.0
H <sub>2</sub> O soluble P <sub>2</sub> O <sub>5</sub> % by weight min	:	16.0	42.5
		16.0	Total P <sub>2</sub> O <sub>5</sub> 46.0


### Triple super PO<sub>4</sub>

It is produced by reacting



### Dicalcium Phosphate

It is prepared by reacting the ortho H<sub>3</sub>PO<sub>4</sub> with Rock PO<sub>4</sub> or CaO.



In the national lab. Pune, it was prepared on semi commercial scale by reacting R.P. with HCl.

### Advantage

40 -44% total P<sub>2</sub>O<sub>5</sub> of which 35 – 40% soluble in Neutral ammonium citrate solution. It has got good excellent physical condition and high P<sub>2</sub>O<sub>5</sub> content, the over head costs are less.

### Tricalcium phosphate


It is prepared by deflourinating the R.P and melting it with silica in a furnace (1) The materials is H<sub>2</sub>O insoluble but citrate soluble. (2) It contains 20 – 24% citrate soluble P<sub>2</sub>O<sub>5</sub>. (3) It has an excellent physical condition and can be stored even openly.

### Meta phosphates

- (i) Calcium meta PO<sub>4</sub>
- (ii) Potassium meta PO<sub>4</sub>

#### (i) Calcium meta PO<sub>4</sub>

Phosphorus is burnt in a furnace where gas and gaseous combustion products pass through the lump. R.P.O<sub>4</sub> is packed in the upper part of the furnace. The R. PO<sub>4</sub> and hot vapour are chemically combined to form calcium meta PO<sub>4</sub> which contains 62 – 64% citrate soluble P<sub>2</sub>O<sub>5</sub>. high grade 'P' fertilizer used for direct soil application and for mixed fertilizer manufacture.



#### Potassium Meta PO<sub>4</sub>

This material is prepared in the same manner by substituting with KCl instead of Rock PO<sub>4</sub>. It contains 60% P<sub>2</sub>O<sub>5</sub> of which 57 to 60% citrate soluble.

### Bone meal

It is prepared by crushing of bones. Available in 2 form.

	Total P <sub>2</sub> O <sub>5</sub>	Citrate soluble	Total N
1. Raw Bone Meal	20%	8%	3
2. Steamed Bone Meal	22%	16%	Trace

Finer the size of the bone meal the availability of P<sub>2</sub>O<sub>5</sub> is more. Particle size is <2.24mm is considered suitable for fertilizer. Bone meal is produced in our country is 30,000 to 35,000 MT/Year.

After removal of gelatin. Finer materials are steamed and exported chiefly for feeding live stocks. The coarse particles are used as fertilizers.

### **Steamed Bone Meal or Bone $\text{PO}_4$**

Steamed Bone Meal or Bone  $\text{PO}_4$  is white grayish material, powder, light weight.

### **Method of manufacturing**

Young bones are boiled and steamed at high pressure for removing fats that are used for manufacturing of getatin sticks. The bones are then ground for easy handling and distribution. Daring application it should. be mixed with saw dust calcium soil because it is already light in weight.

Chemically it is  $\text{Ca}_3(\text{PO}_4)_2$  and it is commonly refused as **Bone phosphate of lime**. Generally young bones contain less p and more N than older bones.

### **Crop Response**

- The availability is superior to Rock  $\text{PO}_4$  but slightly lower to basic slag.
- Liming the soil depress the solubility.
- Highly suited for home gardens and flower gardens. Since it does not have salt damage.

### **Basic slag**

It contains 3 – 8 and  $\text{P}_2\text{O}_5$  it can be used for neutralizing soil acidify as it contains **Calcium** as **Calcium silicate** and many trace elements.

Basic slag is mixed with powdered R.  $\text{PO}_4 + \text{H}_2 \text{SO}_4$  to produce ‘Pelophos’ (at Rourkala) containing 17%  $\text{P}_2\text{O}_5$  of which 16% is citrate soluble.

### **Mussoorie $\text{PO}_4$**

The name **Mussoorie Phosphorus** is derived form the place of its origin **Mussoorrie** Uttar Pradesh, India (dt. Dehradun)

The  $\text{P}_2\text{O}_5$  content ranges from – Fresh ore : 16 – 25% and weathered ore : 23- 32%

### **Composition of the blended material is given below**

1.	Phosphorus as $\text{P}_2\text{O}_5$	:	20 – 24 %
2.	Calcium as CaO	:	37 - 45%
3.	Carbondioxide as $\text{CO}_2$	:	15 – 20 %
4.	Silica as $\text{SiO}_2$	:	5 – 10 %
5.	Magnesium as MgO	:	4 – 7 %
6.	Total Iron as $\text{Fe}_2\text{O}_3$	:	2 – 5%
7.	Fluorine as F	:	1 – 3%
8.	Water as $\text{H}_2\text{O}$	:	1- 3 %
9.	Aluminium as $\text{Al}_2\text{O}_3$	:	0.5 – 1.5 %
10.	Sulphur as S	:	0.5 – 1.0%
11.	Potassium as $\text{K}_2\text{O}$	:	0.1 – 0.5%
12.	Vanadium as Va	:	250 – 500 PPM



13.	Barium as Ba	:	400 – 1000 PPM
14.	Zinc as Zn	:	100 – 150 PPM
15.	Chromium as Cr	:	40 – 60 PPM
16.	Nickel as Ni	:	25 – 150 PPM
17.	Molybdenum as Mo	:	5 – 75 PPM
18.	Copper as Cu	:	10 – 50 PPM

### Classification of K fertilizers

- K fertilizer carrying chloride - Muriate of potash KCl.
- K fertilizer without chloride -  $K_2SO_4$ ,  $KNO_3$   
For tobacco and potato  $\rightarrow K_2SO_4$  is used to avoid Cl injury.

### KCl manufacture

Raw materials  $\rightarrow$  sylvinite mineral. It is mixture of KCl and NaCl. KCl is much more soluble in hot water than in cold  $H_2O$ . If solution saturated with both NaCl and KCl, the concentration of NaCl decreases as the temperature increases. KCl is extracted from sylvinite utilizing solubility characteristic KCl is removed, evaporated, crystallized and powdered.

### Sources of potash

- Mineral deposits - Impt. Source of K fertilizers.
- By product from industry
- Synthesized K fertilizer ---  $KNO_3 \rightarrow 44 - 45\% K_2O$ .

#### 1. Increasesal deposits

##### a. $H_2O$ soluble minceasesals

1. Sylvinite	KCl	:	63 % $K_2O$ (mix of K Cl and NaCl)
2. Carnolite	$MgCl_2 \cdot 6 H_2O$	:	17% $K_2O$ (mix of K Cl and NaCl)
3. Longbeinite	$K_2SO_4 \cdot 2 MgSO_4$	:	27% $K_2O$ (mix of K Cl and NaCl)
4. Kyanite	$KCl \cdot MgSO_4 \cdot 3 H_2O$	:	19% $K_2O$ (mix of K Cl and NaCl)

##### b. Less soluble in $H_2O$

Poly halite  $K_2SO_4 \cdot 2 MgSO_4 \cdot 2 CaSO_4 \cdot 2 H_2O$  : 15%  $K_2O$

##### c. Insoluble in $H_2O$

- Feldspar  $KAlSi_3O_8$
- Mica – Biotite or muscovite
- Glauconite  $KFeSiO_6 \cdot 2 H_2O$ .

### 2. Industrial Bi product

- Flue dust from
  - Steel industry = 3 – 18%
  - Cement industry = 3 – 13%
- Pot. Schoentie
  - obtained from salt pitters
  - double salt of K & Mg.
  - Very good K fertilizer for acid soil.

### Potassium sulphate

Dissolving longbeinite( $K_2SO_4 \cdot 2 MgSO_4$ ) in  $H_2O$  in which concentration Solution of KCl is added. Then  $K_2SO_4$  is precipitated, separated and purified.



This reaction is carried out in a special furnace consists of cast iron muffle with rotating ploughs to agitate the reaction mixture. The gaseous HCl is cooled and absorbed in  $H_2O$ .

### Properties

White crystalline product contains 54%  $K_2O$ .

Sp. gravity 2.662

Melting point  $1067^{\circ}C$ .

Solubility in  $H_2O$  = 13% by wt.

### Secondary nutrient

#### Calcium

**Sources** Dolomite, Calcite, Apatite, Calcium feldspar.

As a general rule, coarse textured, humid region soils formed from rocks low in calcium containing minerals are low in their content of this element. Soils that are fine textured and formed from rocks with high in calcium content minerals are much higher in their content both exchange calcium and total calcium

Gypsum $CaCO_3 \cdot 2H_2O$	:	22.3	SSP	:	19.6
Calcium nitrate	:	19.4	TSP	:	14.3

#### Magnesium

<b>Sources</b>	Earth crust	:	1.93%
	Biotite		
	Dolomite		
	Chlorite		
	Basic slag	:	3.4
	Epsom salt	:	9.6

### Organic manures

#### Farm yard manure

“Consists of decomposed or partly decomposed mixture of dung, urine and the litter. It is a mixture of vegetative refuse from the farm and it mostly consist of the nutrient elements required by the plant.

It is very good source of manure to replace the deficient of nutrient.

#### Composition of dung

75 – 80 % is made up of moisture.

20 – 25% is insoluble materials of fodder or food. Chief components of insoluble materials can be **woody fibre, starch, fats, bile pigments, mucus**. In addition to this it also may contain **skatole and Indole** by product derived from the digestion of food material. It also contain **Fatty acids, Ca, Mg  $PO_4$ , Alkaline soap etc.**

#### Nutrient content of dung %

	Horse	Cattle	Buff	Sheep	Pig	Poultry
H <sub>2</sub> O	76.5	80.0	81.0	61.9	80.7	57.0
Organic Mater	21.0	15.2	12.7	33.1	17.0	29.3
Mineral matter	3.3	3.8	5.3	4.7	3.0	5.6
N	0.47	0.30	0.26	.70	.59	1.46
P <sub>2</sub> O <sub>5</sub>	.38	.18	.18	.51	.46	1.17
K <sub>2</sub> O	.30	.18	.17	.29	.43	.62
Lime (CaO)	.17	0.36	.46	.46	.07	

Poultry manure is very rich in nutrient content compare to others. Next to poultry sheep manure is fairly rich, cattle and buffaloe dung is poor in nutrient content.

### Urine

It contains 96% H<sub>2</sub>O and A% is made up of dissolved substances primarily **urea** derived from the **protein catabolism** in the **blood** and partly from **leucine and tyrosine** as a result of **tryptic digestion** in the intestine. It is also derived as a by product of disintegration in the various organs of the animal body. Other was to subs in urine in addition to urea are **uric** and **Hyppuric** acid.

- |                 |  |                           |
|-----------------|--|---------------------------|
| Inorganic salts | <ul style="list-style-type: none"> <li>a. sodium chloride – 1%.</li> <li>b. Sodium hydrogen Po<sub>4</sub></li> <li>c. Ca and Mg Po<sub>4</sub></li> <li>d. K and Na So<sub>4</sub></li> </ul> | } in varying proportions. |
|-----------------|--|---------------------------|

Nearly 50% of N content of FYM comes form **Urine**.

### Composition of Urine

	Horse	Cattle	Buff	Sheep	Pig
H <sub>2</sub> O	89.6	92.6	81.0	86.3	96.6
Organic Mater	8	4.8	-	9.3	1.5
Mineral matter	8	2.1	-	4.6	1.0
N	1.29	1.21	0.62	1.47	0.38
P <sub>2</sub> O <sub>5</sub>	.01	.01	Traces	0.05	.10
K <sub>2</sub> O	1.39	1.35	1.61	1.96	.99
Lime (CaO)	.45	.01	-	.16	.01

### Litter

It is used as a bedding material in foreign country. In Indian condition it refuse to rejects and waste materials. It includes straws of cereals, stalks of sunflower etc. If is used to conserve the loss of NH<sub>3</sub> from urine. If urine is treated with soil (or) soil is soaked with urine (or) matured straws of sorghum, Cumbu, there is a wide range of C : N ratio.

Eg. Sorghum 50 : 1

### Characteristics of Farm yard manure

#### a. Moisture :

75% is made up of moisture. In general moisture content is 50-80% and dry matter 20-50%

**b. It is low analysis manure :**

one lone of FYM will supply 5 kg N, 2.5 kg  $P_2O_5$  , 5 kg  $K_2O$ .

**c. Residual effect of FYM**

In FYM nutrients are present in complex form as proteins or microbial tissues. During the process of decomposition the microbial tissues are decomposed and form a source of N. In well decomposed FYM about  $\frac{1}{2}$  N, K,  $\frac{1}{6}$  P readily available to first crop.

Remaining part will left in the soil for the succeeding crops. In well decomposed FYM the C:N ration is 12-15 : Partly decomposed FYM the C:N ration is 20 :. IN such case the residual effect is more for 2 or 3 succeeding crops.

**d. Nutrient is unbalanced.**

FYM is poor in NPK content is particularly  $P_2O_5$ . So it is considered as unbalanced. FYM is mixed with super (or) Rock  $PO_4$  before addition not the field,  $P_2O_5$  content can be increased.

FYM + super  $\rightarrow$  Rein forked. So it is considered us complete manure. It also supplied secondary and micronutrients.

**Advantages of farm yard manure**

1. It is a complete manure. It supplies major, micro and trace elements.
2. It improves soil fertility.
3. It improves physical, Chemical, Biological and Properties of soils.

**Green Manures**

**Green manure :**

Growing the plants i.e. sun hemp and ploughing insitu (or) adding green matter obtained from the field.

**Green leaf manure**

Addition of green or plant tissues obtained from trees, herbs, shrubs pruning and unwanted weeds.

<b>Green manure</b>	<b>N</b>	<b><math>P_2O_5</math></b>	<b><math>K_2O</math></b>
Sunhemp	0.5	0.1	0.4
Danicha	<b>1.1</b>	<b>0.2</b>	<b>0.4</b>
Sesbania	0.7	0.1	0.6
Pelli pesera	0.7	0.2	0.4
Cluster bean	0.3	0.1	0.3
Cow pea	0.7	0.2	0.6
<b>Green leaf manure</b>			
Kolungi	0.7	0.1	0.4
Calotropis	0.3	0.1	0.6
Pungam	<b>1.2</b>	<b>0.2</b>	<b>0.8</b>
Glyricidia	0.8	0.1	0.7
Puvarasu	0.8	0.2	0.9

### Benefit of Green manure or Green leaf manures

- a. Addition of organic matter.
- b. Adds Nutrient – Macro, Secondary and micro elements.
- c. It improves physical condition of soil.
- d. It is useful for the reclamation of problem soils.
- e. It act as a cover or catch crop prevent soil erosion, conserves moisture, prevent nutrients, Leaching.
- f. When leguminous GM is grown they fix the atmospheric – N by the roots and improves N status of the soil.

### Characteristics of Green manure

1. It should **have rapid growth**.  
Farmers can fit the GM in the crop rotation because of shorter duration – sun hemp grown very faster than other GM crops with in 45 days it attains 4-5' ht.
2. It should be **abundant and succulent**.  
GM should be heavy yielder and succulent. It contains more of H<sub>2</sub>O and less dry matter, decomposition will be rapid and easy.
3. It should have **ability to grow poor soils**.  
Alkaline soil will able to grow **Danicha**

### Factor affecting decomposition

- a. **Organisms present**
  - Bacteria, Fungi, Actinomycetes
  - Aerobic - Rapid decomposition
  - Anaerobic - Slow decomposition.
- b. **Temperature** : Optimum temperature for decomposition is **30-35°C** under this condition the growth of organism will be more.
- c. **Aeration** : Multiplication of organism is more, aerobic condition decomposition is fast.
- d. **Moisture** : Proper decomposition and growth of organism H<sub>2</sub>O is needed.
- e. **Soil factors** : Physical chemical and biological clay soil decomposition is slow.
- f. **Nature of GM** :
  - If it is succulent - quick decomposition
  - Non- succulent - slow decomposition

### Concentrated organics

1. Oil cakes
2. Various meals is Blood meal, Fish meal, Bone meal, Salughter house.
3. Sullage : It contains waste waters received from. Bath room and kitchen waste H<sub>2</sub>O only.  
Sewage : Night soil with water.

Sludge : Solid portions of Night soil. It is allowed for fermentation.

### Oil cakes

In India about 2m.t. of oil cakes are produced / year. Out of this only 85% is edible and 15% is non edible portion. The nutrient present in the oil cakes are insoluble in H<sub>2</sub>O but which are readily available to the plants. (within 10 days). The **Mahuva** cake takes 1- 1 ½ month time for perfect mineralization. The nutrient content of the oil cake.

N	=	2.5 - 8.0 %
P <sub>2</sub> O <sub>5</sub>	=	1.0 - 3.0%
K <sub>2</sub> O	=	1.0 - 2.0%

### Benefits of organic manures

1. It increases the yield of crop through the supply of nutrient.
2. It improves soil fertility through promoting the physical chemical and biological properties.
3. It corrects the deficiency of secondary and micro elements.
4. Oil cakes and various meals are equally or slightly even better than inorganic fertilizers.
5. Slow acting. It may not be completely decomposed state. So the loss of nutrients are minimized.
6. Quality of crops are improved particularly fruit and vegetable.

### Factors affecting the efficient use of fertilizers and manures

#### I. Soil factors

- a. Total nutrient content
- b. Availability of nutrient
- c. Soil reaction
- d. Soil Erosion
- e. Soil texture
- f. Soil drainage
- g. Soil management practices
- h. Presence of impermeable layers

#### II. Climatic and environmental factors

- a. Temperature
- b. Rainfall
- c. Irrigation
- d. Evaporation
- e. Length of growing season
- f. Intensity of sunlight
- g. Duration of sunlight.

#### III. Crop factor

1. Nutrient requirement of crops
2. Crop sequence
3. Crop adaptability to soil factors
4. Crop production practices
5. Crop residues

#### IV. Systems of farming

#### V. Fertilizer characters

### Compost (or) Artificial FYM (or) Synthetic FYM

“Dark coloured powdered material derived from decomposition of plant refuse by the microbes in the presence of little amount of soluble nitrogen”.

## History

During 1921, Rotham Stead, **Hutchinson and Richards** they produce the material called as **synthetic FYM** from straws. They mix the straw with soluble N, P, Ash and allow to decompose under aerobic condition. The Organism which is responsible is “**Cytophaga hutchinsoni**”. Finally they got a manure which is closely resemble to FYM which is called **Artificial or synthetic FYM**.

## Principles in compost making

### a. Composition of refuse material

It consists of	(a) Cellulose or Hemi cellulose	:	60 – 80%
	(b) Lignin	:	15 – 20%
	(c) H <sub>2</sub> O soluble material	:	5 – 12%
	(d) Protein	:	2 -3 %

### b. Mechanism of the decomposition brought out by micro organic.

### c. The knowledge of the metabolism of the micro organic involved.

## Factors affecting the compost

1. Refuse materials - Obtained from farm, straw leaves, Garbage, street sweeping materials.
2. Suitable starter / inoculum's - Which will provided fungi or bacteria for efficient decomposition.
3. Soluble N - Which increases the bacterial activity.
4. Aeration
5. Moisture
6. Temperature

## Methods of compost making

- a. ADCO process - Hutchin son and Richards
- b. Activated process - Fowler and Rege.
- c. Indore Process - Howard and ward.
- d. Bangalore Method - C.N. Acharya.
- e. Coimbatore Method - AC & RI.

### a. ADCO process

Agriculture Development Company – ADCO. They use ADCO powder for rapid decomposition. The plant refuse (or) which can not fed to cattle is used. The refuse material is spread uniformly on the floor to a thickness of 9 – 12”. Then the adco powder is sprinkled @ 7% ie. 7kg/ 100 kg of material.

Adco powder contains soluble N, AM SO<sub>4</sub>, AM – CO<sub>3</sub> certain amounts of P<sub>2</sub> K, lime, rock PO<sub>4</sub>, Ash, KCl. This material is moistened. This can be repeated normally 6 layers. Periodically moistened for quick decomposition. It can be allowed for 3 – 4 months. It is **aerobic** and **Heap method**. For proper decomposition this material is turned or moistened and restate in the form of heap.

### b. Activated process

In this method, instead of adco powder, we use some inoculum or activator to decompose the material quickly. It is a **heap system** and turning is given. The starter should be **dung urine, Urine mixed earth, sewage and sludge** etc. The straw is uniformly spread and moistened and added sewage sludge. This can start the reaction. This must be respected and after moistened, the level was maintained and periodical turning is given. After 3 – 4 months, the entire mass is converted into Brown colour. They called it as **Starter**. 1/3 of it is used for compost making.

### Composting of sugar cane Thrash

#### Raw materials

- |                       |                      |
|-----------------------|----------------------|
| 1. Trash              | : 100 kg / one tonne |
| 2. Pleurotus          | : 3 bottles          |
| 3. Trichoderma viridi | : 20 grams           |
| 4. Urea               | : 5Kg                |

Final compost contains

- |                               |          |
|-------------------------------|----------|
| N                             | = 0.70 % |
| P <sub>2</sub> O <sub>5</sub> | = 0.25%  |
| K <sub>2</sub> O              | = 0.70%  |

	→ 20 kg dugongs 60 lit H <sub>2</sub> O
	→ 100 kg S. trash
	→ 100 g Trash derma
	→ 100 g S. trash
	→ 1 kg urea
	→ 100 kg S. trash
	→ 1 bottle pleurotus
	→ S. Trash - 100 kg

- ❖ Composting is done in trenches. The size of the trench will be 5 x 3 m.
- ❖ S. trash was cut into small spices and spread uniformly on the floor under shade.
- ❖ As in figure.
- ❖ Like wise 10 layers may prepared and finally covered with 250 Kg of clay soil or Red soil for compacting the beds Sprinkle the H<sub>2</sub>O once in a week.

### Long term and permanent manorial experiments to monitor soil fertility, sol productivity and sustainability

PME will provide valuable information on the impact of long term adoption of nutrient management systems with varying sources, types and combinations of plant nutrient inputs on soil fertility and productivity.

The world's oldest long term agricultural field experiments were begun between 1843 and 1856 by Lawes and Gilbert at Rothamsted in England. Eight of this experiments still continue. All these experiments were started as Agronomic experiment to determine the nutrient requirements of agriculture crops.

#### In Tamil Nadu

Eleven PME are being conducted in various locations, viz.,



TNAU	-	Coimbatore
AC & RI	-	Madurai
AC & RI	-	Killikulam
ARS	-	Kovilapatti
ARS	-	Vamban
IRRI	-	Aduthurai
SWMRS	-	Thanjavur.
SRS	-	Cudallore.
ORS	-	Tindivanam
RRS	-	Paiyur
TRS	-	Yethapur

These experiment represent different soil conditions and cropping systems and are most valuable because the data generated from them provide the basis for drawing conclusions on soil sustainability under continuous maintenance with deficient nutrient input systems both under rainfed and irrigated conditions.

The old permanent manorial experiment started in Red soil, on the lines of Rothamsted manorial experiment has been in progress this institute since 1909. So far 140 crops have been raised and the results shown that the higher yields could be obtained by the combined application of NPK. The cattle manure plots maintained the soil fertility. The O.C. content increased upto 30cm depth and decreased thereafter.

In 1925, a New PME was started with the same treatments under irrigated conditions. So far 126 crop have been completed. Results revealed that application of NPK, MP and cattle manure are recorded higher yield over other treatments.

### **Fertilizer control order – 1957** **Essential commodities Act – 1955**

#### **Objective**

The main of this FCO is to regulate equitable distribution of fertilizers and making fertilizers available at fair prices to the consumers.

#### **Essential features of the order**

- a. No manufacture or dealer shall sell any fertilizer at a price exceeding the max- price fixed by the central government.
- b. Every manufacture or dealer shall give a cash or bill memorandum to a purchases.
- c. No person shall carry on the business of selling fertilizer at any place except under and in accordance with the terms and conditions of the license granted to him under this order.
- d. The State Government may appoint such number of persons as it thinks necessary to be inspectors of fertilizers.
- e. No person shall export or attempt to export of any fertilizer from any state.
- f. Any Inspector of fertilizers appointed under FCO can enter and search any place, Seize any fertilizer in respect of which he has reason to believe that nay prevision of this order is being controversial along with the packages, coverings.

- g. Every manufacture of manure mixture and straight fertilizers shall legibly mark on the containers of such fertilizers the following particulars namely,

**A. On the containers of manure mixtures**

1. Register number if any, brand and name of the manure mixture.
2. The N, Total  $P_2O_5$ ,  $H_2O$  Soluble  $P_2O_5$  and  $K_2O$  contents denoted by the letters N, P(T), P(S) and K respectively.
3. Registered name of the firm and trade mark, if any.
4. The crop or crops for which the manure mixture is suitable and in the case of general mixture for all crops that expression "All crops"

**B. On the containers of straight fertilizers**

1. Name of the manufacture.
  2. Trade mark if any
  3. Name of the fertilizer
  4. percentage by wt of N/P/K.
  5. Gross and Net content by weight in kilograms.
8. As per this order, a dealer may retain at any time one bag or containers of each variety of fertilizer in an open unsealed condition for the purpose of retail sale.
9. Every containers shall be so sealed that the content, there of cannot be tampered with, with out breaking the seal. For fertilizer manufactured in India, if packed in bags and stitched in hand shall bear "Lead" seals and lead seals are not necessary of such bags are machine stitched.
10. The fertilizer inspector can draw samples of fertilizer for analysis in accordance with the procedure of drawal of samples and send them to state fertilizer laboratories for testing in accordance with the procedure prescribed in this order.

As per this order, there is strict specification for each and every fertilizer for their nutrient content moisture and ingredient if any.

**Tolerance or permissible limits**

- a. Fertilizer with definite compounds like  $(NH_4)_2$  SO<sub>4</sub>, Urea, super (or) potash up to 0.2 unit of the nutrient content is above 20%. 0.1 unit of the nutrient content is below 20%.
- b. CAN : 0.3 unit, C. DAP, ASN, Mixture and complex = 0.5unit

The Improvement feature of this FCO is in sampling of the fertilizer so in **drawing samples, the following measures and precautions shall be observed.**

- a. Samples shall not be taken at a place exposed to weather.
- b. Sampling instruments shall be clean and dry when used.
- c. The material being sampled, the sampling instruments and the containers for samples shall be well protected from contamination.
- d. To draw a representative sample, the contents of each containers selected for sapling shall be mixed as thoroughly as possible by suitable means.
- e. The sample shall be placed in suitable, clean, dry and air tight glass or other suitable containers
- f. The sample containers shall be in such size that are almost completely filled by the sample.

- g. Each sample containers shall be sealed air tight after filling and marked with full details of sampling, date of sampling etc.
- h. Sample shall be stored in cool places.

	<b>Lot size</b>	<b>Number</b>	<b>Containers to be selected</b>	
1.	2 – 8	2	101 – 300	6
2.	9 – 27	3	301 – 500	7
3.	28 – 64	4	501 – 800	8
4.	65 – 100	5	801 – 1300	9
			> 1301	10

### **Concepts / approaches for soil test based fertilizer Recommendations**

“Soil testing is a phase where the basic research in soil chemistry culminates and applied research on soil fertility begins. the combined research efforts on soil chemist and soil fertility expert, are reflected in an translated to farmers as fertilizer recommendations”.

#### **(Goswami, 1982)**

1. General / Blanket Recommendation
2. Soil Test rating and fertilizer adjustment
3. Agronomic approach
4. Critical soil test level approach (Cate and Nelson 1965)
5. Foliar diagnosis
6. Liebig’s model
7. FR for certain percentage of yield maximum
8. Deductive approach
9. Inductive approach
10. Targeted yield approach/ prescription procedure.
11. Integrated STCR correlation approach
12. Soil fertility cum soil survey.