

A Scientific approach to Health

AEC 26

UNIT 3 – CREATION OF HEALTHY and CARING RELATIONSHIPS

Building Communication Skills

Communication and its essence

- Personal Process
- Communication is the art of transmitting information, ideas and attitudes from one person to another.
- Communication is the process of meaningful interaction among human beings.
- Involves change in behavior
- Means to influence others
- Expression of thoughts and emotions through words & actions.
- Tool for controlling and motivating people.
- It is a social and emotional process.



Building Communication Skills

LISTENING



FRIENDLINESS



OPEN MINDEDNESS



Essential Communication Skills for Your Career

FEEDBACK



CONFIDENCE



NON-VERBAL COMMUNICATION



Communication Process

Sender: The person who conceptualized the idea and wants it delivered to the recipient.

Encoding: The way the information is described or translated into a message.

Message: The idea, fact or opinion that the sender wants to communicate.

Communication channel: The method of delivering the message.

Noise or Distraction: Hinders successful communication.


Types : Psychological, Physical, Environmental and Semantic.

Receiver: The target audience of the message.

Decoding: The interpretation of the message.

Feedback: The response or action a receiver takes after decoding a message.

Communication Process



The Communication Process

The 3 Important Rules of Communication

In any language, there are three extremely important points to remember when you're communicating with someone.

- 1. Say what you mean.** It can be difficult to express (say) some ideas clearly, but if you're trying to prevent miscommunication, it's important to say exactly what you mean. Be clear and to the point.
- 2. Ask questions.** Communication is two-way, which means you can't be the one doing all the talking. To make sure your listener is *engaged* (interested in what you have to say) and understanding you, ask questions. See #5 below for good types of questions to ask.
- 3. Listen.** We mean *really* listen. Hear what your speaking partner has to say, and try to understand what they mean.

Barriers of Communication

Types of Barriers

- Psychological Barriers
 - Physical Barriers
 - Emotional Barriers
 - Attitudinal Barriers
 - Cultural Barriers
 - Language Barriers
 - Gender Barriers
 - Interpersonal Barriers
- No interest in the conversation.
 - Jumping to conclusions without waiting for the whole message.
 - Inability to Listen to Others.
 - Lack of Transparency & Trust.
 - Communication Styles (when they differ).
 - Fear of offending the other person by expressing your opinions.
 - Cultural Differences & Language.
 - One feels not comfortable sharing your feelings with the other person

Listening

6 KEY ACTIVE LISTENING SKILLS



1. PAY ATTENTION.



2. WITHHOLD
JUDGEMENT.



3. REFLECT.



4. CLARIFY.



5. SUMMARIZE.



6. SHARE.

WAYS TO IMPROVE THE COMMUNICATION SKILLS :

- 1) **Listen without judgment** : The key to good communication is listening well. Save your judging for later after you have heard and understood what was said.
- 2) **Listen with the willingness** :
- 3) **Listen without thinking about what you will say next.**
- 4) **Do not be invested in being right.**
- 5) **If your mind wanders, ask for repetition.**
- 6) In all cases repeat back what you heard and ask if it is correct.
- 7) Listen to yourself.
- 8) Say it honestly, but with consideration for the listener's feelings. Be polite, respectful and sincere.
- 9) **Understand and acknowledge that most things are not black or white, but somewhere in a gray area.**
- 10) Have integrity and build trust. Don't say what you don't mean. Don't promise what you won't or can't fulfill. Follow through with any commitments you make.
- 11) Eye contact - **Eye contact is also a big giveaway; making and maintaining eye contact is a big indicator of confidence.**



Steps to avoid embarrassing in communication with others:

1. **Be patient** Your ideas are important, but no more important than anyone else's.
2. **Listen carefully** If you are busy thinking about what you want to say, you aren't going to hear what the other person is saying.
3. **Take notes if necessary** If the discussion is a long one, notes can help you remember the points you want to make.
4. **Rephrase** what was said in your own words This is the clearest way to ensure you haven't made an assumption about what a statement means.
5. **Don't interrupt** The speaker may be going on to clarify exactly what you are questioning.
6. **Pause and reflect** : Allow some time to let the speaker's comments fully sink in by pausing and reflecting before you jump in to speak.



Friends and Friendship - Education

- Friend : "a person with whom one has a bond of mutual affection, typically one exclusive of family relations"
- A person who has a strong liking for and trust in another person
- A person who helps or supports something She was a friend to environmental causes.
- Friendship : "Friendship is a relationship of mutual affection between people. It is a stronger form of interpersonal bond than an acquaintance or an association, such as a classmate, neighbor, or colleague. "
- It is a dyadic relationship (**Smallest Possible social group**), meaning that it involves a series of interactions between two individuals known to each other.
- It is recognized by both members of the relationship and is characterized by a bond or tie of reciprocated affection.
- It is not compulsory; two individuals choose to form a friendship with each other.



Friends and Friendship - Education

- It is almost always characterized by companionship and shared activities.
- One of the primary goals and motivations of friendship is companionship (company).
- In addition, teenage and adult friendships often perform other functions, such as serving as sources of emotional support and providing opportunities for self-disclosure and intimacy.



Qualities Of A True Friend - A faithful friend is a sure anchor

- Will always be honest with you.
- Loves you for yourself and unconditional
- Gives more than they take.
- Compromises
- Forgives you even without an apology.
- Believe in you.
- Make time for you.
- Respectful to you.
- Thoughtful.
- Stick around you.

“Friendship in Education”

- They provide support and resources and can both encourage and discourage academic achievement.
- Contribute to a positive sense of mental well-being
- Encourage learning and develop interpersonal skills
- Play a significant role in promoting overall health.
- Development of Social Skills
- Better Cooperation
- Better Problem solving abilities
- Better Communication
- Better Academic outcomes



The Value of Relationships and Communication

”WHY COMMUNICATION IS KEY TO HEALTHY RELATIONSHIPS”

- Communication is important in any relationship, as it allows you to effectively share feelings, opinions and expectations.
- GET TO KNOW EACH OTHER : The key to determining whether you're compatible with another person is to communicate with them from the offset.
- AVOID MISUNDERSTANDING : Another reason to communicate effectively is to avoid any misunderstandings, which can lead to a breakdown of communication or the relationship as a whole.
- SET CLEAR EXPECTATIONS : In any relationship, it's important that these expectations are made clear from the offset, to avoid one party upsetting the other without realizing. Always be open and honest and never be afraid to stand by what you believe in. Thus healthy, positive and satisfying relationship.

”HOW TO IMPROVE COMMUNICATION IN A RELATIONSHIP”

- Always talk about important or sensitive issues face-to-face.
- Be open and honest.
- Find the right time for important conversations.
- Don't be insulting or verbally aggressive, even if it's an emotive issue.
- Approach conversations in a calm and relaxed way, in a quiet environment where both parties are comfortable.
- Use positive body language.
- Choose your language carefully.



EXAMPLES OF GOOD COMMUNICATION IN RELATIONSHIPS

- Giving your partner, friend or family member your full attention.
- Giving each other space when needed.
- Face-to-face communication.
- Honesty and openness.
- 24- hour rule – never sleep on an argument.
- Talk about the little things in life.

"EXAMPLES OF BAD COMMUNICATION IN RELATIONSHIPS"

- Belittling each other Openly giving the cold shoulder
- Becoming defensive in heated discussions
- Assuming you know what the other person is thinking
- Arguments that are never resolved and brushed under the carpet
- Inability to compromise
- Few attempts to connect



Relationships for Better or Worsening of Life

- Satisfying relationships not only make us happy, they also influence our long-term health as much as getting enough sleep, eating healthy, and not smoking.
- Many research studies have shown that satisfying relationships are associated with better health, greater happiness, and even longer life.
- Stronger Heart and Healthier Brain
- An unhealthy relationship will cause lot of stress, loneliness and will make us an irritable person which will work as a retarding force in our way of achieving success.
- While in an unhealthy relationship we lose trust on each other and respect for each other decreases.
- A research shows that a healthy relationship have 5 to 1 ratio of positive to negative behaviors and a unhealthy relationship have a much lower ratio of positive to negative behaviors.


Understanding Basic Instincts of Life (More than a Biology)

”Three Basis instincts of Human Life”

- Self-Preservance
- Sexual Instinct
- Social Instinct.



Basic Instincts of life which create self



Changing Health Behaviors Through Social Engineering

Modifying the environment in ways that affect people's ability to practice a particular health behavior –

- Banning certain drugs such as heroin and cocaine
- Requiring vaccinations for school entry
- Using safety containers for medications
- Lowering speed limits
- Raising the drinking age


Changing Health Behaviors Through Social Engineering

- Venues for health habit modification
- Private therapist's office
- Health practitioner's office
- Family
- Self-help groups
- Schools and colleges
- Workplace interventions
- Community-based interventions
- Mass media such as newspaper and magazine articles, published photographs, recordings of television and radio broadcasts, music recorded for mass distribution, advertisements, books
- Cellular phones and landlines
- Internet

UNIT-4 : **Avoiding risks and harmful habits**

**Characteristics of health compromising behaviors,
addictions-recognizing and avoiding, addictive
behaviors,**

Health Related Behaviour

 *Definition – “Any behaviour that affects the health of an individual either positively or negatively is considered to be his/ her health related behaviour.*

 *It includes physical activities, balanced diet, proper intake of water, smoking, alcohol consumption, sleep, meditation, health care precautions, yoga, exercise, etc.*

Health Related Behaviour

*Health Related
Behaviour*

*Health
Promoting
Behaviour*

*Health
Compromising
Behaviour*

Health Related Behaviour



There are number of factors that affects or determine the health related behaviour. For eg :



1. Changing demographics



2. Social and cultural disparities



3. Psychosocial and environmental determinants



4. Individual differences



5. Cognitive factors

Health Related Behaviour



www.examrace.com

Health Promoting Behaviour



There are number of health related behaviour that are considered to be positive and have a positive effect on our body. Such behaviour are promoted and advised to put in use for the betterment of our health. These type of behavior are known as health promoting behaviour.

Health Promoting Behaviour

1. Regular
Physical activities

3. No drug or
alcohol consumption

2. Balanced diet
& a low fat diet

4. Meditation, etc.

Health Promoting Behaviour

- ★ It lower the death rate, lower the risk of death rate due to coronary diseases, promotes healthy habits and lifestyle of living and lower the blood pressure problems.

Health Compromising Behaviour

- ★ There are certain types of health related behaviour that deteriorates our health and pose a negative impact on the health.
- ★ Such type of health behaviours come under the category of health compromising behaviour.
- ★ In order to live long and healthy, the individual should avoid engaging in such type of activities.

Health Compromising Behaviour

1. Alcohol
and drug use.

3. Unhealthy
eating habits

5. No exercise or
physical activities, etc.


2. Smoking

4. Stress
inducing activities

Health Compromising Behaviour

- ★ Such type of behavior increases the risk of coronary heart diseases, blood pressure issues, breathing problems, indigestion, increased death rate and lower quality of life.

Models of Health Related Behaviour



Models of Health Related Behaviour

**Behavioural
Economics
Theory**

**Cognitive
Social Learning
Theory**

**Health
Belief Model**

**Theory of
Planned of
Behaviour**

Conclusion

★ The health related behaviour vary in intensity, duration, frequency and manner in its effect on individual health.

★ The health practitioners and experts are integrating all sorts of theories and trying to figure out methods to promote healthy behaviour and induce health protective behaviour in individual.

Conclusion



Success have been found in smoking cessation, adherence to physical activities and medication, reduced risks of CAD diseases.

Recognizing and avoiding of addictions

- Someone with an addiction will crave a substance or other behavioral habits. They'll often ignore other areas of life to fulfill or support their desires.
- Lack of control, or inability to stay away from a substance or behavior
- Decreased socialization, like abandoning commitments or ignoring relationships
- Ignoring risk factors, like sharing needles despite potential consequences



- Physical effects, like withdrawal symptoms or needing higher dosage for effect
- Problems at school or work — frequently missing school or work, a sudden disinterest in school activities or work, or a drop in grades or work performance.
- Physical health issues — lack of energy and motivation, weight loss or gain, or red eyes, neglected appearance, lack of interest in clothing, grooming or looks.
- Changes in behavior — exaggerated efforts to bar family members from entering his or her room or being secretive about where he or she goes with friends; or drastic changes in behavior and in relationships with family and friends.

- Money issues — sudden requests for money without a reasonable explanation; or your discovery that money is missing or has been stolen or that items have disappeared from your home, indicating maybe they're being sold to support drug use.



Recovering from addictions

- Admit There Is A Problem. The hardest part to recovery is admitting you have an addiction.
- Reflect On Your Addiction.
- Seek Professional Support.
- Appreciate The Benefits of Sobriety.
- Evidence-Based Care.
- Identify Your Triggers.
- Change Your Environment.
- Exercise.

How addiction develops and addictive behaviors

- Substance Use Disorder (SUD) is a complex condition in which there is uncontrolled use of a substance, despite harmful consequences.
- People with SUD have an intense focus on using a certain substance(s) such as alcohol, tobacco, or illicit drugs, to the point where the person's ability to function in day to day life becomes impaired.
- People keep using the substance even when they know it is causing or will cause problems.
- The most severe SUDs are sometimes called addictions

UNIT-4 : Avoiding risks and harmful habits

Types of addictions, influencing factors for addictions, Differences between addictive people and non addictive people and their behavior with society

Addictions

What is Addiction?

- . Addictive behaviour: A behaviour based on the pathological need for the substance or activity, may involve the abuse of substances, such as alcohol or tobacco/opium or the excessive ingestion of high caloric food, resulting in extreme obesity
- . Addiction is a chronic disorder with biological, psychological, social and environmental factors influencing its development and maintenance.
- . Addiction occurs when the act of using a substance takes over these circuits and increases the urge to consume more and more of the substance in order to achieve the same rewarding effect.

Types of Addictions

- Alcoholism/Tobacco
- Drugs-Opium, Marijuana, Cocaine, Cannabis
- Internet/browsing addiction
- Cell phone addiction
- Video gaming
- Pornography

TYPES OF ADDICTION



Sedatives/Anxiolytics/
inhalants

Hyper obesity/Morbid
obesity

ADDICTION



More about Addiction

- About half the risk for addiction is genetic
- Genes affect the degree of reward that individuals experience when initially using a substance (e.g., drugs) or engaging in certain behaviours (e.g., gambling), as well as the way the body processes alcohol or other drugs.
- Heightened desire to re-experience use of the substance or behaviour, potentially influenced by psycho- logical(e.g., stress, history of trauma), social (e.g., family or friends & use of a substance), and environ- mental factors (e.g., accessibility of a substance, low cost) can lead to regular use/exposure, with chronic use/exposure leading to brain changes
- All addictions are part of mental disorders!

Reasons for taking drugs

- To feel good — feeling of pleasure, “high” or "intoxication". Euphoria
- To feel better — relieve stress, forget problems, or feel numb
- To do better — improve performance or thinking
- Curiosity and peer pressure or experimenting

Factors Leading to Addiction



Symptoms of substance use disorder

- . Impaired control: a craving or strong urge to use the substance; desire or failed attempts to cut down or control substance use
- . Social problems: substance use causes failure to complete major tasks at work, school or home; social, work or leisure activities are given up or cut back because of substance use
- . Risky use: substance is used in risky settings; continued use despite known problems
- . Spending a good deal of time getting the drug, using the drug or recovering from the effects of the drug.
- . Experiencing withdrawal symptoms. when you attempt to stop taking the drug.

- Drug effects: Dependence: Involves physiological need for substance; tolerance (need for larger amounts to get the same effect); withdrawal symptoms (different for each substance);
- Feeling that you have to use the drug regularly — daily or even several times a day.
- Having intense urges for the drug that block out any other thoughts.
- Over time, needing more of the drug to get the same effect.

- Failing in your attempts to stop using the drug.
- Taking larger amounts of the drug over a longer period of time than you intended
- Making certain that you maintain a supply of the drug Spending money on the drug, even though you can't afford it.
- Not meeting obligations and work responsibilities, or cutting back on social or recreational activities because of drug use.

- Continuing to use the drug, even though you know
- it's causing problems in your life or causing you physical or psychological harm.
- Doing things to get the drug that you normally wouldn't do, such as stealing.
- Driving or doing other risky activities when you're under the influence of the drug.

Consequences

- Brain changes include alterations in cortical (pre-frontal cortex) and sub-cortical (limbic system) regions involving the neuro-circuitry of reward, motivation, memory, impulse control and judgment.
- This can lead to dramatic increases in cravings for a drug or activity, as well as impairments in the ability to successfully regulate this impulse, despite the knowledge and experience of many consequences related to the addictive behaviour.

Complications

- Getting a communicable disease.
- Other health problems. Drug addiction can lead to a range of both short-term and long-term mental and physical health problems. These depend on what drug is taken.
- While driving more likely do other dangerous activities while under the influence causing accidents.
- People who are addicted to drugs die by suicide.

- Family problems . Behavioral changes may cause marital or family conflict.
- Work issues. Decline in performance at work, absenteeism and eventual loss of employment.
- Problems at school. Academic performance and motivation to excel in school.

- Legal issues. While buying or possessing illegal drugs, stealing to support the drug addiction, driving while under the influence of drugs or alcohol.
- Financial problems.

Differences between addictive people and non addictive people and their behavior with society

Risk Factors	Protective Factors
<i>Aggressive behavior in childhood</i>	<i>Good self-control</i>
<i>Lack of parental supervision</i>	<i>Parental monitoring and support</i>
<i>Poor social skills</i>	<i>Positive relationships</i>
<i>Drug experimentation</i>	<i>Academic competence</i>
<i>Availability of drugs at school</i>	<i>School anti-drug policies</i>
<i>Community poverty</i>	<i>Neighborhood pride</i>

What other factors increase the risk of addiction?

- Early use: Although taking drugs at any age can lead to addiction, research shows that the earlier people begin to use drugs, the more likely they are to develop serious problems.
- This may be due to the harmful effect that drugs can have on the developing brain.
- It also may result from a mix of early social and biological risk factors, including lack of a stable home or family, exposure to physical or sexual abuse, genes, or mental illness.

How to recover from addictions

Treating Substance Use Disorders (SUDs)

- The first step is recognition of the problem
- A formal assessment of symptoms: Medical professional
- Multiple types of treatment: A combination of medication and individual or group therapy is most effective.
- Treatment approaches that address an individual's specific situation and any co-occurring medical, psychiatric, and social problems is optimal for leading to sustained recovery.

- Varies depending on substance and circumstances
- Treating substance use disorder is challenging and includes one or more of the following:
- Acute detoxification,
 - Prevention and management of withdrawal
 - Cessation (or rarely, reduction) of use
 - Maintenance of abstinence

What works

- Cognitive-behavioural therapy: CBT can help addicted patients overcome substance abuse by teaching them to recognize and avoid destructive thoughts and behaviours.
- Motivational interviewing: This therapy technique involves structured conversations that help patients increase their motivation to overcome substance abuse.
- Contingency management. Provide tangible incentives to encourage patients to stay off drugs
- Psychodrama and group therapies
- Alcoholic Anonymous and Narcotic anonymous
- Rehab centers and Therapeutic Communities and other tertiary care centers

13 principles of effective drug addiction treatment

- 1. Addiction is a complex, but treatable, disease that affects brain function and behavior**
- 2. No single treatment is appropriate for everyone.**
- 3. Treatment needs to be readily available.**
- 4. Effective treatment attends to multiple needs of the individual, not just his or her drugabuse.**
- 5. Remaining in treatment for an adequate period of time is critical.**

6. Counselling: individual and/or group —and other behavioral therapies are the most commonly used forms of drug abuse treatment.

7. Medications are an important element of treatment for many patients, especially when combined with counseling and other behavioral therapies.

8. An individual's treatment and services plan must be assessed continually and modified as necessary to ensure it meets his or her changing needs.

9. Many drug-addicted individuals also have other mental disorders.

10. Medically assisted detoxification is only the first stage of addiction treatment and by itself does little to change long-term drug abuse.

11. Treatment does not need to be voluntary to be effective.

12. Drug use during treatment must be monitored continuously, as lapses during treatment do occur.

13. Treatment programs should assess patients for the presence of HIV/AIDS, hepatitis B and C, tuberculosis and other infectious diseases, as well as provide targeted risk-reduction counseling to help patients modify or change behaviors that place them at risk of contracting or

UNIT-4 : **Avoiding risks and harmful habits**

Effects and health hazards from addictions

Health Hazards

- Health issues: Cardiovascular, digestive and renal impairment, strokes, HIV/Aids, Cancer, Hepatitis B and C, Lungs, Fetal Alcohol Syndromes, etc.
- Mental health issues: Anxiety, depression, Schizophrenias, organic mental disorders due to substance abuse, cognitive dysfunction, personality disorder, etc
- Social health issues: Stigma, disrupted families, anti-social behaviours, higher crime records and law & order problems.

Health Effects

The damage caused by substance abuse depends on the type of drug abused and the duration of abuse. The following factors also influence the severity of the problems that follow drug abuse:

Age: The damage is greater among adolescents as well those who are above 50 years of age.

Nutrition status and living conditions: Poor quality and inadequate quantity of food, unhygienic living conditions and poor access to health care increase the problems caused by drug abuse.

Health condition prior to abuse: The damage is less among individuals whose physical condition is good, as compared to individuals whose health was poor even before the onset of addiction.

Genetic loading factor: A family history of a specific medical or psychiatric problem makes the individual more susceptible to the same problems in comparison to another drug abuser who is not genetically prone to the disorder.

Medical and psychiatric complications due to substance abuse:

a) Problems due to intoxication

- Trauma or other physical injury: Mood changes caused by drugs reduce inhibitions as well as increase negative feelings leading to violence.
- Driving vehicles or operating machinery under the influence of alcohol/ drugs causes accidents and injury.
- Poor judgment and poor coordination can cause physical injury or even death.
- Perceptual distortions can also result in accidents.
- Acute intoxication reaction.
- Coma and overdose deaths.

b) Problems due to withdrawal-Physical withdrawal symptoms of varying intensity occur as part of withdrawal.

c) Psychiatric disorders associated with substance abuse-Psychoactive substances can induce psychotic disorders during or immediately after use such as cannabis.

d) Systemic disorders associated with substance abuse

Dermatological complications-

When the same vein is used to inject the drug, repeated injections leave marks on the vein. The discoloration along the vein leaves dark lines called needle tracks.

Tattooing and scarring in IV drug users at injecting site.

Use of infected needles can cause infection at the injection site.

Pruritis and dermatitis

Cardio-vascular complications

- Infection in the heart
- Blocking circulation: Repeated injections at the same site can cause blood clots or air bubbles (emboli) that can block the blood supply in the veins.
- Ventricular fibrillation: This may lead to hemorrhage and sudden death.

Pulmonary complications

- Embolus formation
- Pulmonary oedema
- Aspiration pneumonia
- Pharyngitis, bronchitis, pneumonia and tuberculosis are common infective respiratory diseases among drug abusers.

Hepatic complications

- Hepatitis: Hepatitis B, C & D are blood borne infections that can be transmitted through sharing of needles.
- Alcoholic hepatitis
- Fatty liver
- Alcoholic cirrhosis
- Hepatic trauma

Complications in the reproductive system

- The incidence of sexually transmitted diseases is higher among drug abusers.
- Drug abuse can reduce sex drive and cause impotence.
- Reduced ovulation and menstrual abnormality are seen among female drug abusers.
- Children born to mothers who abuse drugs and alcohol during their pregnancy usually have low birth weight and other problems in their physical and mental development.

Neuromuscular complications

- Cerebrovascular accidents
- Dementia
- Neuritis
- Bacterial meningitis

Haematopoietic complications

- Reduced count of platelets in the blood may lead to bleeding disorders.
- Bone marrow depression: The count of both white and red blood corpuscles may fall.
- Poor nutritional status can cause anemia, especially among alcoholics.

Endocrine system disorders

- Ganja, opiate and alcohol abuse lowers the testosterone levels in the body,
- Incidence of diabetes mellitus increases among alcoholics.

Renal function disorders:

- Damage to renal cells cause nephropathy.


Gastro-intestinal disorders

- Gastritis (inflammation of the lining of the stomach) and peptic ulcers can develop due to poor food habits as well as due to the effect of the drugs abused.
- Acute and chronic pancreatitis.
- The incidence of cancer all along the gastro intestinal tract from the mouth to the anus is higher among alcoholics.


Cancer-

- As most drug and alcohol abusers also abuse tobacco products, the risk of cancer is high.


The Risks of Parental Substance Use Disorders on Children: Alcohol and Illegal Drugs



The Risks of Parental Substance Use Disorders on Children: Alcohol and Illegal Drugs



The Risks of Parental Substance Use Disorders on Children: Prescription Drugs and Pain Medications



(Breshears, 2009; National Institute on Drug Abuse, 2018a)

Diagnosing Substance Use Disorders: DSM-5 Criteria

Severe
6+ Criteria

Moderate
4-5 Criteria

Mild
2-3 Criteria



1. Impaired Control

- Larger amounts or over a longer time than originally intended
- Persistent desire to cut down
- A great deal of time spent obtaining the substance
- Intense craving

2. Social Impairment

- Failure to fulfill work or school obligations
- Recurrent social or interpersonal problems
- Withdraw from social or recreational activities

3. Risky Use

- Recurrent use in situations physically hazardous
- Continued use despite persistent physical or psychological problem that is likely to have been caused or exacerbated by use

4. Pharmacological Criteria

- Tolerance: Need for markedly increased dose to achieve the desired effect
- Withdrawal: Syndrome that occurs when blood or tissue concentrations of a substance decline in an individual who had maintained prolonged heavy use

Overview of the Treatment Process



**Early Identification, Screening,
and Brief Intervention**

Done at earliest point possible



**Comprehensive
Assessment**

Determine extent and severity
of disease



Stabilization

Via medically supervised detoxification,
when necessary



**Timely and Appropriate
Substance Use Disorder Treatment**


Address substance use disorder and co-
occurring issues



**Continuing Care
and Recovery Support**


Help parents sustain recovery, maintain
family safety and stability

(American Society of Addiction Medicine, 2014)



Mental Health Problems (1/2)

- Mental Health Disorders are strongly linked to drug use and dependence.
- Internalizing Disorders (PTSD, Depression, Anxiety disorders, Bipolar disorder)
 - Brain responses are heightened in response to stress.
 - Tendency to self-medicate the anxiety & depression this process causes.



Mental Health Problems (2/2)

- Externalizing Disorders (Conduct Disorder, Attention Deficit Hyperactivity Disorder, Oppositional Defiant Disorder, Antisocial Personality Disorder)
 - Low level of arousal in these disorders is related to an insensitivity to consequences and a need for more stimulation.
 - Heightens risk for continued drug use to relieve symptoms
 - Tend to be resistant to substance abuse treatment
 - Exacerbates stress reactivity problems

Personality & Temperament

- A difficult temperament and certain personality characteristics are consistently related to heightened risk for drug use.
 - Impulsivity
 - Aggressiveness
 - Sensation or novelty-seeking
 - Negative affect
 - Impaired judgment
- High activity level
- Risk taking tendencies
- Lack of regard for negative consequences
- Lack of pain avoidance responses
- Abnormal levels of arousal in response to stress.



Importance of Personality in Adolescence

- Normal adolescence is characterized by greater reward anticipation, sensitivity, and sensation seeking—particularly social rewards (e.g., peer regard, gains in social status).
 - It follows that adolescence is the period during which drug use onset is most common.
 - And, therefore, that adolescents with especially high levels of any combination of these traits are at heightened risk.
- Preventive Implications: These traits can be redirected through psychosocial means to decrease risk for drug use. Prevention programs must be designed to specifically redirect this developmental track.

SAMHSA'S Strategic Prevention Framework

Assessment-collect data to determine the population needs, resources, and gaps.

Capacity-mobilize and/or build up resources to meet goals.

Planning-development of a comprehensive plan.

Implementation-carrying out of the plan, using evidence-based prevention programs.

Evaluation-monitoring the implementation, measuring impact, and determining needed improvements.

Always be mindful of:

Sustainability-process of integrating prevention into ongoing operations.

Cultural Competence-interacting with audiences from diverse backgrounds.



Prevention

WHO has recommended the prevention of substance abuse as primary prevention, early intervention, treatment, care, recovery, rehabilitation and social reintegration measures, along with measures aimed at minimizing the adverse public health and social consequences of drug abuse.

Successful prevention of substance use and risk reduction is thus an essential approach to achieve better public health outcomes in following disorders:

The prevention of substance-induced mental disorders and reductions in injuries and violence (traffic and domestic injuries, child abuse, and gender-based, sexual and other violence), reduction in communicable diseases such as HIV, viral hepatitis and tuberculosis, sexual and reproductive health problems (sexually transmitted infections, unplanned pregnancies and complicated pregnancies) and reduction in noncommunicable diseases (such as cancer, cardiovascular diseases and liver diseases).

Prevention of drug abuse can be achieved by-

i) To take effective and practical primary prevention measures that protect people, in particular children and youth, from drug use initiation:

- by providing them with accurate information about the risks of drug abuse,
- by developing supportive parenting and healthy social environments and
- by promoting skills and opportunities to choose healthy lifestyles and
- by ensuring equal access to education and vocational training;

The family is the strongest factor in protecting children from drug use and other risky behaviour. Families that are happy communicate openly and provide a healthy environment and a positive future for their children.

Schools can cultivate sound values in their students in relation to health and drug use through the development of knowledge and providing them with coping skills to face situations which can lead to drug abuse. It is also necessary to target those who are not in school. Therefore, outreach efforts on the streets are often helpful.

Workplace: Employers can promote the health of employees by preventing substance abuse and assisting those with a drug dependence problem.

ii) To take effective measures to prevent progression to severe drug use disorders through targeted interventions for people at risk for such progression.

iii) Raising awareness of drug-driving and its impact on road safety including deaths, injuries, property damage and harm to other road users among the general public. Policy-makers can reduce drug-driving by enforcing drug driving laws. It is important not only to enforce drug driving laws but also to counsel and, when appropriate, treat drivers found to be impaired by drugs; especially those who are repeat offenders or those with drug-use disorders.

iv) Mass media can play an important role in substance abuse prevention. They can be used to raise awareness concerning not only the dangers of drug abuse but also the dangers of specific practices such as injecting drugs.

National Programmes related to prevention of substance abuse:

Ministry of Health and Family Welfare (MoHFW) and Ministry of Social Justice and Empowerment (MSJE), Government of India are involved with alcohol and drug demand reduction policies and drug de-addiction programme (DDAP) in the country. Under the DDAP de-addiction centres have been established in association with various District Hospitals and psychiatry departments of Medical Colleges. The centre at All India Institute of Medical Sciences, Delhi (AIIMS) has been designated as the “National Drug Dependence Treatment Centre” (NDDTC) and functions as a resource centre for the DDAP of MoHFW, GoI.

Various E- Health web based programmes related to prevention of substance abuse such as alcohol e-Help (alcoholwebindia.in/intervention) operated by NDDTC, AIIMS, Delhi (alcoholwebindia.in/) and m-Cessation programme (nhp.gov.in/quit-tobacco) for tobacco cessation have been initiated by MoHFW, GoI.

National AIDS Control Programme (NACP), MoHFW, GoI has implemented various targeted interventions programme for IDUs such as Needle syringe programme, Opioid substitution therapy.

The MSJE, GoI, has formulated the central sector scheme of assistance for prevention of alcoholism and substance (drugs) abuse and social defence services which has been updated in 2015.

The Department of Revenue, Central government acts as administrator of the Narcotic Drugs and Psychotropic Substances (NDPS) Act, 1985 and the Prevention of Illicit Traffic in Narcotic Drugs and Psychotropic Substances Act, 1988;

The Narcotics Control Bureau, Ministry of Home Affairs, Govt functions as an enforcement agency for NDPS Act.

Tips for prevention of substance abuse: youth

Don't be afraid to say no.

Connect with your friends and avoid negative peer pressure.

Make connections with your parents or other adults.
Get educated about alcohol and drugs.