Dowi

Mark

Switch:

Marketi

Automa

all Char

CONTACT **NEWSLETTER** HOME RESOURCES SALESmanago EARTH HEALTH PHYSICS SCIENCE **TECHNOLOGY**

SEPTEMBER 21, 2019 | TURNING CO2 INTO FUELS, PLASTICS AND

HOME **SCIENCE NEWS**

HOT TOPICS

Neuroscientists Make Major Breakthrough in 200-Year-Old Puzzle – New Psychophysical Law

TOPICS: Behavioral Science

Champalimaud Centre For The Unknown Neuroscience

SUBSCRIBE

SEARCH ...

SciTechDaily: Home of the best science and technology news since 1998. Keep up with the latest scitech news via email or social media.

E-mail









Submit

Time

By CHAMPALIMAUD CENTRE FOR THE UNKNOWN SEPTEMBER 21, 2019









About 200 years ago, the German physician Ernst Heinrich Weber made a seemingly innocuous observation which led to the birth of the discipline of Psychophysics — the science relating physical stimuli in the world and the sensations they evoke in the mind of a subject. Weber asked subjects to say which of two slightly different weights was heavier. From these experiments, he discovered that the probability that a subject will make the right choice only depends on the ratio between the weights.

For instance, if a subject is correct 75% of the time when comparing a weight of 1 Kg and a weight of 1.1 Kg, then she will also be correct 75% of the time when comparing two weights of 2 and 2.2 Kg — or, in



POPULAR ARTICLES



AUGUST 21, 2019

After 127 Years, Physics Riddle Finally Solved

Researcher proves that boat wakes can be skewed. He solved a 127-year-old physics problem on paper and proved that off-centered boat wakes could exist. Five...

general, any pair of weights where one is 10% heavier than the other. This simple but precise rule opened the door to the quantification of behavior in terms of mathematical 'laws'.

READ MORE

Weber's observations have since been generalized to all sensory modalities across many animal species, leading to what is now known as Weber's Law. It is the oldest and most firmly established law in psychophysics. Psychophysical laws describe precise rules of perception and are important because they can be used to obtain mathematical explanations of behavior in terms of brain processes, just like the precise patterns of movement of the planets in the sky were useful to understand gravitation.

Many explanations for Weber's law have been proposed over the years. Although they can all account for Weber's findings, no experimental test had been found to identify which model was correct. Thus, the puzzle of a mathematical explanation of Weber's law remained open.

Now, a team of researchers at the Champalimaud Centre for the Unknown in Lisbon, Portugal, has discovered that Weber's Law can be described as the consequence of a new psychophysical rule involving the time that it takes to make a choice, not just the outcome of the decision. The team has shown that this new rule is sufficient to derive a unique and accurate mathematical model describing the cognitive process underlying Weber's Law. Their results are described in an article published in the scientific journal *Nature Neuroscience*.

Time is key

In this new study, Alfonso Renart, the principal investigator that led the study, and his team, trained rats to discriminate between two sounds of slightly



AUGUST
21, 2019

NASA'S
Spitzer
Space
Telescope
Gets a
Rare Look
at the
Surface
of a
Rocky
Exoplanet



AUGUST 21, 2019 New, Healthier 'Butter' Spread Made Mostly of Water



AUGUST 21, 2019 Fake News Can Lead to False Memories



AUGUST 20, 2019 Merging

different intensities. They constructed minuscule headphones, specially adapted to the head of rats, and used them to deliver sounds simultaneously to both ears.

In each trial, the sound on one of the two speakers would be slightly louder, and the rat's job was to report which speaker played the louder sound by orienting towards the corresponding side. "This behavior is natural to rats, because they orient their heads towards the source of a sound, just like we do," explains José Pardo-Vazquez, one of the article's coauthors. The rats could experience the sound as long as they needed in order to make up their mind. Thus, every attempt provided a choice, and a decision time.

"Our experiments confirmed that the animals' behavior matched Weber's Law," says Pardo-Vazquez. "Their ability to tell which of the two sounds was louder only depended on the ratio between the sounds' intensities. If the rat had to compare the intensities of two sounds that were played softly, its accuracy was just as good as with a pair of sounds that were played loudly, as long as both pairs had the same intensity ratio."

Then, the team started to analyze in detail how long the rats took to make their decisions, a step that turned out to be critical. "Typically, studies of Weber's law focussed on the accuracy of the discrimination, which is what Weber himself described," Pardo-Vazquez explains. "Surprisingly, the time taken to decide has received little attention." The team realized decision times and the loudness of the pair of sounds were linked — the louder the sounds, the shorter the decision time. In fact, they showed that the nature of this link was unique and mathematically precise, making the decision times observed, for instance, in a discrimination between two quiet sounds, exactly proportional to the decision times measured when the

Galaxies Spotted by Hubble Space Telescope



AUGUST 20, 2019 Rocky, Earth-Sized Exoplanet Is Missing an Atmosphe re [Video]



AUGUST
20, 2019

Physicists
Design
Experime
nt to Pin
Down the
Origin of
the
Elements



AUGUST
19, 2019
Black
Hole
Swallowin
g a
Neutron
Star
Detected

subject discriminated between two loud sounds — as long as their relative intensities were constant.

TAGS

Beyond Weber's Law

The team had, in fact, discovered a new "psychophysical law," which they referred to as 'Time-Intensity Equivalence in Discrimination' (TIED), because it linked the overall intensity of a pair of sounds and the time it took to discriminate between them. The TIED is more stringent than Weber's Law, because it not only links the accuracy of pairs of discriminations, it also links their associated decision times. "The precision of this relationship between the decision times in our experiments is amazing," says Pardo-Vazquez, "it is unusual that the behavior of animals can be described with such mathematical precision."

To investigate if the TIED held also in different conditions, the team ran the same kind of experiment with human subjects, obtaining similar results. They also analyzed experiments done by others in which rats perform olfactory discriminations of odor mixtures, again with the same outcome. "It is still early to say if the TIED is as general as Weber's law, but the fact that we obtained the same results in two species and across two sensory modalities is an encouraging first step," Pardo-Vazquez concluded.

Closing in on the right model

Dozens of mathematical models have been proposed through the years to explain Weber's law, but there was no clear experimental test to distinguish them. The researchers reasoned that the TIED offered a way forward. Their analysis revealed that in order to be consistent with the TIED, a mathematical model of the discrimination task would need to satisfy a set of strict conditions. "This was fantastic," says Juan Castiñeiras,

Astronomy

Astrophysics Biology

Black Hole Brain Activity Cancer Cassini-Huygens Mission Cell Biology Chandra X-ray Observatory Chemistry Climate Change

Cosmology Disease

Earth Earth Science Engineering European Southern **Observatory European Space** Agency Evolutionary Biology **Exoplanet Genetics Green** Technology Harvard-Smithsonian Center For **Astrophysics** Harvard University Health **Hubble Space** Telescope Images JPL Mars Materials Science Max Planck Institute Medicine MIT Nanoscience Nanotechnology NASA Neurology Neuroscience Physics Planetary Science Popular

Quantum Physics Saturn UCLA
Yale University

another co-author of the study. "The TIED constrained the world of possible explanations, and thereby resolved the ambiguity between the many proposed models of Weber's law." A previously proposed model by the psychologist Stephen Link at the end of the 80s came close to the solution, but missed an important condition describing how the intensity of sensory stimuli is encoded by the activity of sensory neurons.

The final step was to take this set of conditions, and construct a model to test how accurately it accounted for the behavior of the rats. "We analyzed the simplest model with the fewest possible number of parameters," explains Castiñeiras. When the values of these parameters were chosen to maximize the similarity to the behavior of the rats, they discovered that the fit of the model was remarkable. "Even the simplest model effectively captured everything we could measure with almost no error. This greatly strengthened our confidence that the model describes something true about how perception works," says Renart.

Precise experiments and theories lead to definite progress

These results stand out in their field because of the precision of both the new psychophysical rule and of the mathematical model that describes the experimental data. "Although less frequently observed, in biology and in the study of behavior — like in physics — precise experimental results permit precise explanations which resolve previous ambiguities and therefore constitute progress," says Renart. For instance, their results suggest that one of the main theories in psychophysics was not adequate to describe the TIED. "Producing mathematical explanations that rule out competing theories is very rare in neuroscience, because there is always the possibility to slightly modify one model to make it

compatible with the experimental data," Castiñeiras points out. "We showed that a very influential theory in psychophysics (called Signal Detection Theory) did not model decision times and could therefore not describe the TIED. It missed the essence of the explanation of Weber's law."

One of the next goals of the team is to understand how the mathematical model they have identified is implemented by the brain: "We want to determine which brain areas are important in our task, and how neurons in these circuits carry out the different computational elements of the model," Renart concludes.

Reference: "The mechanistic foundation of Weber's law" by Jose L. Pardo-Vazquez, Juan R. Castiñeiras-de Saa, Mafalda Valente, Iris Damião, Tiago Costa, M. Inês Vicente, André G. Mendonça, Zachary F. Mainen and Alfonso Renart, 12 August 2019, Nature Neuroscience.

DOI: 10.1038/s41593-019-0439-7





MORE ON SCITECHDAILY











SCIENCE Zipf's Law from Scale-free Geometry - Like Violates the Galaxies, Like

SCIENCE **Nanoparticle Temporarily** Second Law of

SCIENCE Electronically **Linked Brains of** Rats Communicate

BIOLOGY **Neuroscientists Identify Brain Circuit That Controls**

Thermodynamic Directly to Solve Decision-Cities Behavioral **Making Under Puzzles** Conflict BIOLOGY **SPACE** Scientists in **Human Stem** New **Neuroscientists New York City Cells Allow Fundamental** Discover Paraplegic Rats Law Stipulates **Fundamental** Use the Evolution of Pheromones to Rule of Brain to Walk and **Track Rats Regain Sensory Galaxy Clusters Plasticity** Perception

BE THE FIRST TO COMMENT

ON "NEUROSCIENTISTS MAKE MAJOR BREAKTHROUGH IN 200-YEAR-OLD PUZZLE – NEW PSYCHOPHYSICAL LAW"

Leave a comment

Email address is optional. If provided, your email will not be published or shared.

Comment			
Name			
Email			

POST COMMENT

Confirm you are NOT a spammer

Mach' den E

Du weißt nicht, wie sind? Dieser Englis

VfbW e.V. Hamburg

FOLLOW SCITECHDAILY	SCITECH NEWS	LATEST NEWS	
Facebook	Biology News	Finally Know How It Works Turning CO2 Into Fuels,	
Twitter	Chemistry News		
Pinterest	Earth News		
Newsletter	Health News	Plastics and Other Valuable Products	
RSS	Physics News	Einstein's General Relativity	

Science News	Validated 10 Years Ahead of Schedule by 'Ringing' Black Hole	
Space News		
Technology News	Where to Park Your Car? Here's the Efficient Way, According to Math. [Video]	
	Understanding the Mysteries of the Cosmos: New Clues to Dark Matter From Primordial Black Holes	

COPYRIGHT © 1998 - 2019 SCITECHDAILY. ALL RIGHTS RESERVED.

EDITORIAL BOARD | SCIENCE NEWS | PRIVACY POLICY TERMS OF USE