c

c

$\frac{2min[\mu_1,\mu_2]}{1+\eta}$.

$-ct$

$V >=$

$\frac{1-\eta}{2}||(y,u)||^2$

$||(y,u)|| <=$

$\sqrt{\left(\frac{1+\eta}{1-\eta}\right)||(y(0),u(0))||e^{-ct}}$

$(x_i(t),_i(t))$

$(x*,u*)$

$\alpha_i(t)$

$\beta_i(t)$

ECS 334: Linear Control Systems

Vedanshi Vaghela & G.V.S Sriram

Spring 2026

1 Abstract

This report studies a new distributed adaptive continuous-time convex optimization algorithm over weight-balanced directed and strongly connceted graphs with adaptive constants. The study can be further extended to weight-unbalanced settings and event-triggered communication among agents. We present the problem formulation, key analytical results, and equilibrium characterization within the convergence analysis. Simulation results are provided to demonstrate the algorithm's behavior and to validate the theoretical findings. These types of algorithms are particularly useful for optimization involving non-convex global cost functions, unknown network connectivity, and nonlinear system dynamics.

2 Introduction

In recent decades, distributed optimization has attracted significant attention due to its wide range of applications, in operations of swarm of robots, sensor networks, resource allocation, and distributed learning. A central problem in this area involves minimizing a global objective function expressed as the sum of local cost functions, where agents rely only on local computation and communication with neighboring nodes. While a large body of work has focused on discrete-time, consensus-based algorithms, continuous-time approaches have emerged as an attractive alternative due to their analytical tractability and the ability to leverage classical stability tools for convergence analysis.

Existing continuous-time distributed optimization algorithms have achieved notable progress, particularly over undirected and weight-balanced directed graphs [3] [2]. However, these methods rely on restrictive assumptions such as global knowledge of network parameters and strong convexity of gradients. These requirements limit their applicability in large-scale or uncertain networks, where connectivity information may be unavailable and objective functions may exhibit nonlinear [1] and/or non-convex behavior [4]. Moreover, the extension to directed communication topologies introduces additional challenges due to asymmetric information flow and coupled system dynamics.

3 Notation

Let $\mathbb{R}$, $\mathbb{R}^n$, and $\mathbb{R}^{n\times m}$ denote the sets of real numbers, real n-dimensional vectors, and real $n \times m$ matrices, respectively. Let $\mathbb{R}_{>0}$ denote the set of positive real numbers. The transpose of a vector or matrix is denoted by $(\cdot)^T$. The identity matrix of dimension n is denoted by I_n, and the all-ones vector in $\mathbb{R}^n$ is denoted by $\mathbf{1}_n$.

For a vector $x \in \mathbb{R}^n$, $\|x\|$ denotes its Euclidean norm. The ith eigenvalue of a matrix A is denoted by $\lambda_i(A)$. The symbol $\otimes$ denotes the Kronecker product.

For vectors $u_1, \ldots, u_m$, we write $u = [u_1, \ldots, u_m]$ to denote the aggregated vector. In a networked system, local variables associated with agent i are denoted using a superscript; for example, x^i denotes the state of agent i. If $p^i \in \mathbb{R}^d$ is a variable of agent i, then the aggregated (along columns) variable of the N-agent network is written as

$$p = [p^1, \ldots, p^N] \in (\mathbb{R}^d)^N.$$

For convenience, we use $\mathbf{r} \in \mathbb{R}^N$ and $\mathbf{R} \in \mathbb{R}^{N\times(N-1)}$, defined by

$$\mathbf{r} = \frac{1}{\sqrt{N}}\mathbf{1}_N, \qquad \mathbf{r}^T\mathbf{R} = 0, \qquad \mathbf{R}^T\mathbf{R} = I_{N-1}. \tag{3.1}$$

A directed graph is denoted by $\mathcal{G} = (\mathcal{V}, \mathcal{E}, A)$, where $\mathcal{V} = \{1, 2, \ldots, n\}$ is the set of agents, $\mathcal{E}$ is the set of directed edges, and $A = [a_{ij}] \in \mathbb{R}^{n\times n}$ is the weighted adjacency matrix. The Laplacian matrix of the graph is denoted by $L = D - A$, where $D = \mathrm{diag}(d_1, \ldots, d_n)$ is the out-degree matrix with $d_i = \sum_{j=1}^n a_{ij}$.

$$\text{A digraph } G \text{ is weight-balanced} \quad \Longleftrightarrow \quad \big(\mathbf{1}_N^T L = 0\big) \quad \Longleftrightarrow \quad \big(\forall x \in \mathbb{R}^N,\ x^T L x \geq 0\big)$$

Each agent $i \in \mathcal{V}$ is associated with a local cost function $f_i : \mathbb{R}^p \to \mathbb{R}$, and the global optimization problem is to minimize the aggregate cost

$$f(x) = \sum_{i=1}^n f_i(x).$$

The gradient of f_i is denoted by $\nabla f_i(x)$, and the gradient of the global cost is denoted by $\nabla f(x)$.

A set $\Omega \subseteq \mathbb{R}^p$ is said to be convex if, for all $x, y \in \Omega$ and all $\lambda \in [0, 1]$,

$$\lambda x + (1 - \lambda)y \in \Omega.$$

A differentiable function f is said to be strongly convex on a convex set $\Omega \subseteq \mathbb{R}^p$ if there exists $m > 0$ such that

$$(x - y)^T\big(\nabla f(x) - \nabla f(y)\big) \geq m\|x - y\|^2, \quad \forall x, y \in \Omega.$$

A function f is said to be locally Lipschitz if, for every $x \in \mathbb{R}^p$, there exists a neighborhood W of x and a constant $M \geq 0$ such that

$$\|f(y) - f(z)\| \leq M\|y - z\|, \quad \forall y, z \in W.$$

4 Problem Statement

The formulation in this section follows [4]. Consider a set of N agents interacting over a directed communication graph. Each agent $i \in \{1, 2, \ldots, N\}$ is associated with a local cost function

$f_i : \mathbb{R}^n \to \mathbb{R}$. The global cost function is defined as

$$f(z) = \sum_{i=1}^N f_i(z).$$

The objective is to design a continuous-time distributed algorithm such that each agent can solve the optimization problem

$$\min_{z\in\mathbb{R}^n} f(z) \tag{4.1}$$

using only its own information and information received from its neighboring agents.

4.1 Assumptions

The following assumptions are adopted throughout this report.

Assumption 1. The global cost function f is differentiable and strongly convex over $\mathbb{R}^n$. Each local cost function f_i is differentiable, and its gradient is locally Lipschitz on $\mathbb{R}^n$.

Under Assumption 1, the strong convexity of the global cost function guarantees that the optimization problem admits a unique solution.

Note that this assumption does not require each local cost function f_i to be convex. Instead, it suffices that their aggregation satisfies the strong convexity condition mentioned above, which can arise even if some of the individual f_i are nonconvex.

Assumption 2. The communication graph $\mathcal{G}$ is strongly connected.

Assumption 3. The initial condititons of the internal states of the agents sum to 0

4.2 Equivalent Distributed Formulation

Under Assumption 2, the optimization problem can be reformulated in a distributed form as

$$\min_{x_i \in \mathbb{R}^n} \tilde{F}(x) = \sum_{i=1}^{N} f_i(x_i), \quad \text{subject to } (L \otimes I_n)x = 0 \tag{4.2}$$

where x_i is the state of agent i, and

$$x = [x_1^T, x_2^T, \ldots, x_N^T]^T$$

denotes the network state.

The constraint $(L \otimes I_n)x = 0$ represents the consensus condition. In particular,

$$(L \otimes I_n)x = 0 \implies \exists \tau \in \mathbb{R}^n \text{ such that } x = \mathbf{1}_N \otimes \tau.$$

The proof of this equivalence follows the same argument as Lemma 3.1 in [2]. Therefore, all agents agree on a common decision variable, and the original optimization problem is equivalent to the distributed constrained formulation above. This reformulation makes the problem suitable for distributed algorithm design and analysis.

4.3 Project Objectives

The objectives of this project are as follows:

- Study the assumptions and conditions required for the convergence of the distributed optimization algorithm.
- Derive the equilibrium points and establish the convergence proof for the algorithm that solves the stated optimization problem.
- Provide simulation results to support the theoretical analysis.

Report Structure: Section 5 presents the derivation of the equilibrium points and the proof of convergence by considering an appropriate Lyapunov function and showing that the algorithm converges to the equilibrium points. Section 6 presents simulations for a few prominent directed graph topologies, and Section 7 provides an analysis of the simulation results together with concluding remarks.

5 Distributed Adaptive Continuous-Time Convex Optimization Algorithm on Weight-Balanced and Strongly Connected Directed Graphs

In this section, we present a fully distributed algorithm for solving the optimization problem in Equation (2). The algorithm is inspired by the adaptive continuous-time distributed optimization method proposed in [4], with modifications inspired by [3].

For each agent $i \in \{1, 2, \ldots, N\}$, consider the dynamics

$$\dot{v}^i = \gamma_1(\alpha_i + \beta_i) \sum_{j=1}^{N} a_{ij}(x^i - x^j), \tag{5.1}$$

$$\dot{x}^i = -\gamma_2 \nabla f_i(x^i) - \gamma_1(\alpha_i + \beta_i) \sum_{j=1}^{N} a_{ij}(x^i - x^j) - v^i, \tag{5.2}$$

$$\dot{\alpha}_i = \beta_i. \tag{5.3}$$

where

$$\beta_i = e_i^T e_i, \qquad e_i = \sum_{j=1}^{N} a_{ij}(x^i - x^j).$$

Here, $v^i \in \mathbb{R}^n$ denotes the internal state of agent i, e_i denotes the relative consensus error, and $\gamma_1, \gamma_2 \in \mathbb{R}_{>0}$ are control gains. The quantities α_i and β_i act as adaptive coupling gains, with $\alpha_i(0) > 0$.

The algorithm is distributed in the sense that each agent updates its state using only its own information and information received from its neighbors. In the modified form considered here, Equation (5.2) contains just the term v^i instead of

$$\sum_{j=1}^{N} a_{ij}(v^i - v^j).$$

As a result, neighboring agents are not required to exchange their internal states through the communication graph.

The overall network dynamics can be written compactly as

$$\dot{v} = \gamma_1 \big[(\hat{\alpha} + \hat{\beta}) \otimes I_n\big](L \otimes I_n)x, \tag{5.4}$$

$$\dot{x} = -\gamma_2 \tilde{f}(x) - \gamma_1 \big[(\hat{\alpha} + \hat{\beta}) \otimes I_n\big](L \otimes I_n)x - v, \tag{5.5}$$

$$\dot{\hat{\alpha}} = \hat{\beta}, \tag{5.6}$$

where

$$\hat{\alpha} = \mathrm{diag}(\alpha_1, \alpha_2, \ldots, \alpha_N), \qquad \hat{\beta} = \mathrm{diag}\left(\|e_1\|^2, \ldots, \|e_N\|^2\right),$$
$$e = (L \otimes I_n)x,$$
$$v = [(v^1)^T, (v^2)^T, \ldots, (v^N)^T]^T,$$

and

$$\tilde{f}(x) = [(\nabla f_1(x^1))^T, (\nabla f_2(x^2))^T, \ldots, (\nabla f_N(x^N))^T]^T.$$

This compact form is useful for analyzing the equilibrium points and convergence properties of the algorithm in the subsequent sections.

5.1 Equilibrium Points

To determine the equilibrium points, we set

$$\dot{x} = 0, \qquad \dot{v} = 0, \qquad \dot{\hat{\alpha}} = 0,$$

and denote an equilibrium by $(\bar{x}, \bar{v})$.

From (5.6), the condition $\dot{\hat{\alpha}} = 0$ implies $\hat{\beta} = 0$. Since

$$\hat{\beta} = \mathrm{diag}(\|e_1\|^2, \ldots, \|e_N\|^2),$$

we obtain $e_i = 0$ for all i, and therefore

$$(L \otimes I_n)\bar{x} = 0. \tag{5.7}$$

Hence $\bar{x} \in \ker(L \otimes I_n)$, which implies

$$\bar{x} = \mathbf{1}_N \otimes \tilde{\theta}$$

for some $\tilde{\theta} \in \mathbb{R}^n$.

Next, from the equilibrium condition (5.5), we have

$$\bar{v} = -\gamma_2 \nabla \tilde{f}(\bar{x}) \quad \Longleftrightarrow \quad \bar{v}^i = -\gamma_2 \nabla f_i(\bar{x}^i), \quad \forall i \in \{1, \ldots, N\}. \tag{5.8}$$

Summing over i and using Assumption 3 gives

$$\sum_{i=1}^{N} \bar{v}^i = 0 \quad \Longrightarrow \quad \sum_{i=1}^{N} \nabla f_i(\tilde{\theta}) = 0.$$

Equivalently,

$$\nabla \left(\sum_{i=1}^{N} f_i(\tilde{\theta}) \right) = 0. \tag{5.9}$$

By Assumption 1, the global cost function is strongly convex, so its stationary point is unique. Therefore,

$$\tilde{\theta} = x^* = \arg\min_x \sum_{i=1}^{N} f_i(x).$$

Substituting this into the consensus form of $\bar{x}$ yields

$$\bar{x} = \mathbf{1}_N \otimes x^*.$$

Thus, every equilibrium point is a consensus configuration at the global minimizer of (4.2). □

5.2 Proof of Convergence

Theorem 1. Under Assumptions 1 and 2, if the communication graph $\mathcal{G}$ is weight-balanced, then the dynamic algorithm in (5.1)–(5.3) solves the distributed optimization problem in (4.2) for any initial conditions $x^i(0), v^i(0) \in \mathbb{R}^n$ exponentially. Furthermore, the dynamic coupling gains α_i converge to finite steady-state values exponentially.

Proof: The methodology combines ideas from [4] and [3], namely equilibrium shifting and Lyapunov analysis. The proof proceeds step-wise for clarity.

Step 1: Equilibrium Shift. Transfer the equilibrium $(\bar{x}, \bar{v})$ to the origin via

$$\mathbf{u} = v - \bar{v}, \qquad \mathbf{y} = x - \bar{x}, \tag{5.10}$$

so that $\dot{\mathbf{u}} = \dot{v}$ and $\dot{\mathbf{y}} = \dot{x}$ since $\bar{v}$ and $\bar{x}$ are constants.
Step 2: Gradient Deviation. Define for each agent i the local gradient deviation

$$h_i(x^i) := \nabla f_i(x^i) - \nabla f_i(x^*), \tag{5.11}$$

with stack $\mathbf{h}(x) = \tilde{f}(x) - \tilde{f}(x^*)$. Note that $\mathbf{h}(x^*) = \mathbf{0}$ by construction. Introducing $\mathbf{h}$ rather than working directly with $\tilde{f}$ allows the consensus and optimality error components to decouple cleanly in the Lyapunov analysis.
Error Dynamics. Substituting (5.10) into (5.4)–(5.6), and noting that $\bar{x} \in \ker(L \otimes I_n)$ together with (5.11), we obtain

$$\dot{\mathbf{u}} = -\gamma_1 \big[(\hat{\alpha} + \hat{\beta}) \otimes I_n\big](L \otimes I_n)\,\mathbf{y}, \tag{5.12}$$

$$\dot{\mathbf{y}} = -\gamma_2\,\mathbf{h}(\mathbf{y}) - \gamma_1 \big[(\hat{\alpha} + \hat{\beta}) \otimes I_n\big](L \otimes I_n)\,\mathbf{y} - \mathbf{u}, \tag{5.13}$$

$$\dot{\hat{\alpha}} = \hat{\beta}, \tag{5.14}$$

using (5.10)and (5.6), we get the consensus error be

$$\mathbf{e} = (L \otimes I_n)\,\mathbf{y}. \tag{5.15}$$

For ease of notation we use $K = \hat{\alpha} + \hat{\beta}$ and $\mathcal{L} = (L \otimes I_n)$

Step 3: Lyapunov Analysis. Consider the Lyapunov candidate:

$$V(y, u) = \frac{1}{2}\|y\|^2 + \frac{1}{2}\|u\|^2 + \eta\, y^\top u \tag{5.16}$$

As V is positive definite, invoking Young's inequality, we get

$$|y^\top u| \;\le\; \|y\|\,\|u\| \;\le\; \tfrac{1}{2}\|y\|^2 + \tfrac{1}{2}\|u\|^2$$

Hence, if we adjust the value of η in within the open range of -1 to 1, $V(y, u)$ is positive definite. Now, taking the time derivative of V, we get

$$\dot{V} = y^T\dot{y} + u^T\dot{u} + \eta(u^T\dot{y} + y^T\dot{u})$$

Subtituting (5.10) expressions of $\dot{y}$ and $\dot{u}$, we get

$$\dot{V} = -y^T h(y) - y^T K\mathcal{L}y - y^T u + u^T K\mathcal{L}y + \eta y^T K\mathcal{L}Ly - \eta u^T h(y) - \eta u^T K\mathcal{L}y - \eta\|u\|^2 \tag{5.17}$$

Further simplification gives

$$\dot{V} = -y^T h(y) - (1-\eta)y^T K\mathcal{L}y - (1+\eta)\|u\|^2 - \eta u^T h(y) - \eta u^T K\mathcal{L}y \tag{5.18}$$

Now, we bound the individual terms in the above equation. Since each f_i has a Lipschitz gradient, $h(y)$ is Lipschitz. Therefore, by the Cauchy-Schwarz inequality,

$$y^T h(y) \le \|y\|\,\|h(y)\|.$$

Moreover, since the global cost function is strongly convex, we have

$$y^T h(y) \geq M\|y\|^2.$$

Let k_m denote the minimum eigenvalue of K, and let l denote the algebraic connectivity, i.e., the second-smallest eigenvalue of L. By definition, $k_m \geq 0$ and $k_m = \min_i(\alpha_i + \beta_i)$. Hence,

$$-(1-\eta)y^T K\mathcal{L}y \leq -(1-\eta)lk_m\|y\|^2.$$

For the remaining terms, Young's inequality gives

$$|\eta u^T h(y)| \leq \frac{\eta M}{2}\|y\|^2 + \frac{\eta}{2M}\|u\|^2$$

and

$$|\eta u^T K\mathcal{L}y| \leq \frac{\eta\|L\|\,\|K\|}{2\epsilon}\|y\|^2 + \frac{\eta\epsilon}{2}\|u\|^2.$$

umptions, as f_i is strongly Lipshitz, we geth t——$^2 Foranyeq > 0$ to be chosen. Combining all the above bounds, we get

$$\dot{V} <= -\left[M + (1-\eta)lk_m - \frac{\eta M}{2bothth} - \frac{\eta||K||||L||}{2\epsilon}\right]||x||^2 - \left[(1+\eta) - \frac{\eta}{2M} - \frac{\eta\epsilon}{2}\right]||u||^2 \tag{5.19}$$

In the above equation we adjust the following parameters to ensure the negative definiteness of $\dot{V}$. We set $\epsilon = \frac{1}{M}$ to the balance the coefficient of u. We shall choose η small enough such that both the coefficients are positive.

$$\eta < \frac{2m}{M(1+||L||||K||)}$$

With the above choices, there exists $\mu_1, \mu_2 > 0$ such that $\dot{V} <= -\mu_1||y||^2 - \mu_2||u||^2 <= -cV$ for some constant

6 Simulations

The simulations were implemented in MATLAB; the source code is available here.

Simulation 1. Four agents with two-dimensional states interact over a square graph with heterogeneous local objectives. Using $T = 100$ and $dt = 0.001$, the states converge to a common optimizer.



(a) State trajectories (b) Pairwise disagreement (c) Evolution of x_1 and x_2

Figure 1: Simulation 1 results.

Simulation 2. Four agents with two-dimensional states interact over the same graph, but with nonlinear local objectives including Rosenbrock, double-well, sinusoidal, and coupled quadratic terms. Using $T = 200$ and $dt = 0.001$, the states still converge to a common consensus point.



(a) State trajectories (b) Pairwise disagreement (c) Evolution of x_1 and x_2

Figure 2: Simulation 2 results.

7 Conclusion

By the proof of convergence produced above the dynamics in (5.4)–(5.6) solves the distributed problem (4.2), and the simuations validate the observations.

References

[1] Agniva Banerjee, Ashish Shukla, Yashaswini Mathur, and Arijit Sen. Continuous-time adaptive distributed optimization driven geometric-mean within a weight-unbalanced network. pages 37–42, 12 2025.

[2] Bahman Gharesifard and Jorge Cortes. Distributed continuous-time convex optimization on weight-balanced digraphs, 2012.

[3] Solmaz S. Kia, Jorge Cortes, and Sonia Martinez. Distributed convex optimization via continuous-time coordination algorithms with discrete-time communication, 2014.

[4] Zhenhong Li, Zhengtao Ding, Junyong Sun, and Zhongkui Li. Distributed adaptive convex optimization on directed graphs via continuous-time algorithms. *IEEE Transactions on Automatic Control*, 63(5):1434–1441, 2018.