

Arbitrarily many independent observers can share the nonlocality of a single maximally entangled qubit pair

Peter Brown¹
joint work with Roger Colbeck²

Nov 26, 2020


Paper: Phys. Rev. Lett. 125, 090401 (2020) or arXiv:2003.12105

¹ENS de Lyon, France

²University of York, UK


Motivation I

Bell-nonlocality



Motivation I


Bell-nonlocality



- Nonlocal correlations are the foundation for many device independent protocols

Motivation I


Bell-nonlocality



- Nonlocal correlations are the foundation for many device independent protocols
- There are measurements that do not destroy the entanglement between the two halves of the state.

Motivation I


Bell-nonlocality



- Nonlocal correlations are the foundation for many device independent protocols
- There are measurements that do not destroy the entanglement between the two halves of the state.
- Can we use this remaining entanglement to generate more nonlocal correlations?

The scenario


We focus on the following scenario introduced in [SGGP15].



- All inputs/outputs are binary - inputs chosen uniformly.

The scenario

We focus on the following scenario introduced in [SGGP15].




- All inputs/outputs are binary - inputs chosen uniformly.
- Quantum state passed to next Bob (but not the input/output information)

$$\rho_{AB^n} = \frac{1}{2} \sum_{b_{n-1}y_{n-1}} (I \otimes F_{b_{n-1}|y_{n-1}}^{1/2}) \rho_{AB^{n-1}} (I \otimes F_{b_{n-1}|y_{n-1}}^{1/2})$$

The scenario

We focus on the following scenario introduced in [SGGP15].




Main question

- All
- Question: Suppose Alice and Bob¹ share the state ρ_{AB^1} . Then what is the maximum number of Bob's that can achieve an expected CHSH violation with Alice?

The scenario II

Previous works:


- [SGGP15]: 2-qubit maximally entangled state + heavily biased inputs \implies unbounded #-violations.
- [SGGP15]: Numerical evidence to suggest that **without** biasing at most **two** Bobs could violate.
- [MMH16]: Give a proof of this latter statement but under a restricted class of measurements.



The scenario II

Previous works:

- [SGGP15]: 2-qubit maximally entangled state + heavily biased inputs \implies unbounded $\#$ -violations.
- [SGGP15]: Numerical evidence to suggest that **without** biasing at most **two** Bobs could violate.
- [MMH16]: Give a proof of this latter statement but under a restricted class of measurements.



In this work we show this statement is in general false.

- We construct an explicit measurement strategy for any $n \in \mathbb{N}$ such that n Bobs can violate.
- Also extend this strategy to a larger class of states including any pure two-qubit state.

The strategy

We consider qubit POVMs of the form $\{M, I - M\}$ where the effects are given by $M = I/2 + \gamma(\cos(\varphi)\sigma_z + \sin(\varphi)\sigma_x)/2$, where $\varphi \in [-\pi, \pi]$ and $\gamma \in [0, 1]$ is the *sharpness*.

The strategy


We consider qubit POVMs of the form $\{M, I - M\}$ where the effects are given by $M = I/2 + \gamma(\cos(\varphi)\sigma_z + \sin(\varphi)\sigma_x)/2$, where $\varphi \in [-\pi, \pi]$ and $\gamma \in [0, 1]$ is the *sharpness*.

Alice's measurements: For $\theta \in (0, \pi/4]$

$$A_{0|0} = \frac{I + \cos(\theta)\sigma_z + \sin(\theta)\sigma_x}{2}$$

and

$$A_{0|1} = \frac{I + \cos(\theta)\sigma_z - \sin(\theta)\sigma_x}{2}.$$



The strategy


We consider qubit POVMs of the form $\{M, I - M\}$ where the effects are given by $M = I/2 + \gamma(\cos(\varphi)\sigma_z + \sin(\varphi)\sigma_x)/2$, where $\varphi \in [-\pi, \pi]$ and $\gamma \in [0, 1]$ is the *sharpness*.

Alice's measurements: For $\theta \in (0, \pi/4]$

$$A_{0|0} = \frac{I + \cos(\theta)\sigma_z + \sin(\theta)\sigma_x}{2}$$

and

$$A_{0|1} = \frac{I + \cos(\theta)\sigma_z - \sin(\theta)\sigma_x}{2}.$$



Bob^k's measurements: For $\gamma_k \in (0, 1)$

$$B_{0|0}^k = \frac{I + \sigma_z}{2}$$

and

$$B_{0|1}^k = \frac{I + \gamma_k \sigma_x}{2}.$$

The strategy II

If Alice and Bob¹ start with the state $|\psi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ then

$$\text{CHSH}_{AB^k} = 2^{2-k} \left(\gamma_k \sin(\theta) + \cos(\theta) \prod_{j=1}^{k-1} \left(1 + \sqrt{1 - \gamma_j^2} \right) \right).$$

The strategy II

If Alice and Bob¹ start with the state $|\psi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ then

$$\text{CHSH}_{AB^k} = 2^{2-k} \left(\gamma_k \sin(\theta) + \cos(\theta) \prod_{j=1}^{k-1} \left(1 + \sqrt{1 - \gamma_j^2} \right) \right).$$

Theorem

For any $n \in \mathbb{N}$ there exists $\theta \in (0, \pi/4]$ and $(\gamma_1, \dots, \gamma_n) \in (0, 1)^n$ such that $\text{CHSH}_{AB^k} > 2$ for all $1 \leq k \leq n$.

The strategy II

If Alice and Bob¹ start with the state $|\psi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ then

$$\text{CHSH}_{AB^k} = 2^{2-k} \left(\gamma_k \sin(\theta) + \cos(\theta) \prod_{j=1}^{k-1} \left(1 + \sqrt{1 - \gamma_j^2} \right) \right).$$

Theorem

For any $n \in \mathbb{N}$ there exists $\theta \in (0, \pi/4]$ and $(\gamma_1, \dots, \gamma_n) \in (0, 1)^n$ such that $\text{CHSH}_{AB^k} > 2$ for all $1 \leq k \leq n$.

Sketch:

$$\text{CHSH}_{AB^k} > 2 \iff \gamma_k > \frac{2^{k-1} - \cos(\theta) \prod_{j=1}^{k-1} (1 + \sqrt{1 - \gamma_j^2})}{\sin(\theta)}$$

So for $\epsilon > 0$ set

$$\gamma_k := \begin{cases} (1 + \epsilon) \frac{2^{k-1} - \cos(\theta) \prod_{j=1}^{k-1} (1 + \sqrt{1 - \gamma_j^2})}{\sin(\theta)} & \text{if } 0 \leq \gamma_{k-1} \leq 1 \\ \infty & \text{otherwise} \end{cases}$$

Show you can always choose θ small enough such that $0 < \gamma_1 < \dots < \gamma_n < 1$.

Summary/Further work

- Shown unbounded violations for scenario introduced in [SGGP15].

Summary/Further work

- Shown unbounded violations for scenario introduced in [SGGP15].
- Strategy can also be extended to larger class of states including all two-qubit pure states.

Summary/Further work

- Shown unbounded violations for scenario introduced in [SGGP15].
- Strategy can also be extended to larger class of states including all two-qubit pure states.
- Many works have investigated limitations in the sequential scenario and found **strong** limitations:
 - Steering: [SHDH⁺19, SDMM18]
 - Entanglement witnessing: [BMSS18]
 - Other Bell-inequalities: [KP19, DGS⁺19]
 - Tripartite settings: [SDS⁺19, MDG⁺20]

However the analyses use a restricted class of measurements as in [MMH16]. Worth rethinking some of the results within a more general measurement scheme.

Summary/Further work

- Shown unbounded violations for scenario introduced in [SGGP15].
- Strategy can also be extended to larger class of states including all two-qubit pure states.
- Many works have investigated limitations in the sequential scenario and found **strong** limitations:
 - Steering: [SHDH⁺19, SDMM18]
 - Entanglement witnessing: [BMSS18]
 - Other Bell-inequalities: [KP19, DGS⁺19]
 - Tripartite settings: [SDS⁺19, MDG⁺20]

However the analyses use a restricted class of measurements as in [MMH16]. Worth rethinking some of the results within a more general measurement scheme.

- Can we translate sequential schemes into a practical advantage?

Summary/Further work

- Shown unbounded violations for scenario introduced in [SGGP15].
- Strategy can also be extended to larger class of states including all two-qubit pure states.
- Many works have investigated limitations in the sequential scenario and found **strong** limitations:
 - Steering: [SHDH⁺19, SDMM18]
 - Entanglement witnessing: [BMSS18]
 - Other Bell-inequalities: [KP19, DGS⁺19]
 - Tripartite settings: [SDS⁺19, MDG⁺20]

However the analyses use a restricted class of measurements as in [MMH16]. Worth rethinking some of the results within a more general measurement scheme.

- Can we translate sequential schemes into a practical advantage?
- Scenario where we also have a sequence of Alices?

Bibliography



Anindita Bera, Shiladitya Mal, Aditi Sen(De), and Ujjwal Sen.

Witnessing bipartite entanglement sequentially by multiple observers.

Physical Review A, 98:062304, 2018.



Debarshi Das, Arkaprabha Ghosal, Souradeep Sasmal, Shiladitya Mal, and A. S. Majumdar.

Facets of bipartite nonlocality sharing by multiple observers via sequential measurements.

Physical Review A, 99:022305, 2019.



Asmita Kumari and A. K. Pan.

Sharing nonlocality and nontrivial preparation contextuality using the same family of bell expressions.

Physical Review A, 100:062130, 2019.



Ananda G. Maity, Debarshi Das, Arkaprabha Ghosal, Arup Roy, and A. S. Majumdar.

Detection of genuine tripartite entanglement by multiple sequential observers.

Physical Review A, 101:042340, 2020.



Shiladitya Mal, Archan Majumdar, and Dipankar Home.

Sharing of nonlocality of a single member of an entangled pair of qubits is not possible by more than two unbiased observers on the other wing.

Mathematics, 4:48, 2016.



Souradeep Sasmal, Debarshi Das, Shiladitya Mal, and A. S. Majumdar.

Steering a single system sequentially by multiple observers.

Physical Review A, 98:012305, 2018.



Sutapa Saha, Debarshi Das, Souradeep Sasmal, Debasis Sarkar, Kaushiki Mukherjee, Arup Roy, and Some Sankar Bhattacharya.

Sharing of tripartite nonlocality by multiple observers measuring sequentially at one side.

Quantum Information Processing, 18:42, 2019.



Ralph Silva, Nicolas Gisin, Yelena Guryanova, and Sandu Popescu.

Multiple observers can share the nonlocality of half of an entangled pair by using optimal weak measurements.

Physical Review Letters, 114:250401, 2015.




Akshata Shenoy H., Sébastien Designolle, Flavien Hirsch, Ralph Silva, Nicolas Gisin, and Nicolas Brunner.

Unbounded sequence of observers exhibiting Einstein-Podolsky-Rosen steering.

Physical Review A, 99:022317, 2019.

Unbounded violations don't scale so well

We have to pick smaller and smaller θ to allow more Bob's to violate.
Analytical and numerical evidence to suggest θ must decrease double exponentially fast with n .



Pretty bad for the CHSH violations. . .

$$\text{CHSH}_{AB^n} < 2 + 2^{2-n}\theta.$$