PHENIX PWG Meeting

Run 15 pp J/ψ Multiplicity Analysis

PHENIX HI PWG Meeting

Zhaozhong Shi

Los Alamos National Laboratory

02/03/2022





Overview

- Implement the unbinned Roofit to extract the raw yield from the fits to the invariant mass of distribution of J/ψ
- Validation of the fitting results by doing toy studies
- Report the fit parameters including the J/ ψ signal raw yield and mean peak invariant mass in a differential multiplicity binning
- The codes for J/ψ raw yield extraction can be found at github: https://github.com/MYOMAO/PHENIXJPsiAna






Roofit Framework

New framework based on ROOT to fit on dataset based on the extended maximum likelihood dedicated for high energy and nuclear physics

Reference: https://root.cern/manual/roofit/

Extended Unbinned Maximum Likelihood



Advantages:

- Improve fitting performance compared to the traditional binned
- Remove potential bias due to binning
- Used in CMS heavy flavor physics analysis





Model for J/ψ Mass Fitting

Model Function for Roofit for Nominal Results

Signal

Double crystal ball function with the same mean but different width, alpha, and N to model the tail in the low dimuon mass region:

$$F_s(m_{J/\psi}; \alpha, n, \mu, \sigma) = N \exp(-(x-\mu)^2/2\sigma^2)$$
 and $F_s(x; \alpha, n, \mu, \sigma) = N A(B - (x-\mu)/\sigma)^{-n}$

Background

First order polynomial:

$$F_B(m_{J/\psi}) = ax + b$$

• The model will be directly applied to fit the J/ ψ invariant mass in data

$$F(m_{J/\psi}) = S \cdot F_s + B \cdot F_B$$

To extract the signal raw yield S and background raw yield B


Systematics:

Signal function: vary with double Gaussian and single crystal ball functions Background function: vary with exponential decay and quadratic functions





Fitting Performance for Inclusive MB




- The fits look good from a glance
- Unity pull (mean = 0 and width 1) for both signal yield and J/ψ mean -> good closure for fit -> validate the roofit model and fitting results





Fitting Performance Separated for North




- Again, the fits look good from a glance
- Unity pull (mean = 0 and width 1) for both signal yield and J/ψ mean -> good closure for fit -> validate the roofit model and fitting results





Fitting Performance Separated for South



- Again, good closure of the signal yield and J/ψ mean parameters
- In this case the signal raw yield of South and North do approximately add up compared to the inclusive case
 - South + North = 12249 + 8496= 20745
 - Inclusive = 22579







FVTX Event Multiplicity Distribution



FVTX South Multiplicity Distribution






- Total number of events:49320
- Based on the results, we decide to have a multiplicity binning of [0,2,5,8,12,18]





Fitting Performance for Multiplicity [0,2)





- The fits all look great
- Here the signal region is defined as J/ψ PDG mass ± 0.4 GeV/c²
- S and B are obtained from the fit parameters and integrating the F_s and F_B over the signal region
- The event multiplicitySouth + North = FVTX_N + FVTX_S
- The event multiplicity for North = FVTX_N
- The event multiplicity for South = FVTX_S
- In this case they do not add up because of the multiplicity cut





Fitting Performance for Multiplicity [2,5)







Again, the fits look great





Fitting Performance for Multiplicity [5,8)







Again, the fits look great





Fitting Performance for Multiplicity [8,12)






Again, the fits look great





Fitting Performance for Multiplicity [12,19)




 Marginal signal is observed at this multiplicity bin – very low statistics for either FVTX North or South with >= 12 tracks





Inclusive MB Multiplicity at [12,19)





- The fits look great
- We can see clear signal event at the highest multiplicity bin up to about 18 for 12 <= FVTX_N + FVTX_S <= 18





J/ψ Signal Raw Yield and Mean - Inclusive







• The value of J/ ψ mean peak of the combined FVTX North and South is stable within about 20 MeV/c², ranging from 3.13 GeV/c² to 3.15 GeV/c², close to but systematically above the J/ ψ mass: 3.096 GeV/c² from PDG





J/ψ Signal Raw Yield and Mean - North







• The value of J/ ψ mean peak from the FVTX North is stable within about 60 MeV/c², ranging from 3.14 GeV/c² to 3.20 GeV/c², close to but systematically above the J/ ψ mass: 3.096 GeV/c² from PDG. However, the highest multiplicity bin





J/ψ Signal Raw Yield and Mean - South






• The value of J/ ψ mean peak from the FVTX South is stable within about 100 MeV/c², ranging from 3.14 GeV/c² to 3.04 GeV/c², close to the J/ ψ mass: 3.096 GeV/c² from PDG. However, the highest multiplicity bin





Preliminary J/ψ Signal Raw Yield Ratios




- Need to correct with trigger bias efficiency ratio to obtain the final results
- Overall, no significant dependence on multiplicity
- No significant modification





J/ψ Signal Raw Yield Ratios South and North



- Overall good agreement between south and north (symmetric pp collisions

 parity symmetric)
- Need to look into the mean multiplicity in each bin for the x-axis





To Do List

- Optimize our fitting results with the best mass window
- Complete the systematic studies and estimate the systematic uncertainties of the J/ ψ signal raw yield extraction
- Complete the studies to estimate the trigger efficiency ratios
- Present our final results in abscissae manners rather than a bin format (bin center) for the x-axis (asymmetric X center for the multiplicity in the x-axis)
- Produce the preliminary results for J/ ψ corrected yield ratio as a function of event multiplicity
- Complete the documentation and prepare for the presentation in QM 2022



