

Geometry of Surfaces

5CCM223A/6CCM223B

Video 15

Length of curves on surfaces

Jürgen Berndt

King's College London

Let $\sigma : U \rightarrow \mathbb{R}^3$ be a regular surface patch and

$$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3, \quad t \mapsto \sigma(u(t), v(t))$$

be a curve on the surface $\mathcal{S} = \sigma(U)$. The CHAIN RULE implies

$$\dot{\gamma}(t) = \sigma_u(u(t), v(t))\dot{u}(t) + \sigma_v(u(t), v(t))\dot{v}(t) \quad \cdot = \frac{d}{dt}$$

or briefly

$$\dot{\gamma} = \sigma_u \dot{u} + \sigma_v \dot{v}$$

Then

$$\|\dot{\gamma}\|^2 = (\sigma_u \cdot \sigma_u)\dot{u}^2 + 2(\sigma_u \cdot \sigma_v)\dot{u}\dot{v} + (\sigma_v \cdot \sigma_v)\dot{v}^2 = E\dot{u}^2 + 2F\dot{u}\dot{v} + G\dot{v}^2$$

with

$$E = \sigma_u \cdot \sigma_u, \quad F = \sigma_u \cdot \sigma_v, \quad G = \sigma_v \cdot \sigma_v$$

Let $t_0 \in (\alpha, \beta)$ and consider the arc length function

$$\begin{aligned} s(t) &= \int_{t_0}^t \left\| \frac{d\gamma}{dx} \right\| dx = \int_{t_0}^t \left(E \frac{du^2}{dx^2} + 2F \frac{du}{dx} \frac{dv}{dx} + G \frac{dv^2}{dx^2} \right)^{\frac{1}{2}} dx \\ &= \int_{t_0}^t (Edu^2 + 2Fdudv + Gdv^2)^{\frac{1}{2}} \end{aligned}$$

The expression

$$Edu^2 + 2Fdudv + Gdv^2$$

is called the **first fundamental form** of σ . Since $s(t) = \int_{t_0}^t (ds^2)^{\frac{1}{2}}$, we also write

$$ds^2 = Edu^2 + 2Fdudv + Gdv^2$$

The first fundamental form depends on the parametrization but not on the curve γ

First fundamental form of plane

$$\sigma(u, v) = a + up + vq$$

with $a, p, q \in \mathbb{R}^3$ and p, q linearly independent. Then $\sigma_u = p$ and $\sigma_v = q$. Thus

$$E = \sigma_u \cdot \sigma_u = \|p\|^2, \quad F = \sigma_u \cdot \sigma_v = p \cdot q, \quad G = \sigma_v \cdot \sigma_v = \|q\|^2$$

We can always choose $p, q \in \mathbb{R}^3$ with $\|p\| = 1 = \|q\|$ and $p \cdot q = 0$. Then the first fundamental form of the plane is of the form

$$ds^2 = du^2 + dv^2$$

First fundamental form of sphere

$$\sigma(\theta, \varphi) = (\cos(\theta) \cos(\varphi), \cos(\theta) \sin(\varphi), \sin(\theta))$$

$$\sigma_\theta(\theta, \varphi) = (-\sin(\theta) \cos(\varphi), -\sin(\theta) \sin(\varphi), \cos(\theta))$$

$$\sigma_\varphi(\theta, \varphi) = (-\cos(\theta) \sin(\varphi), \cos(\theta) \cos(\varphi), 0)$$

The coefficients of the first fundamental form are

$$E = \sigma_\theta \cdot \sigma_\theta = 1, \quad F = \sigma_\theta \cdot \sigma_\varphi = 0, \quad G = \sigma_\varphi \cdot \sigma_\varphi = \cos(\theta)^2$$

The first fundamental form of the sphere is

$$ds^2 = d\theta^2 + \cos(\theta)^2 d\varphi^2$$

First fundamental form of generalized cylinder

$$\sigma(u, v) = (f(u), g(u), v)$$

$$\sigma_u(u, v) = (\dot{f}(u), \dot{g}(u), 0)$$

$$\sigma_v(u, v) = (0, 0, 1)$$

The coefficients of the first fundamental form are

$$E = \sigma_u \cdot \sigma_u = \dot{f}^2 + \dot{g}^2, \quad F = \sigma_u \cdot \sigma_v = 0, \quad G = \sigma_v \cdot \sigma_v = 1$$

Since σ is regular, also γ is regular. If we parametrize γ by arc length, then $E = 1$ and the first fundamental form is

$$ds^2 = du^2 + dv^2$$



This is the same expression as for the plane!

Reparametrizations change E, F, G but not ds^2 (Exercise!)

Every surface has a **conformal parametrization**, that is, a parametrization for which $E = G$ and $F = 0$.

Proposition. *A surface parametrization $\sigma : U \rightarrow \mathbb{R}^3$ is conformal if and only if for any two curves $\pi_1 = (u_1, v_1)$, $\pi_2 = (u_2, v_2)$ in U with $\pi_1(t_0) = (a, b) = \pi_2(t_0)$ the angle of intersection of π_1 and π_2 at (a, b) is equal to the angle of intersection of $\gamma_1 = \sigma \circ \pi_1$ and $\gamma_2 = \sigma \circ \pi_2$ at $\sigma(a, b)$*

Briefly: Conformal parametrizations are angle-preserving parametrizations



Proof. Assume σ is conformal. Let θ be the angle between $\dot{\gamma}_1(t_0)$ and $\dot{\gamma}_2(t_0)$. We have $\dot{\gamma}_i = \sigma_u \dot{u}_i + \sigma_v \dot{v}_i$ and

$$\begin{aligned} \cos(\theta) &= \frac{\dot{\gamma}_1 \cdot \dot{\gamma}_2}{\|\dot{\gamma}_1\| \|\dot{\gamma}_2\|}(t_0) \\ &= \frac{E \dot{u}_1 \dot{u}_2 + F(\dot{u}_1 \dot{v}_2 + \dot{v}_1 \dot{u}_2) + G \dot{v}_1 \dot{v}_2}{(E \dot{u}_1^2 + 2F \dot{u}_1 \dot{v}_1 + G \dot{v}_1^2)^{\frac{1}{2}} (E \dot{u}_2^2 + 2F \dot{u}_2 \dot{v}_2 + G \dot{v}_2^2)^{\frac{1}{2}}}(t_0) \quad (*) \end{aligned}$$

Since σ is conformal, we have $E = G$ and $F = 0$ and thus

$$\cos(\theta) = \frac{\dot{u}_1 \dot{u}_2 + \dot{v}_1 \dot{v}_2}{(\dot{u}_1^2 + \dot{v}_1^2)^{\frac{1}{2}} (\dot{u}_2^2 + \dot{v}_2^2)^{\frac{1}{2}}} \quad (**)$$

which is the angle between $\dot{\pi}_1(t_0)$ and $\dot{\pi}_2(t_0)$. Thus σ is angle-preserving

Conversely, assume σ is angle-preserving, which means $(*) = (**)$ for all curves π_1, π_2 with $\pi_1(t_0) = (a, b) = \pi_2(t_0)$. Take

$$\pi_1(t) = (a + t, b) , \pi_2(t) = (a, b + t) , t_0 = 0$$

Then $(*) = (**)$ gives $0 = F$. Take

$$\pi_1(t) = (a + t, b + t) , \pi_2(t) = (a + t, b - t) , t_0 = 0$$

Then $(*) = (**)$ gives $0 = E - G$.

Geometry of Surfaces

5CCM223A/6CCM223B

Video 16

Isometries of surfaces

Jürgen Berndt

King's College London

Plane and round cylinder have same first fundamental form.

plane \sim round cylinder

Length of curve remains unchanged when deforming plane to round cylinder

plane $\not\sim$ round sphere

expect different first fundamental form

$\sigma_1 : U_1 \rightarrow \mathbb{R}^3$, $\sigma_2 : U_2 \rightarrow \mathbb{R}^3$ regular surface patches,
 $\mathcal{S}_1 = \sigma_1(U_1)$, $\mathcal{S}_2 = \sigma_2(U_2)$

A map $f : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ is **smooth** if there exists a smooth map
 $F : U_1 \rightarrow U_2$ with $f \circ \sigma_1 = \sigma_2 \circ F$

$$\begin{array}{ccc} U_1 & \xrightarrow{F} & U_2 \\ \sigma_1 \downarrow & & \downarrow \sigma_2 \\ \mathcal{S}_1 & \xrightarrow{f} & \mathcal{S}_2 \end{array}$$

A map $f : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ is a **diffeomorphism** if f is bijective and f, f^{-1}
are smooth maps ($\iff F : U_1 \rightarrow U_2$ bijective and F, F^{-1} smooth)
Note: If $\mathcal{S}_1 = \mathcal{S}_2 = \mathcal{S}$, $f = \text{id}_{\mathcal{S}}$, then F is a reparametrization map

A diffeomorphism $f : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ preserving the length of curves is called an **isometry**

\mathcal{S}_1 and \mathcal{S}_2 are **isometric** (or σ_1 and σ_2 are isometric) if there exists an isometry $f : \mathcal{S}_1 \rightarrow \mathcal{S}_2$. We write **$\mathcal{S}_1 \cong \mathcal{S}_2$** if \mathcal{S}_1 and \mathcal{S}_2 are isometric

The concept of isometric surfaces is independent of reparametrizations

Theorem. *Two surfaces are isometric if and only if they have reparametrizations $\sigma_1 : U_1 \rightarrow \mathbb{R}^3$ and $\sigma_2 : U_2 \rightarrow \mathbb{R}^3$ with the same first fundamental form*

Proof. Let $\tilde{\sigma}_1 : U_1 \rightarrow \mathbb{R}^3$, $\tilde{\sigma}_2 : U_2 \rightarrow \mathbb{R}^3$ be regular surface patches, $S_1 = \tilde{\sigma}_1(U_1)$, $S_2 = \tilde{\sigma}_2(U_2)$. Let $f : S_1 \rightarrow S_2$ be an isometry. There exists a bijective smooth map $F : U_1 \rightarrow U_2$ so that

$$\begin{array}{ccc} U_1 & \xrightarrow{F} & U_2 \\ \tilde{\sigma}_1 \downarrow & & \downarrow \tilde{\sigma}_2 \\ S_1 & \xrightarrow{f} & S_2 \end{array} \qquad \begin{array}{ccc} U = U_1 & \xrightarrow{F} & U_2 \\ \sigma_1 = \tilde{\sigma}_1 \downarrow & \searrow \sigma_2 & \downarrow \tilde{\sigma}_2 \\ S_1 & \xrightarrow{f} & S_2 \end{array}$$

Define $U = U_1$, $\sigma_1 = \tilde{\sigma}_1 : U \rightarrow S_1$, $\sigma_2 = \tilde{\sigma}_2 \circ F : U \rightarrow S_2$. σ_2 is a reparametrization of $\tilde{\sigma}_2$. Consider curves

$$\gamma_1(t) = \sigma_1(u(t), v(t)), \quad \gamma_2(t) = \sigma_2(u(t), v(t)) = (f \circ \sigma_1)(u(t), v(t))$$

Since f is an isometry, γ_1, γ_2 have the same length

$$\int_{t_0}^{t_1} (E_1 \dot{u}^2 + 2F_1 \dot{u}\dot{v} + G_1 \dot{v}^2)^{\frac{1}{2}} dt = \int_{t_0}^{t_1} (E_2 \dot{u}^2 + 2F_2 \dot{u}\dot{v} + G_2 \dot{v}^2)^{\frac{1}{2}} dt$$

Put $u_0 = u(t_0)$, $v_0 = v(t_0)$

$$(u(t), v(t)) = (u_0 + t - t_0, v_0) \implies E_1 = E_2$$

$$(u(t), v(t)) = (u_0, v_0 + t - t_0) \implies G_1 = G_2$$

$$(u(t), v(t)) = (u_0 + t - t_0, v_0 + t - t_0) \implies F_1 = F_2$$

Thus σ_1 and σ_2 have the same first fundamental form

Conversely, assume that σ_1 and σ_2 have the same first fundamental form. Define $f : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ by $f(\sigma_1(u, v)) = \sigma_2(u, v)$. Let

$$\gamma_1(t) = \sigma_1(u(t), v(t)), \gamma_2(t) = \sigma_2(u(t), v(t)) = (f \circ \sigma_1)(u(t), v(t))$$

Length of γ_1, γ_2 is obtained by integrating $(E_2 \dot{u}^2 + 2F_2 \dot{u}\dot{v} + G_2 \dot{v}^2)^{\frac{1}{2}}$, which is the same for both surfaces by assumption. It follows that f preserves the length of curves and hence is an isometry.

plane \cong generalized cylinder \cong generalized cone

A **tangent developable** is the union of tangent lines to a regular curve γ in \mathbb{R}^3 (can assume $\|\dot{\gamma}\| = 1$):

$$\sigma(u, v) = \gamma(u) + v\dot{\gamma}(u)$$


$$\sigma_u(u, v) = \dot{\gamma}(u) + v\ddot{\gamma}(u)$$

$$\sigma_v(u, v) = \dot{\gamma}(u)$$

$$\begin{aligned}(\sigma_u \times \sigma_v)(u, v) &= v\ddot{\gamma}(u) \times \dot{\gamma}(u) = v(\dot{\mathbf{t}} \times \mathbf{t})(u) \\ &= v\kappa(u)(\mathbf{n} \times \mathbf{t})(u) = -v\kappa(u)\mathbf{b}(u)\end{aligned}$$

Thus σ regular if and only if $\kappa > 0$ and $v \neq 0$

The tangent developable of a circular helix



Theorem. *Any tangent developable is isometric to (part of) a plane.*

Proof. $\sigma(u, v) = \gamma(u) + v\dot{\gamma}(u)$

$$\sigma_u(u, v) = \dot{\gamma}(u) + v\ddot{\gamma}(u)$$

$$\sigma_v(u, v) = \dot{\gamma}(u)$$

$$E = (\sigma_u \cdot \sigma_u)(u, v) = \|\dot{\gamma}(u) + v\ddot{\gamma}(u)\|^2 = 1 + v^2\kappa(u)^2$$

$$F = (\sigma_u \cdot \sigma_v)(u, v) = (\dot{\gamma}(u) + v\ddot{\gamma}(u)) \cdot \dot{\gamma}(u) = 1$$

$$G = (\sigma_v \cdot \sigma_v)(u, v) = \|\dot{\gamma}(u)\|^2 = 1$$

$$ds^2 = (1 + v^2\kappa(u)^2)du^2 + 2dudv + dv^2$$

Tangent developable of a plane curve $\tilde{\gamma}$ with curvature κ has the same first fundamental form (tangents fill out parts of a plane).
Thus both surfaces are isometric

One can prove that a surface that is isometric to (part of) a plane is (part of)

- plane
- generalized cylinder
- generalized cone
- tangent developable

Geometry of Surfaces

5CCM223A/6CCM223B

Video 17


Surface area

Jürgen Berndt

King's College London

Let $\sigma : U \rightarrow \mathbb{R}^3$ be a regular surface patch. Let R be a rectangle in U with side lengths Δu and Δv . Then, for small Δu and Δv , the area of $\sigma(R)$ is approximately the area of the parallelogram with sides $\sigma_u \Delta u$ and $\sigma_v \Delta v$:

$$\|\sigma_u \Delta u \times \sigma_v \Delta v\| = \|\sigma_u \times \sigma_v\| \Delta u \Delta v$$



Let $\sigma : U \rightarrow \mathbb{R}^3$ be a regular surface patch and $R \subseteq U$. The area of $\sigma(R)$ is

$$\mathcal{A}_\sigma(R) = \iint_R \|\sigma_u \times \sigma_v\| du dv$$

Let θ be the angle between σ_u and σ_v . Then

$$\begin{aligned} \|\sigma_u \times \sigma_v\|^2 &= \|\sigma_u\|^2 \|\sigma_v\|^2 \underbrace{\sin(\theta)^2}_{1 - \cos(\theta)^2} \\ &= \|\sigma_u\|^2 \|\sigma_v\|^2 - \underbrace{\|\sigma_u\|^2 \|\sigma_v\|^2 \cos(\theta)^2}_{(\sigma_u \cdot \sigma_v)^2} = EG - F^2 \end{aligned}$$

Therefore

$$\mathcal{A}_\sigma(R) = \iint_R (EG - F^2)^{\frac{1}{2}} du dv$$

Proposition. *Area is unchanged by reparametrizations.*

Proof. Let $\sigma : U \rightarrow \mathbb{R}^3$ be a regular surface patch and $\tilde{\sigma} : \tilde{U} \rightarrow \mathbb{R}^3$ be a reparametrization of σ with reparametrization map $\phi : U \rightarrow \tilde{U}$:

$$\tilde{\sigma}(\tilde{u}, \tilde{v}) = \tilde{\sigma}(\phi(u, v)) = \sigma(u, v)$$

The CHAIN RULE gives

$$\sigma_u = \tilde{\sigma}_{\tilde{u}} \frac{\partial \tilde{u}}{\partial u} + \tilde{\sigma}_{\tilde{v}} \frac{\partial \tilde{v}}{\partial u}, \quad \sigma_v = \tilde{\sigma}_{\tilde{u}} \frac{\partial \tilde{u}}{\partial v} + \tilde{\sigma}_{\tilde{v}} \frac{\partial \tilde{v}}{\partial v}$$
$$\sigma_u \times \sigma_v = \underbrace{\left(\frac{\partial \tilde{u}}{\partial u} \frac{\partial \tilde{v}}{\partial v} - \frac{\partial \tilde{u}}{\partial v} \frac{\partial \tilde{v}}{\partial u} \right)}_{\det(J\phi)} \tilde{\sigma}_{\tilde{u}} \times \tilde{\sigma}_{\tilde{v}}$$

Thus

$$\sigma_u \times \sigma_v = \det(J\phi) \tilde{\sigma}_{\tilde{u}} \times \tilde{\sigma}_{\tilde{v}}$$

Altogether,

$$\begin{aligned}\mathcal{A}_\sigma(R) &= \iint_R \|\sigma_u \times \sigma_v\| du dv \\ &= \iint_R |\det(J\phi)| \|\tilde{\sigma}_{\tilde{u}} \times \tilde{\sigma}_{\tilde{v}}\| du dv \\ &= \iint_R \|\tilde{\sigma}_{\tilde{u}} \times \tilde{\sigma}_{\tilde{v}}\| d\tilde{u} d\tilde{v} = \mathcal{A}_{\tilde{\sigma}}(R)\end{aligned}$$

by CHANGE OF VARIABLES FORMULA