

King's College London

UNIVERSITY OF LONDON

This paper is part of an examination of the College counting towards the award of a degree. Examinations are governed by the College Regulations under the authority of the Academic Board.

FOLLOW THE INSTRUCTIONS YOU HAVE BEEN GIVEN ON HOW TO UPLOAD YOUR SOLUTIONS

BSc AND MSci EXAMINATION

6CCM223B GEOMETRY OF SURFACES

SUMMER 2021

TIME ALLOWED: TWO HOURS

THIS PAPER CONSISTS OF TWO SECTIONS, SECTION A AND SECTION B.

SECTION A CONTRIBUTES 45% OF THE TOTAL MARKS FOR THE PAPER.

ANSWER ALL QUESTIONS IN SECTION A.

ALL QUESTIONS IN SECTION B CARRY EQUAL MARKS, BUT IF MORE THAN TWO QUESTIONS ARE ATTEMPTED, THEN ONLY THE BEST TWO WILL COUNT.

YOU MAY CONSULT LECTURE NOTES AND USE A CALCULATOR.

Section A

All ten questions in Section A carry equal marks.

Answer all questions for full marks.

- A 1.** Calculate the arc length of $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$, $t \mapsto (3t - 2, 4t + 5)$ starting at $\gamma(0)$.
- A 2.** Calculate the curvature of $\gamma : \mathbb{R} \rightarrow \mathbb{R}^3$, $t \mapsto (t, t^2, \sin(t))$ at $\gamma(0)$.
- A 3.** Calculate the torsion of $\gamma : \mathbb{R} \rightarrow \mathbb{R}^3$, $t \mapsto (\sin(t), \cos(t), e^t)$ at $\gamma(0)$.
- A 4.** Calculate the tangent plane of the surface $\sigma(u, v) = (u \cos(v), u \sin(v), v)$ at $\sigma(0, 0)$.
- A 5.** Calculate the geodesic curvature of the curve $\gamma(s) = (-\frac{1}{2} \sin(2s), \frac{1}{2}, \frac{1}{2} \cos(2s))$ on the surface $\sigma(u, v) = (u, 2(u^2 + v^2), v)$.
- A 6.** Calculate the first fundamental form of the surface $\sigma(u, v) = (e^u v, e^{-u} v, u)$.
- A 7.** Calculate the second fundamental form of the surface $\sigma(u, v) = (u + v, u - v, uv)$ at $\sigma(0, 0)$.
- A 8.** Calculate the principal curvatures of the surface \mathcal{S} at $p \in \mathcal{S}$ whose coefficients of the first and second fundamental form at p are given by $E = 2$, $F = 2$, $G = 4$, $L = 4$, $M = 1$, $N = 2$.
- A 9.** Calculate the Gaussian curvature of the surface $\sigma(u, v) = (u^3 v, v, u - v^2)$ at $\sigma(1, 0)$.
- A 10.** Calculate the mean curvature of the surface $\sigma(u, v) = (e^u v, \sin(u)v, u \cos(v))$ at $\sigma(0, 0)$.

Section B

All three questions in Section B carry equal marks.
Answer TWO questions for full marks.

B 11. (i) Show that the smooth curve

$$\gamma : \mathbb{R} \rightarrow \mathbb{R}^3, \quad t \mapsto (t^3, t^6, t^9)$$

is not regular. Find a regular curve with the same image as γ .

(ii) Find a unit speed reparametrization of the curve

$$\gamma : \mathbb{R} \rightarrow \mathbb{R}^3, \quad t \mapsto (4t, 4 \cosh(t), 3 \sinh(t)).$$

(iii) Let $\gamma : \mathbb{R} \rightarrow \mathbb{R}^3$ be a unit speed curve with constant curvature $\kappa > 0$. Assume that $\gamma(\mathbb{R})$ is contained in a plane. Use the FRENET-SERRET EQUATIONS to prove that $\gamma(\mathbb{R})$ lies on a circle.

B 12. (i) Show that

$$\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3, (u, v) \mapsto (u + \cosh(v), u + \sinh(v), 1 + 2ue^{-v})$$

is a regular surface patch parametrizing a part of a hyperbolic paraboloid. Determine this part explicitly.

(ii) Let $U = \mathbb{R} \times (-\frac{\pi}{2}, \frac{\pi}{2})$ and

$$\sigma : U \rightarrow \mathbb{R}^3, (u, v) \mapsto (u \cos(v), u \sin(v), \ln(\cos(v)) + u).$$

Let $v_0 \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $a, b \in \mathbb{R}$ with $a < b$, and define the curve

$$\gamma_{v_0} : (a, b) \rightarrow \mathbb{R}^3, t \mapsto \sigma(t, v_0).$$

Prove that the length of γ_{v_0} does not depend on v_0 .

(iii) Let

$$\sigma : U \rightarrow \mathbb{R}^3, (u, v) \mapsto \sigma(u, v)$$

be a regular surface patch and assume that the first fundamental form of σ is of the form

$$ds^2 = du^2 + g(u)dv^2$$

with some smooth function $g(u)$. Write down the geodesic equations for a unit speed curve

$$\gamma(t) = \sigma(u(t), v(t))$$

on σ and use them to determine the geodesic $\gamma(t)$ for the initial values

$$\gamma(0) = (u_0, v_0) \in U \text{ and } \dot{\gamma}(0) = \frac{\sigma_u(u_0, v_0)}{\|\sigma_u(u_0, v_0)\|}.$$

- B 13.** (i) Let $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^3$ be a unit speed curve, $U = (\alpha, \beta) \times (-\pi, \pi)$, $r \in \mathbb{R}^+$. Assume that the curvature κ of γ satisfies $0 < \kappa < \frac{1}{r}$. Define

$$\sigma : U \rightarrow \mathbb{R}^3, (u, v) \mapsto \gamma(u) + r(\cos(v)\mathbf{n}(u) + \sin(v)\mathbf{b}(u)),$$

where \mathbf{n} is the principal normal and \mathbf{b} is the binormal of γ . Assume that σ is a regular surface patch. Compute the Gauss map of the surface and show that the curve

$$\alpha(t) = \sigma(u_0, \frac{t}{r}) \quad ((u_0, \frac{t}{r}) \in U)$$

is a geodesic on the surface.

- (ii) Let $\sigma_i : U_i \rightarrow \mathbb{R}^3$, $i \in \{1, 2\}$, be two regular surface patches with Gaussian curvature K_i and mean curvature H_i . Are the follow statements true or false? In each case, justify your answer.
- (a) If the two surfaces are isometric to each other, then $K_1 = K_2$;
 - (b) If the two surfaces are isometric to each other, then $H_1 = H_2$.
- (iii) Consider a triangulation of the sphere S^2 with 12 vertices and 20 faces. How many edges are there in this triangulation? Justify your answer.

Solutions

For each question I state one possible solution that is based on the material taught in the course. For some questions, in particular proofs, there are of course other solutions for which a student can get full marks.

A 1. To get the arc length function $s(t)$ starting at $\gamma(0)$, we calculate

$$\begin{aligned}\gamma(t) &= (3t - 2, 4t + 5) \\ \gamma'(t) &= (3, 4) \\ \|\gamma'(t)\| &= \sqrt{3^2 + 4^2} = \sqrt{25} = 5 \\ s(t) &= \int_0^t \|\gamma'(u)\| du = \int_0^t 5 du = 5t\end{aligned}$$

A 2. To get the curvature $\kappa(0)$ of γ at $\gamma(0)$, we calculate

$$\begin{aligned}\gamma(t) &= (t, t^2, \sin(t)) \\ \gamma'(t) &= (1, 2t, \cos(t)) \\ \gamma''(t) &= (0, 2, -\sin(t)) \\ \gamma'(0) \times \gamma''(0) &= (1, 0, 1) \times (0, 2, 0) = (-2, 0, 2) \\ \kappa(0) &= \frac{\|\gamma'(0) \times \gamma''(0)\|}{\|\gamma'(0)\|^3} = \frac{\|(-2, 0, 2)\|}{\|(1, 0, 1)\|^3} = \frac{\sqrt{8}}{\sqrt{2}^3} = 1\end{aligned}$$

A 3. To get the torsion $\tau(0)$ of γ at $\gamma(0)$, we calculate

$$\begin{aligned}\gamma(t) &= (\sin(t), \cos(t), e^t) \\ \gamma'(t) &= (\cos(t), -\sin(t), e^t) \\ \gamma''(t) &= (-\sin(t), -\cos(t), e^t) \\ \gamma'''(t) &= (-\cos(t), \sin(t), e^t) \\ \gamma'(0) \times \gamma''(0) &= (1, 0, 1) \times (0, -1, 1) = (1, -1, -1) \\ \gamma'''(0) &= (-1, 0, 1) \\ \tau(0) &= \frac{(\gamma'(0) \times \gamma''(0)) \cdot \gamma'''(0)}{\|\gamma'(0) \times \gamma''(0)\|^2} = \frac{-2}{3} = -\frac{2}{3}\end{aligned}$$

- A 4.** To get the tangent plane $T_{\sigma(0,0)}\mathcal{S}$ of the surface \mathcal{S} given by $\sigma(u, v) = (u \cos(v), u \sin(v), v)$ at $\sigma(0, 0)$, we calculate

$$\begin{aligned}\sigma_u(u, v) &= (\cos(v), \sin(v), 0) \\ \sigma_v(u, v) &= (-u \sin(v), u \cos(v), 1) \\ \sigma_u(0, 0) &= (1, 0, 0) \\ \sigma_v(0, 0) &= (0, 0, 1) \\ T_{\sigma(0,0)}\mathcal{S} &= \{(x, 0, z) \in \mathbb{R}^3 : x, z \in \mathbb{R}\}\end{aligned}$$

- A 5.** To get the geodesic curvature $\kappa_g(s)$ of $\gamma(s) = (-\frac{1}{2} \sin(2s), \frac{1}{2}, \frac{1}{2} \cos(2s))$ on the surface $\sigma(u, v) = (u, 2(u^2+v^2), v)$, we observe that $\gamma(s) = \sigma(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s))$ and calculate

$$\begin{aligned}\dot{\gamma}(s) &= (-\cos(2s), 0, -\sin(2s)) \text{ (thus } \gamma \text{ is unit speed)} \\ \ddot{\gamma}(s) &= (2 \sin(2s), 0, -2 \cos(2s)) \\ \sigma_u(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s)) &= (1, -2 \sin(2s), 0) \\ \sigma_v(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s)) &= (0, 2 \cos(2s), 1) \\ (\sigma_u \times \sigma_v)(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s)) &= (-2 \sin(2s), -1, 2 \cos(2s)) \\ \mathbf{N}(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s)) &= \frac{1}{\sqrt{5}}(-2 \sin(2s), -1, 2 \cos(2s)) \\ \mathbf{N}(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s)) \times \dot{\gamma}(s) &= \frac{1}{\sqrt{5}}(\sin(2s), -2, -\cos(2s)) \\ \kappa_g(s) &= \ddot{\gamma}(s) \cdot (\mathbf{N}(-\frac{1}{2} \sin(2s), \frac{1}{2} \cos(2s)) \times \dot{\gamma}(s)) = \frac{2}{\sqrt{5}}\end{aligned}$$

- A 6.** To get the first fundamental form ds^2 of the surface $\sigma(u, v) = (e^u v, e^{-u} v, u)$, we calculate

$$\begin{aligned}\sigma_u(u, v) &= (e^u v, -e^{-u} v, 1) \\ \sigma_v(u, v) &= (e^u, e^{-u}, 0) \\ E(u, v) &= \sigma_u(u, v) \cdot \sigma_u(u, v) = e^{2u} v^2 + e^{-2u} v^2 + 1 = 1 + 2 \cosh(2u) v^2 \\ F(u, v) &= \sigma_u(u, v) \cdot \sigma_v(u, v) = e^{2u} v - e^{-2u} v = 2 \sinh(2u) v \\ G(u, v) &= \sigma_v(u, v) \cdot \sigma_v(u, v) = e^{2u} + e^{-2u} = 2 \cosh(2u) \\ ds^2 &= (1 + 2 \cosh(2u) v^2) du^2 + 4 \sinh(2u) v du dv + 2 \cosh(2u) dv^2\end{aligned}$$

A 7. To get the second fundamental form $II(0,0)$ of $\sigma(u,v) = (u+v, u-v, uv)$ at $\sigma(0,0)$, we calculate

$$\begin{aligned}
\sigma_u(u,v) &= (1, 1, v) \\
\sigma_v(u,v) &= (1, -1, u) \\
\sigma_u(0,0) \times \sigma_v(0,0) &= (1, 1, 0) \times (1, -1, 0) = (0, 0, -2) \\
\mathbf{N}(0,0) &= (0, 0, -1) \\
\sigma_{uu}(u,v) &= (0, 0, 0) \\
\sigma_{uv}(u,v) &= (0, 0, 1) \\
\sigma_{vv}(u,v) &= (0, 0, 0) \\
L(0,0) &= \sigma_{uu}(0,0) \cdot \mathbf{N}(0,0) = 0 \\
M(0,0) &= \sigma_{uv}(0,0) \cdot \mathbf{N}(0,0) = -1 \\
N(0,0) &= \sigma_{vv}(0,0) \cdot \mathbf{N}(0,0) = 0 \\
II(0,0) &= 2M(0,0)dudv = -2dudv
\end{aligned}$$

A 8. To get the principal curvatures of \mathcal{S} at p , we calculate

$$\begin{aligned}
0 &= \det \begin{pmatrix} L - \kappa E & M - \kappa F \\ M - \kappa F & N - \kappa G \end{pmatrix} = \det \begin{pmatrix} 4 - 2\kappa & 1 - 2\kappa \\ 1 - 2\kappa & 2 - 4\kappa \end{pmatrix} \\
&= (4 - 2\kappa)(2 - 4\kappa) - (1 - 2\kappa)^2 = 8 - 16\kappa - 4\kappa + 8\kappa^2 - 1 + 4\kappa - 4\kappa^2 \\
&= 4\kappa^2 - 16\kappa + 7 = 4 \left(\kappa^2 - 4\kappa + \frac{7}{4} \right) = 4 \left((\kappa - 2)^2 - \frac{9}{4} \right)
\end{aligned}$$

which has solutions $2 \pm \frac{3}{2}$. Thus the principal curvatures of \mathcal{S} at p are $\frac{1}{2}$ and $\frac{7}{2}$.

A 9. To get the Gaussian curvature $K(1, 0)$ of the surface $\sigma(u, v) = (u^3v, v, u - v^2)$ at $\sigma(1, 0)$, we calculate

$$\begin{aligned}
\sigma_u(u, v) &= (3u^2v, 0, 1) \\
\sigma_v(u, v) &= (u^3, 1, -2v) \\
(\sigma_u \times \sigma_v)(1, 0) &= (0, 0, 1) \times (1, 1, 0) = (-1, 1, 0) \\
\mathbf{N}(1, 0) &= \frac{1}{\sqrt{2}}(-1, 1, 0) \\
\sigma_{uu}(u, v) &= (6uv, 0, 0) \\
\sigma_{uv}(u, v) &= (3u^2, 0, 0) \\
\sigma_{vv}(u, v) &= (0, 0, -2) \\
E(1, 0) &= \sigma_u(1, 0) \cdot \sigma_u(1, 0) = 1 \\
F(1, 0) &= \sigma_u(1, 0) \cdot \sigma_v(1, 0) = 0 \\
G(1, 0) &= \sigma_v(1, 0) \cdot \sigma_v(1, 0) = 2 \\
L(1, 0) &= \sigma_{uu}(1, 0) \cdot \mathbf{N}(1, 0) = 0 \\
M(1, 0) &= \sigma_{uv}(1, 0) \cdot \mathbf{N}(1, 0) = -\frac{3}{\sqrt{2}} \\
N(1, 0) &= \sigma_{vv}(1, 0) \cdot \mathbf{N}(1, 0) = 0 \\
K(1, 0) &= \frac{LN - M^2}{EG - F^2}(1, 0) = -\frac{9}{4}
\end{aligned}$$

A 10. To get the mean curvature $H(0, 0)$ of the surface $\sigma(u, v) = (e^u v, \sin(u)v, u \cos(v))$ at $\sigma(0, 0)$, we calculate

$$\begin{aligned}
\sigma_u(u, v) &= (e^u v, \cos(u)v, \cos(v)) \\
\sigma_v(u, v) &= (e^u, \sin(u), -u \sin(v)) \\
(\sigma_u \times \sigma_v)(0, 0) &= (0, 0, 1) \times (1, 0, 0) = (0, 1, 0) \\
\mathbf{N}(0, 0) &= (0, 1, 0) \\
\sigma_{uu}(u, v) &= (e^u v, -\sin(u)v, 0) \\
\sigma_{uv}(u, v) &= (e^u, \cos(u), -\sin(v)) \\
\sigma_{vv}(u, v) &= (0, 0, -u \cos(v)) \\
E(0, 0) &= \sigma_u(0, 0) \cdot \sigma_u(0, 0) = 1 \\
F(0, 0) &= \sigma_u(0, 0) \cdot \sigma_v(0, 0) = 0 \\
G(0, 0) &= \sigma_v(0, 0) \cdot \sigma_v(0, 0) = 1 \\
L(0, 0) &= \sigma_{uu}(0, 0) \cdot \mathbf{N}(0, 0) = 0 \\
M(0, 0) &= \sigma_{uv}(0, 0) \cdot \mathbf{N}(0, 0) = 1 \\
N(0, 0) &= \sigma_{vv}(0, 0) \cdot \mathbf{N}(0, 0) = 0 \\
H(0, 0) &= \frac{LG - 2MF + NE}{2(EG - F^2)}(0, 0) = \frac{0}{2} = 0
\end{aligned}$$

- B 11.** (i) We have $\gamma(t) = (t^3, t^6, t^9)$ and $\gamma'(t) = (3t^2, 6t^5, 9t^8)$. Then $\gamma'(0) = (0, 0, 0)$ and hence γ is not regular. Consider the curve

$$\tilde{\gamma} : \mathbb{R} \rightarrow \mathbb{R}^3, \quad t \mapsto (t, t^2, t^3)$$

Then $\tilde{\gamma}'(t) = (1, 2t, 3t^2) \neq 0$ and hence $\tilde{\gamma}$ is regular [4 marks]. Furthermore, we have $\tilde{\gamma}(\tilde{t}) = \gamma(t)$ if and only if $(\tilde{t}, \tilde{t}^2, \tilde{t}^3) = (t^3, t^6, t^9)$, that is, if and only if, $\tilde{t} = t^3$, $\tilde{t}^2 = t^6$ and $\tilde{t}^3 = t^9$. Since the map $\mathbb{R} \rightarrow \mathbb{R}$, $t \mapsto t^3$ is bijective, it follows that $\tilde{\gamma}$ has the same image as γ [4 marks].

- (ii) We have $\gamma(t) = (4t, 4 \cosh(t), 3 \sinh(t))$ and $\gamma'(t) = (4, 4 \sinh(t), 3 \cosh(t))$. Thus

$$\|\gamma'(t)\|^2 = 16 + 16 \sinh(t)^2 + 9 \cosh(t)^2 = 25 \cosh(t)^2,$$

where we used $\cosh(t)^2 - \sinh(t)^2 = 1$. The arc length function is

$$s(t) = \int_0^t \|\gamma'(u)\| du = 5 \int_0^t \cosh(u) du = 5(\sinh(t) - \sinh(0)) = 5 \sinh(t)$$

[4 marks]. This implies $t = \sinh^{-1}(\frac{1}{5}s)$ and hence

$$\begin{aligned} \mathbb{R} \rightarrow \mathbb{R}^3, \quad s \mapsto & \left(4 \sinh^{-1}\left(\frac{1}{5}s\right), 4 \cosh\left(\sinh^{-1}\left(\frac{1}{5}s\right)\right), 3 \sinh\left(\sinh^{-1}\left(\frac{1}{5}s\right)\right)\right) \\ & = \left(4 \sinh^{-1}\left(\frac{1}{5}s\right), 4\sqrt{1 + \frac{1}{25}s^2}, \frac{3}{5}s\right) \end{aligned}$$

is a unit speed parametrization of γ [4 marks].

- (iii) Assume that γ lies in the plane $\Pi = \{(x, y, z) \in \mathbb{R}^3 : (x, y, z) \cdot a = d\}$ with $a \in \mathbb{R}^3$, $\|a\| = 1$, $d \in \mathbb{R}$. Consider the curve $\tilde{\gamma} = \gamma + \frac{1}{\kappa}\mathbf{n}$. Using the Frenet-Serret equation $\dot{\mathbf{n}} = -\kappa\mathbf{t} + \tau\mathbf{b}$, we get

$$\dot{\tilde{\gamma}} = \dot{\gamma} + \frac{1}{\kappa}\dot{\mathbf{n}} = \mathbf{t} + \frac{1}{\kappa}(-\kappa\mathbf{t} + \tau\mathbf{b}) = \frac{\tau}{\kappa}\mathbf{b} = 0$$

since γ is a planar curve and the torsion of planar curves vanishes. It follows that there exists $x_0 \in \mathbb{R}^3$ so that $\tilde{\gamma} = x_0$ and thus $\gamma + \frac{1}{\kappa}\mathbf{n} = x_0$. This tells us that γ lies on the sphere in \mathbb{R}^3 with center x_0 and radius $\frac{1}{\kappa}$ [6 marks]. Since γ is contained in Π , γ lies on the (non-empty) intersection of the sphere and the plane Π , which is a circle, or a point (which cannot happen since γ is unit speed and hence non-constant) [3 marks].

- B 12.** (i) Clearly, σ is smooth [1 mark]. Assume that $\sigma(u, v) = \sigma(u', v')$. Then $u + \cosh(v) = u' + \cosh(v')$ and $u + \sinh(v) = u' + \sinh(v')$. Subtracting these two equations gives $e^{-v} = \cosh(v) - \sinh(v) = \cosh(v') - \sinh(v') = e^{-v'}$ and thus $v = v'$. Inserting this into $u + \cosh(v) = u' + \cosh(v')$ implies $u = u'$. Thus σ is injective [2 marks]. We have

$$\begin{aligned}\sigma_u(u, v) &= (1, 1, 2e^{-v}), \\ \sigma_v(u, v) &= (\sinh(v), \cosh(v), -2ue^{-v}).\end{aligned}$$

Since $\sinh(v) \neq \cosh(v)$ for all $v \in \mathbb{R}$, $\sigma_u(u, v)$ and $\sigma_v(u, v)$ are linearly independent for all $(u, v) \in \mathbb{R}^2$ [2 marks]. It follows that σ is a regular surface patch. Putting $x = u + \cosh(v)$, $y = u + \sinh(v)$ and $z = 1 + 2ue^{-v}$, we see that $z = x^2 - y^2$. Thus σ parametrizes part of the hyperbolic paraboloid $z = x^2 - y^2$ [2 marks]. We have $x - y = e^{-v}$ and $x + y = 2u + e^v$. Thus $x - y > 0$ and $v = -\ln(x - y) = \ln(\frac{1}{x - y})$. Then $u = \frac{1}{2}(x + y - e^v) = \frac{1}{2}(x + y - \frac{1}{x - y})$. Altogether we see that σ parametrizes the part of the hyperbolic paraboloid $z = x^2 - y^2$ given by $x > y$ [3 marks].

- (ii) We have $\gamma'_{v_0}(t) = \sigma_u(t, v_0) = (\cos(v_0), \sin(v_0), 1)$ and hence $\|\gamma'_{v_0}(t)\|^2 = 2$ [2 marks]. The length of γ_{v_0} is $\int_a^b \|\gamma'_{v_0}(t)\| dt = \sqrt{2}(b - a)$ [2 marks] and independent of v_0 [1 mark].
- (iii) The general geodesic equations are

$$\begin{aligned}\frac{d}{dt}(E\dot{u} + F\dot{v}) &= \frac{1}{2}(E_u\dot{u}^2 + 2F_u\dot{u}\dot{v} + G_u\dot{v}^2), \\ \frac{d}{dt}(F\dot{u} + G\dot{v}) &= \frac{1}{2}(E_v\dot{u}^2 + 2F_v\dot{u}\dot{v} + G_v\dot{v}^2)\end{aligned}$$

[2 marks]. By assumption, we have $E = 1$, $F = 0$ and $G = g(u)$. Inserting this into the geodesic equations gives

$$\ddot{u} = \frac{1}{2}g'(u)\dot{v}^2, \quad g'(u)\dot{u}\dot{v} + g(u)\ddot{v} = 0. \quad (*)$$

[2 marks]. We have $\dot{\gamma}(0) = \sigma_u(u_0, v_0)\dot{u}(0) + \sigma_v(u_0, v_0)\dot{v}(0)$. From the initial condition $\dot{\gamma}(0) = \frac{\sigma_u(u_0, v_0)}{\|\sigma_u(u_0, v_0)\|}$ we therefore get $\dot{u}(0) = \frac{1}{\|\sigma_u(u_0, v_0)\|} = a$ and $\dot{v}(0) = 0$ [2 marks]. We thus have to solve (*) subject to the initial conditions $u(0) = u_0$, $v(0) = v_0$, $\dot{u}(0) = a$, $\dot{v}(0) = 0$ [1 mark]. The obvious solution is $u(t) = at + u_0$ and $v(t) = v_0$ [2 marks]. The geodesic $\gamma(t) = \sigma(u(t), v(t))$ with $\gamma(0) = \sigma(u_0, v_0)$ and $\dot{\gamma}(0) = a\sigma_u(u_0, v_0)$ is therefore $\gamma(t) = \sigma(at + u_0, v_0)$ [1 mark].

- B 13.** (i) We have $\sigma(u, v) = \gamma(u) + r(\cos(v)\mathbf{n}(u) + \sin(v)\mathbf{b}(u))$. Using the Frenet-Serret equations we get

$$\begin{aligned}\sigma_u(u, v) &= \mathbf{t}(u) + r(\cos(v)\dot{\mathbf{n}}(u) + \sin(v)\dot{\mathbf{b}}(u)) \\ &= \mathbf{t}(u) + r(\cos(v)(-\kappa(u)\mathbf{t}(u) + \tau(u)\mathbf{b}(u)) + \sin(v)(-\tau(u)\mathbf{n}(u))) \\ &= (1 - r\kappa(u)\cos(v))\mathbf{t}(u) - r\tau(u)\sin(v)\mathbf{n}(u) + r\tau(u)\cos(v)\mathbf{b}(u), \\ \sigma_v(u, v) &= r(-\sin(v)\mathbf{n}(u) + \cos(v)\mathbf{b}(u)) = -r\sin(v)\mathbf{n}(u) + r\cos(v)\mathbf{b}(u)\end{aligned}$$

[2 marks]. Using the fact that $\mathbf{t} \times \mathbf{n} = \mathbf{b}$, $\mathbf{n} \times \mathbf{b} = \mathbf{t}$ and $\mathbf{b} \times \mathbf{t} = \mathbf{n}$, we get

$$(\sigma_u \times \sigma_v)(u, v) = r(r\kappa(u)\cos(v) - 1)(\cos(v)\mathbf{n}(u) + \sin(v)\mathbf{b}(u)).$$

Since $0 < \kappa(u) < \frac{1}{r}$, we have $r\kappa(u)\cos(v) - 1 < 0$. Hence the unit normal \mathbf{N} of σ is

$$\mathbf{N}(u, v) = \frac{(\sigma_u \times \sigma_v)(u, v)}{\|(\sigma_u \times \sigma_v)(u, v)\|} = -\cos(v)\mathbf{n}(u) - \sin(v)\mathbf{b}(u)$$

[3 marks]. The Gauss map is the map

$$\sigma(U) \rightarrow S^2, \sigma(u, v) \mapsto \mathbf{N}(u, v) = -\cos(v)\mathbf{n}(u) - \sin(v)\mathbf{b}(u)$$

[1 mark]. For $\alpha(t) = \sigma(u_0, \frac{t}{r}) = \gamma(u_0) + r(\cos(\frac{t}{r})\mathbf{n}(u_0) + \sin(\frac{t}{r})\mathbf{b}(u_0))$ we get

$$\begin{aligned}\dot{\alpha}(t) &= -\sin(\frac{t}{r})\mathbf{n}(u_0) + \cos(\frac{t}{r})\mathbf{b}(u_0), \\ \ddot{\alpha}(t) &= \frac{1}{r}(-\cos(\frac{t}{r})\mathbf{n}(u_0) - \sin(\frac{t}{r})\mathbf{b}(u_0)) = \frac{1}{r}\mathbf{N}(u_0, \frac{t}{r}).\end{aligned}$$

Thus α is unit speed and $\ddot{\alpha}(t)$ is parallel to $\mathbf{N}(u_0, \frac{t}{r})$ for all t , which shows that α is a geodesic. [4 marks]

- (ii) (a) True [1 mark]; follows from Theorema Egregium [4 marks].
 (b) False [1 mark]; consider for example corresponding parts of a plane and a round cylinder which are isometric to each other. The plane has $H_1 = 0$ and the round cylinder has $H_2 \neq 0$ [4 marks].
- (iii) The Euler number χ of S^2 is equal to 2 [1 mark] and satisfies $\chi = V - E + F = 12 - E + 20 = 32 - E$ [2 marks]. It follows that there are 30 edges in the triangulation [2 marks].