VIRTUAL RESOLUTIONS FOR A PRODUCT OF PROJECTIVE SPACES

CHRISTINE BERKESCH ZAMAERE, DANIEL ERMAN, AND GREGORY G. SMITH

ABSTRACT. Syzygies capture intricate geometric properties of a subvariety in projective space. However, when the ambient space is a product of projective spaces or a more general smooth projective toric variety, minimal free resolutions over the Cox ring are too long and contain many geometrically superfluous summands. In this paper, we construct some much shorter free complexes that better encode the geometry.

The geometric and algebraic sources of locally-free resolutions have complementary advantages. To see the differences, consider a smooth projective toric variety X together with its $\operatorname{Pic}(X)$ -graded Cox ring S. The local version of the Hilbert Syzygy Theorem implies that any coherent \mathcal{O}_X -module admits a locally-free resolution of length at most $\dim X$; see [Har77, Exercise III.6.9]. The global version of the Hilbert Syzygy Theorem implies that every saturated module over the polynomial ring S has a minimal free resolution of length at most $\dim S - 1$, so any coherent \mathcal{O}_X -module has a locally-free resolution of the same length; see [Cox95, Proposition 3.1]. Unlike the geometric approach, this algebraic method only involves vector bundles that are a direct sum of line bundles. When X is projective space, these geometric and algebraic constructions usually coincide. However, when the Picard number of X is greater than 1, the locally-free resolutions arising from the minimal free resolution of an S-module are longer, and typically much longer, than their geometric counterparts.

To enjoy the best of both worlds, we focus on a more flexible algebraic source for locally-free resolutions. The following definition, beyond providing concise terminology, highlights this source.

Definition 1.1. A free complex $F := [F_0 \longleftarrow F_1 \longleftarrow F_2 \longleftarrow \cdots]$ of Pic(X)-graded S-modules is called a *virtual resolution* of a Pic(X)-graded S-module M if the corresponding complex \widetilde{F} of vector bundles on X is a locally-free resolution of the sheaf \widetilde{M} .

In other words, a virtual resolution is a free complex of S-modules whose higher homology groups are supported on the irrelevant ideal of X. The benefits of allowing a limited amount of homology are already present in other parts of commutative algebra including almost ring theory [GR03], where one accepts homology annihilated by a given idempotent ideal, and phantom homology [HH93], where one admits cycles that are in the tight closure of the boundaries.

For projective space, minimal free resolutions are important in the study of points [GGP95, EP99], curves [Voi02, EL15], surfaces [GP99, DS00], and moduli spaces [Far09, DFS16]. Our overarching goal is to demonstrate that the right analogues for subschemes in a smooth complete toric variety use virtual resolutions rather than minimal free resolutions. This distinction is not apparent on

²⁰¹⁰ Mathematics Subject Classification. 13D02; 14M25, 14F05.

CBZ was partially supported by the NSF Grant DMS-1440537, DE was partially supported by the NSF Grants DMS-1302057 and DMS-1601619, and GGS was partially supported by the NSERC.

projective space because the New Intersection Theorem [Rob87] establishes that a free complex with finite-length higher homology groups has to be at least as long as the minimal free resolution. In contrast, for other toric varieties such as products of projective spaces, allowing irrelevant homology may yield simpler complexes; see Example 1.4.

Throughout this paper, we write $\mathbb{P}^n := \mathbb{P}^{n_1} \times \mathbb{P}^{n_2} \times \cdots \times \mathbb{P}^{n_r}$ for the product of projective spaces with dimension vector $\mathbf{n} := (n_1, n_2, \dots, n_r) \in \mathbb{N}^r$ over a field \mathbb{k} . Let $S := \mathbb{k}[x_{i,j} : 1 \le i \le r, 0 \le j \le n_i]$ be the Cox ring of \mathbb{P}^n and let $B := \bigcap_{i=1}^r \langle x_{i,0}, x_{i,1}, \dots, x_{i,n_i} \rangle$ be its irrelevant ideal. We identify the Picard group of \mathbb{P}^n with \mathbb{Z}^r and partially order the elements via their components. If e_1, e_2, \dots, e_r is the standard basis of \mathbb{Z}^r , then the polynomial ring S has the \mathbb{Z}^r -grading induced by $\deg(x_{i,j}) := e_i$. We first reprove the existence of short virtual resolutions; compare with Corollary 1.14 in [EES15].

Proposition 1.2. Every finitely-generated \mathbb{Z}^r -graded B-saturated S-module has a virtual resolution of length at most $|\mathbf{n}| := n_1 + n_2 + \cdots + n_r = \dim \mathbb{P}^n$.

Since $\dim S - \dim \mathbb{P}^n = r$, we see that a minimal free resolution can be arbitrarily long when compared with a virtual resolution. A proof of Proposition 1.2 appears in Section 2.

Besides having shorter representatives, virtual resolutions also exhibit a closer relationship with Castelnuovo–Mumford regularity than minimal free resolutions. On projective space, Castelnuovo–Mumford regularity has two equivalent descriptions: one arising from the vanishing of sheaf cohomology and another arising from the Betti numbers in a minimal free resolutions. However, on more general toric varieties, the multigraded Castelnuovo–Mumford regularity is not determined by a minimal free resolution; see Theorem 1.5 in [MS04] or Theorem 4.7 in [BC17]. From this perspective, we demonstrate that virtual resolutions improve on minimal free resolutions in two ways. First, Theorem 3.1 proves that the set of virtual resolutions of a module determines its multigraded Castelnuovo–Mumford regularity. Second, the next theorem, from Section 4, demonstrates how to use regularity to extract a virtual resolution from a minimal free resolution.

Theorem 1.3. Let M be a finitely-generated \mathbb{Z}^r -graded B-saturated S-module that is d-regular. If G is the free subcomplex of a minimal free resolution of M consisting of all summands generated in degree at most d + n, then G is a virtual resolution of M.

For convenience, we refer to the free complex G as the *virtual resolution of the pair* (M, d). Algorithm 4.4 shows that it can be constructed without computing the entire minimal free resolution.

The following example illustrates that a virtual resolution can be much shorter and much thinner than the minimal free resolution. Surprisingly, it follows that a majority of the summands in the minimal free resolution are often irrelevant for geometric purposes.

Example 1.4. A hyperelliptic curve C of genus 4 can be embedded as a curve of bidegree (2,8) in $\mathbb{P}^1 \times \mathbb{P}^2$; see [Har77, Theorem IV.5.4]. For instance, the B-saturated S-ideal

$$I := \left\langle \begin{array}{c} x_{1,1}^3 x_{2,0} - x_{1,1}^3 x_{2,1} + x_{1,0}^3 x_{2,2}, x_{1,0}^2 x_{2,0}^2 + x_{1,1}^2 x_{2,1}^2 + x_{1,0} x_{1,1} x_{2,2}^2, x_{1,1}^2 x_{2,0}^3 - x_{1,1}^2 x_{2,0}^2 x_{2,1} - x_{1,0} x_{1,1} x_{2,1}^2 x_{2,2} - x_{1,0}^2 x_{2,2}^2, x_{1,0} x_{1,1} x_{2,0}^3 - x_{1,1}^2 x_{2,0}^2 x_{2,1} - x_{1,0} x_{1,1} x_{2,0}^2 x_{2,1} - x_{1,0} x_{2,1}^2 x_{2,2} - x_{1,0}^2 x_{2,1}^2 x_{2,1} x_{2,2}^2 - x_{1,1}^2 x_{2,0}^2 x_{2,1}^2 - x_{1,1} x_{2,0}^2 x_{2,1}^2 - x_{1,1} x_{2,0}^2 x_{2,1}^2 - x_{1,0} x_{2,0}^2 x_{2,1}^2 - x_{1,0} x_{2,0}^4 x_{2,1}^2 - x_{1,0} x_{2,0}^4 x_{2,1}^2 + x_{1,0} x_{2,0}^2 x_{2,1}^2 x_{2,2}^2 + x_{1,0} x_{2,0}^2 x_{2,1}^2 x_{2,1}^2 x_{2,2}^2 - x_{1,0} x_{2,0}^2 x_{2,1}^2 x_{2,1}^2 x_{2,1}^2 x_{2,2}^2 + x_{1,0}^2 x_{2,0}^2 x_{2,1}^2 x_{2,1}^2 x_{2,2}^2 + x_{1,0}^2 x_{2,0}^2 x_{2,1}^2 x_{2,1$$

defines such a curve. Macaulay [M2] shows that the minimal free resolution of S/I has the form

$$(1.4.1) \begin{array}{c} S(-3,-1)^1 \\ \oplus \\ S(-2,-2)^1 \\ \oplus \\ S(-2,-3)^2 \\ \oplus \\ S(-2,-5)^6 \\ \oplus \\ S(-2,-5)^6 \\ \oplus \\ S(-2,-5)^6 \\ \oplus \\ S(-2,-5)^6 \\ \oplus \\ S(-2,-7)^2 \\ \longleftarrow \\ S(-2,-7)^2 \\ \oplus \\ S(-3,-7)^1 \\ \oplus \\ S(-3,-7)$$

Using the Riemann–Roch Theorem [Har77, Theorem IV.1.3], one verifies that the module S/I is (2,1)-regular, so the virtual resolution of the pair (S/I,(2,1)) has the much simpler form

(1.4.2)
$$S^{1} \leftarrow S(-2,-2)^{1} \leftarrow S(-3,-3)^{3} \leftarrow 0.$$

$$S(-2,-3)^{2}$$

If the ideal $J \subset S$ is the image of the first map in (1.4.2), then we have $J = I \cap Q$ for some ideal Q whose radical contains the irrelevant ideal. Using Proposition 2.4, we can even conclude that S/J is Cohen–Macaulay and J is the ideal of maximal minors of the 4×3 matrix

(1.4.3)
$$\varphi := \begin{bmatrix} x_{2,1}^2 & x_{2,2}^2 & -x_{2,0}^2 \\ -x_{1,1}(x_{2,0} - x_{2,1}) & 0 & x_{1,0}x_{2,2} \\ x_{1,0} & -x_{1,1} & 0 \\ 0 & x_{1,0} & x_{1,1} \end{bmatrix} .$$

As an initial step towards our larger goal, we formulate a novel analogue for properties of points in projective space. Although any punctual subscheme of projective space is arithmetically Cohen–Macaulay, this almost always fails for a zero-dimensional subscheme of \mathbb{P}^n ; see [GVT15]. However, by considering virtual resolutions, we obtain the following variant.

Theorem 1.5. Let $Z \subset \mathbb{P}^n$ be a zero-dimensional scheme and let I be its corresponding B-saturated S-ideal. There exists an S-ideal Q, whose radical contains B, such that the minimal free resolution of $S/(I \cap Q)$ has length |n|. In particular, the minimal free resolution of $S/(I \cap Q)$ is a virtual resolution of S/I.

This theorem, proven in Section 5, does not imply that $S/(I \cap Q)$ is itself Cohen–Macaulay, as the components of Q will often have codimension less than |n|. However, when the ambient variety is $\mathbb{P}^1 \times \mathbb{P}^1$, the ring $S/(I \cap Q)$ will be Cohen–Macaulay of codimension 2. In this case, Corollary 5.2 shows that there is a matrix whose maximal minors cut out Z scheme-theoretically. Proposition 5.8 extends this to general points on any smooth toric surface.

As a second and perhaps more substantial step, we apply virtual resolutions to deformation theory. On projective space, there are three classic situations in which the particular structure of the minimal free resolution allows one to show that all deformations have the same structure: arithmetically

Cohen–Macaulay subschemes of codimension 2, arithmetically Gorenstein subschemes in codimension 3, and complete intersections; see [Har10, Sections 2.8–2.9]. We generalize these results about unobstructed deformations in projective space as follows.

Theorem 1.6. Consider $Y \subset \mathbb{P}^n$ and let I be the corresponding B-saturated S-ideal. Assume that the generators of I have degrees d_1, d_2, \ldots, d_s and that the natural map $(S/I)_{d_i} \to H^0(Y, \mathcal{O}_Y(d_i))$ is an isomorphism for all $1 \leq i \leq s$. If any one of the following conditions hold

- (i) the subscheme Y has codimension 2 and there is $d \in reg(S/I)$ such that the virtual resolution of the pair (S/I, d) has length 2;
- (ii) each factor in \mathbb{P}^n has dimension at least 2, the subscheme Y has codimension 3, and there is $d \in \operatorname{reg}(S/I)$ such that the virtual resolution of the pair (S/I, d) is a self-dual complex (up to a twist) of length 3; or
- (iii) there is $d \in reg(S/I)$ such that virtual resolution of the pair (S/I, d) is a Koszul complex of length codim Y;

then the embedded deformations of Y in \mathbb{P}^n are unobstructed and the component of the multigraded Hilbert scheme of \mathbb{P}^n containing the point corresponding to Y is unirational.

To illustrate this theorem, we can reuse the hyperelliptic curve in Example 1.4.

Example 1.7. By reinterpreting Example 1.4, we see that the hyperelliptic curve $C \subset \mathbb{P}^1 \times \mathbb{P}^2$ satisfies condition (i) in Theorem 1.6. It follows that the embedded deformations of C are unobstructed and the corresponding component of the multigraded Hilbert scheme of $\mathbb{P}^1 \times \mathbb{P}^2$ can be given an explicit unirational parametrization by varying the entries in the 4×3 matrix φ from (1.4.3). \diamond

Three other geometric applications for virtual resolutions are collected in Section 6. The first, Proposition 6.1, provides an unmixedness result for subschemes of \mathbb{P}^n that have a virtual resolution whose length equals its codimension. The second, Proposition 6.5, gives sharp bounds on the Castelnuovo–Mumford regularity of a tensor products of coherent $\mathcal{O}_{\mathbb{P}^n}$ -modules; compare with [Laz04, Prop. 1.8.8]. Lastly, Proposition 6.8 describes new vanishing results for the higher-direct images of sheaves, which are optimal in many cases.

The final section presents some promising directions for future research.

Conventions. In this article, we work in the product $\mathbb{P}^n := \mathbb{P}^{n_1} \times \mathbb{P}^{n_2} \times \cdots \times \mathbb{P}^{n_r}$ of projective spaces with dimension vector $\mathbf{n} := (n_1, n_2, \dots, n_r) \in \mathbb{N}^r$ over a field \mathbb{k} . Its Cox ring is the polynomial ring $S := \mathbb{k}[x_{i,j} : 1 \le i \le r, 0 \le j \le n_i]$ and its irrelevant ideal is $B := \bigcap_{i=1}^r \langle x_{i,0}, x_{i,1}, \dots, x_{i,n_i} \rangle$. The Picard group of \mathbb{P}^n is identified with \mathbb{Z}^r and the elements are partially ordered componentwise. If e_1, e_2, \dots, e_r is the standard basis of \mathbb{Z}^r , then S has the \mathbb{Z}^r -grading induced by $\deg(x_{i,j}) := e_i$. We assume that all S-modules are finitely generated and \mathbb{Z}^r -graded.

Acknowledgements. Some of this research was completed during visits to the Banff International Research Station (BIRS) and the Mathematical Sciences Research Institute (MSRI), and we are very grateful for their hospitality. We also thank Lawrence Ein, David Eisenbud, Craig Huneke, Nathan Ilten, Rob Lazarsfeld, Diane Maclagan, Frank-Olaf Schreyer, and Ian Shipman for helpful conversations.

2. Existence of Short Virtual Resolutions

This section, by proving Proposition 1.2, establishes the existence of virtual resolutions whose length is bounded above by the dimension of \mathbb{P}^n . In particular, these virtual resolutions are typically shorter than a minimal free resolution. Moreover, Proposition 2.4 shows that Proposition 1.2 provides the best possible uniform bound.

Our proof of Proposition 1.2 is based on a minor variation of Beilinson's resolution of the diagonal; compare with Proposition 3.2 in [Căl05] or Lemma 8.27 in [Huy06]. Given an \mathcal{O}_{X_j} -module \mathcal{F}_j for all $1 \leq j \leq n$, their external tensor product is

$$\mathscr{F}_1 \boxtimes \mathscr{F}_2 \boxtimes \cdots \boxtimes \mathscr{F}_m := p_1^* \mathscr{F}_1 \otimes_{\mathscr{O}_{X_1 \times X_2}} p_2^* \mathscr{F}_2 \otimes_{\mathscr{O}_{X_1 \times X_2}} \cdots \otimes_{\mathscr{O}_{X_1 \times X_2}} p_m^* \mathscr{F}_m$$

where p_j denotes the projection map from the Cartesian product $X_1 \times X_2 \times \cdots \times X_m$ to X_j . In particular, for all $u \in \mathbb{Z}^r$, we have $\mathscr{O}_{\mathbb{P}^n}(u) = \mathscr{O}_{\mathbb{P}^{n_1}}(u_1) \boxtimes \mathscr{O}_{\mathbb{P}^{n_2}}(u_2) \boxtimes \cdots \boxtimes \mathscr{O}_{\mathbb{P}^{n_r}}(u_r)$. With this notation, we can describe the resolution of the diagonal $\mathbb{P}^n \hookrightarrow \mathbb{P}^n \times \mathbb{P}^n$.

Lemma 2.1. If $\mathscr{T}^{e_i}_{\mathbb{P}^n} := \mathscr{O}_{\mathbb{P}^{n_1}} \boxtimes \mathscr{O}_{\mathbb{P}^{n_2}} \boxtimes \cdots \boxtimes \mathscr{O}_{\mathbb{P}^{n_{i-1}}} \boxtimes \mathscr{T}_{\mathbb{P}^{n_i}} \boxtimes \mathscr{O}_{\mathbb{P}^{n_{i+1}}} \boxtimes \cdots \boxtimes \mathscr{O}_{\mathbb{P}^{n_r}}$ for $1 \leqslant i \leqslant r$, then the diagonal $\mathbb{P}^n \hookrightarrow \mathbb{P}^n \times \mathbb{P}^n$ is the zero scheme of a global section of $\bigoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^n}(e_i) \boxtimes \mathscr{T}^{e_i}_{\mathbb{P}^n}(-e_i)$. Hence, the diagonal has a locally-free resolution of the form

$$\mathscr{O}_{\mathbb{P}^{m{n}} imes\mathbb{P}^{m{n}}} \longleftarrow igoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^{m{n}}}(-m{e}_i)oxtimes \Omega^{m{e}_i}_{\mathbb{P}^{m{n}}}(m{e}_i) \longleftarrow igoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^{m{n}}}(-m{u})oxtimes \Omega^{m{u}}_{\mathbb{P}^{m{n}}}(m{u}) \longleftarrow \longleftarrow \mathscr{O}_{\mathbb{P}^{m{n}}}(-m{n})oxtimes \Omega^{m{n}}_{\mathbb{P}^{m{n}}}(m{n}), \ 0 \leqslant u \leqslant n \ |u|=2$$

where $\Omega^{a}_{\mathbb{P}^{n}} := \Omega^{a_{1}}_{\mathbb{P}^{n_{1}}} \boxtimes \Omega^{a_{2}}_{\mathbb{P}^{n_{2}}} \boxtimes \cdots \boxtimes \Omega^{a_{r}}_{\mathbb{P}^{n_{r}}}$ is the external tensor product of the exterior powers of the cotangent bundles on the factors of \mathbb{P}^{n} .

Proof. For each $1 \le i \le r$, fix a basis $x_{i,0}, x_{i,1}, \dots, x_{i,n_i}$ for $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(e_i))$. The Euler sequence on \mathbb{P}^{n_i} yields

$$0 \longleftarrow \mathscr{T}^{oldsymbol{e}_i}_{\mathbb{P}^{oldsymbol{n}}} \longleftarrow igoplus_{i=0}^{n_i} \mathscr{O}_{\mathbb{P}^{oldsymbol{n}}}(oldsymbol{e}_i) \stackrel{[x_{i,0} \ x_{i,1} \ \cdots \ x_{i,n_i}]}{-} \mathscr{O}_{\mathbb{P}^{oldsymbol{n}}} \longleftarrow 0;$$

see [CLS11, Theorem 8.1.6]. Knowing the cohomology of line bundles on \mathbb{P}^n , the associated long exact sequence gives $H^0(\mathbb{P}^n, \mathscr{T}^{e_i}_{\mathbb{P}^n}(-e_i)) \cong \bigoplus_{j=0}^{n_i} H^0(\mathbb{P}^n, \mathscr{O}_{\mathbb{P}^n})$. A basis for $\bigoplus_{j=0}^{n_i} H^0(\mathbb{P}^n, \mathscr{O}_{\mathbb{P}^n})$ is given by the dual basis $x_{i,0}^*, x_{i,1}^*, \dots, x_{i,n_i}^*$. Let $\partial/\partial x_{i,j}$ denote the image of $x_{i,j}^*$ in $H^0(\mathbb{P}^n, \mathscr{T}_{\mathbb{P}^n}(-e_i))$. Consider $s \in H^0(\mathbb{P}^n \times \mathbb{P}^n, \bigoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^n}(e_i) \boxtimes \mathscr{T}^{e_i}_{\mathbb{P}^n}(-e_i))$ given by

$$s := \left(\sum_{j=0}^{n_1} x_{1,j} \frac{\partial}{\partial y_{1,j}}, \sum_{j=0}^{n_2} x_{2,j} \frac{\partial}{\partial y_{2,j}}, \dots, \sum_{j=0}^{n_r} x_{r,j} \frac{\partial}{\partial y_{r,j}}\right)$$

where $x_{i,j}$ and $y_{i,j}$ are the coordinates on the first and second factor of $\mathbb{P}^n \times \mathbb{P}^n$ respectively. We claim that the zero scheme of s equals the diagonal in $\mathbb{P}^n \times \mathbb{P}^n$. By symmetry, it suffices to check this on a single affine open neighbourhood. If $x_{1,0}x_{2,0}\cdots x_{r,0} \neq 0$ and $y_{1,0}y_{2,0}\cdots y_{r,0} \neq 0$, then the Euler relations yield

$$\sum_{j=0}^{n_i} x_{i,j} \frac{\partial}{\partial y_{i,j}} = x_{i,0} \left(-\frac{1}{y_{i,0}} \sum_{j=1}^{n_i} y_{i,j} \frac{\partial}{\partial y_{i,j}} \right) + \sum_{j=1}^{n_i} x_{i,j} \frac{\partial}{\partial y_{i,j}} = \frac{1}{y_{i,0}} \sum_{j=1}^{n_i} (x_{i,j} y_{i,0} - x_{i,0} y_{i,j}) \frac{\partial}{\partial y_{i,j}},$$

for each $1 \leqslant i \leqslant r$. It follows that s = 0 if and only if $\frac{x_{i,j}}{x_{i,0}} = \frac{y_{i,j}}{y_{i,0}}$ for all $1 \leqslant i \leqslant r$ and $1 \leqslant j \leqslant n_i$. Hence, the global section s vanishes precisely on the diagonal $\mathbb{P}^n \hookrightarrow \mathbb{P}^n \times \mathbb{P}^n$.

The Koszul complex associated to s is the required locally-free resolution of the diagonal, because \mathbb{P}^n is smooth and the codimension of the diagonal equals the rank of the vector bundle $\bigoplus_{i=1}^r \mathscr{O}_{\mathbb{P}^n}(e_i) \boxtimes \mathscr{T}^{e_i}_{\mathbb{P}^n}(-e_i)$; see [Laz04, Section B.2]. Since $\Omega^{e_i}_{\mathbb{P}^n} = \mathscr{H}om_{\mathscr{O}_{\mathbb{P}^n}}(\mathscr{T}^{e_i}_{\mathbb{P}^n}, \mathscr{O}_{\mathbb{P}^n})$, we have

$$igwedge^k ig(\mathscr{O}_{\mathbb{P}^{m{n}}}(-e_i)oxtimes\Omega^{m{e}_i}_{\mathbb{P}^{m{n}}}(e_i)ig) = igoplus_{m{0}\leqslantm{u}\leqslantm{n}}m{\mathscr{O}}_{\mathbb{P}^{m{n}}}(-m{u})oxtimes\Omega^{m{u}}_{\mathbb{P}^{m{n}}}(m{u})\,,$$

for
$$0 \leqslant k \leqslant |\mathbf{n}|$$
.

Proof of Proposition 1.2. Let p_1 and p_2 be the projections of $\mathbb{P}^n \times \mathbb{P}^n$ onto the first and second factors respectively. For any $u \in \mathbb{Z}^r$, the Fujita Vanishing Theorem [Fuj83, Theorem 1] implies that $\Omega^u_{\mathbb{P}^n}(u+d) \otimes \widetilde{M}$ has no higher cohomology for any sufficiently positive $d \in \mathbb{Z}^r$. Let \mathscr{K} be the locally-free resolution of the diagonal $\mathbb{P}^n \hookrightarrow \mathbb{P}^n \times \mathbb{P}^n$ described in Lemma 2.1. Both hypercohomology spectral sequences, namely

$${}'\mathrm{E}_{2}^{p,q} := \mathrm{H}^{p} \left(\mathbf{R}^{q} \mathrm{p}_{1*} \left(\mathrm{p}_{2}^{*} \widetilde{M}(\boldsymbol{d}) \otimes_{\mathscr{O}_{\mathbb{P}\boldsymbol{n}_{\vee}\mathbb{P}\boldsymbol{n}}} \mathscr{K} \right) \right) \ \ \text{and} \ \ {}''\mathrm{E}_{2}^{p,q} := \mathbf{R}^{p} \mathrm{p}_{1*} \, \mathrm{H}^{q} \left(\mathrm{p}_{2}^{*} \widetilde{M}(\boldsymbol{d}) \otimes_{\mathscr{O}_{\mathbb{P}\boldsymbol{n}_{\vee}\mathbb{P}\boldsymbol{n}}} \mathscr{K} \right),$$

converge to $\mathbf{R}^{p+q}\mathbf{p}_{1*}(\mathbf{p}_2^*\widetilde{M}(\mathbf{d})\otimes_{\mathscr{O}_{\mathbb{P}^n\times\mathbb{P}^n}}\mathscr{K})$; see [EGA3I, Section 12.4]. Since \mathscr{K} is a locally-free resolution of the diagonal, it follows that $''\mathbf{E}_2^{0,0}\cong\widetilde{M}(\mathbf{d})$ and $''\mathbf{E}_2^{p,q}=0$ when either $p\neq 0$ or $q\neq 0$; compare with [Huy06, Proposition 8.28]. Hence, we conclude that

$$\mathbf{R}^{p+q} \mathbf{p}_{1*} \left(\mathbf{p}_2^* \widetilde{M}(\boldsymbol{d}) \otimes_{\mathscr{O}_{\mathbb{P}^n \times \mathbb{P}^n}} \mathscr{K} \right) \cong \begin{cases} \widetilde{M}(\boldsymbol{d}) & \text{if } p = 0 = q, \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand, the first page of the other hypercohomology spectral sequence is

$$\begin{split} {}'\mathbf{E}_{1}^{p,q} &= \mathbf{R}^{q} \mathbf{p_{1}}_{*} \Big(\mathbf{p_{2}^{*}} \widetilde{M}(\boldsymbol{d}) \otimes_{\mathscr{O}_{\mathbb{P}^{n} \times \mathbb{P}^{n}}} \mathscr{K}_{-p} \Big) = \mathbf{R}^{q} \mathbf{p_{1}}_{*} \Bigg(\bigoplus_{\substack{0 \leqslant \boldsymbol{u} \leqslant \boldsymbol{n} \\ |\boldsymbol{u}| = -p}} \mathscr{O}_{\mathbb{P}^{n}} (-\boldsymbol{u}) \boxtimes \Big(\Omega_{\mathbb{P}^{n}}^{\boldsymbol{u}} \otimes \widetilde{M}(\boldsymbol{u} + \boldsymbol{d}) \Big) \Bigg) \\ &= \bigoplus_{\substack{0 \leqslant \boldsymbol{u} \leqslant \boldsymbol{n} \\ |\boldsymbol{u}| = -p}} \mathscr{O}_{\mathbb{P}^{n}} (-\boldsymbol{u}) \otimes_{\mathbb{K}} H^{q} \Big(\mathbb{P}^{n}, \Omega_{\mathbb{P}^{n}}^{\boldsymbol{u}} \otimes \widetilde{M}(\boldsymbol{u} + \boldsymbol{d}) \Big) \,. \end{split}$$

Our positivity assumption on \boldsymbol{d} implies that $H^q\left(\mathbb{P}^{\boldsymbol{n}},\Omega^{\boldsymbol{u}}_{\mathbb{P}^{\boldsymbol{n}}}\otimes\widetilde{\boldsymbol{M}}(\boldsymbol{u}+\boldsymbol{d})\right)=0$ for all q>0, so $'\mathrm{E}_1^{p,q}$ is concentrated in a single row. Applying the functor $\mathscr{F}\mapsto\bigoplus_{\boldsymbol{v}\in\mathbb{N}^r}H^0\left(\mathbb{P}^{\boldsymbol{n}},\mathscr{F}(\boldsymbol{v})\right)$, we obtain a virtual resolution of M in which the i-th module is

$$\bigoplus_{\substack{\mathbf{0}\leqslant \boldsymbol{u}\leqslant \boldsymbol{n}\\|\boldsymbol{u}|=i}} S(-\boldsymbol{u})\otimes_{\Bbbk} H^q\left(\mathbb{P}^{\boldsymbol{n}},\Omega^{\boldsymbol{u}}_{\mathbb{P}^{\boldsymbol{n}}}\otimes\widetilde{M}(\boldsymbol{u}+\boldsymbol{d})\right).$$

The next examples demonstrate why we want more than just these short virtual resolutions arising from the proof of Proposition 1.2.

Example 2.2. Consider the hyperelliptic curve $C \subset \mathbb{P}^1 \times \mathbb{P}^2$ from Example 1.4. Using d := (2,2) in the construction from the proof of Proposition 1.2 yields a virtual resolution of the form

$$S(-2,-2)^{17} \longleftarrow \frac{S(-2,-3)^{26}}{S(-3,-2)^{15}} \longleftarrow \frac{S(-2,-4)^9}{S(-3,-3)^{22}} \longleftarrow S(-3,-4)^7 \longleftarrow 0.$$

Compared to the virtual resolution in (1.4.2), the length of this complex is longer, the rank of the free modules is higher, and the degrees of the generators are larger. \diamond

Example 2.3. If X is the union of m distinct points on $\mathbb{P}^1 \times \mathbb{P}^1$, then for any sufficiently positive $d = (d_1, d_2)$, the construction in the proof of Proposition 1.2 yields a virtual resolution of the form

$$S(-d_1, -d_2)^m \longleftarrow S(-d_1 - 1, -d_2)^m \bigoplus_{S(-d_1, -d_2 - 1)^m} \longleftarrow S(-d_1 - 1, -d_2 - 1)^m \longleftarrow 0.$$

Unlike the minimal free resolution, this Betti table of this free complex is independent of the geometry of the points, so even short virtual resolutions can obscure the geometric information. \diamond

As a counterpoint to Proposition 1.2, we provide a lower bound on the length of a virtual resolution. Extending the well-known result for projective space, we show that the codimension of any associated prime of M gives a lower bound on the length of any virtual resolution of M.

Proposition 2.4. Let M be a finitely-generated \mathbb{Z}^r -graded S-module. Let Q be an associated prime of M that does not contain the irrelevant ideal B and let $F := [F_0 \longleftarrow F_1 \longleftarrow \cdots \longleftarrow F_p \longleftarrow 0]$ be a virtual resolution of M. These hypotheses yield the following.

- (i) We have codim $Q \leq p$.
- (ii) If Q is the prime ideal for a closed point of \mathbb{P}^n , then we have $p \ge |n|$.
- (iii) If $p \le \min\{n_i + 1 : 1 \le i \le r\}$, then F is a minimal free resolution of $H_0(F)$.

Proof. Localizing at the prime ideal Q, F becomes a free S_Q -resolution of M_Q . Part (i) then follows from the fact that, over the local ring S_Q , the projective dimension of a module is always greater than or equal to the codimension of a module; see [Eis95, Proposition 18.2]. Part (ii) is immediate, as $\operatorname{codim} Q = |n|$ if Q is the prime ideal for a closed point of \mathbb{P}^n .

For part (iii), assume by way of contradiction that F is not a minimal free resolution of $H_0(F)$. It follows that $H_j(F) \neq 0$ for some j > 0; choose the maximal such j. Since F is a virtual resolution of M, the module $H_j(F)$ must be supported on the irrelevant ideal B. Setting $P_i := \langle x_{i,0}, x_{i,1}, \dots, x_{i,n_i} \rangle$ to be the component of the irrelevant ideal B corresponding to the factor \mathbb{P}^{n_i} , there is an index i such that $(H_j(F))_{P_i} \neq 0$. Localizing at P_i yields a complex F_{P_i} of the form

$$\cdots \longleftarrow (F_{j-1})_{P_i} \longleftarrow (F_j)_{P_i} \longleftarrow (F_{j+1})_{P_i} \longleftarrow \cdots \longleftarrow (F_p)_{P_i} \longleftarrow 0$$

where $H_j(F_{P_i})$ is supported on the maximal ideal of S_{P_i} . We deduce that $p > p - j \geqslant \operatorname{codim} P_i = n_i + 1$ from the Peskine–Szpiro Acyclicity Lemma; see [Eis95, Lemma 20.11]. However, this contradicts our assumption that $p \leqslant \min\{n_i + 1 : 1 \leqslant i \leqslant r\}$. Therefore, we conclude that the complex F is a minimal free resolution of $H_0(F)$.

The following simple corollary is useful in applications such as Theorem 1.6.

Corollary 2.5. Let I be a B-saturated S-ideal and let $F = [F_0 \longleftarrow F_1 \longleftarrow \cdots \longleftarrow F_p \longleftarrow 0]$ be a virtual resolution of S/I. If $F_0 = S$ and $p < \min\{n_i + 1 : 1 \le i \le r\}$, then the complex F is a minimal free resolution of S/I.

Proof. By part (iii) of Proposition 2.4, the complex F is a minimal free resolution of $H_0(F)$. The hypothesis that $F_0 = S$ implies that $H_0(F) = S/J$ for some ideal J. Since I is B-saturated and F is a virtual resolution of S/I, we deduce that I equals the B-saturation of J. If we had $I \neq J$, then it would follow that S/J has an associated prime Q that contains the irrelevant ideal B. However, the codimension of Q is at least $\min\{n_i + 1 : 1 \le i \le r\}$. As F is a free resolution of S/J, this would yield the contraction $P \ge \min\{n_i + 1 : 1 \le i \le r\}$; see [Eis95, Proposition 18.2].

Just like in projective space, one can find subvarieties of codimension c which do not admit a virtual resolution of length c.

Example 2.6. Working in $\mathbb{P}^2 \times \mathbb{P}^2$, consider the *B*-saturated *S*-ideal $J := \langle x_{1,0}, x_{1,1} \rangle \cap \langle x_{2,0}, x_{2,1} \rangle$. The minimal free resolution of S/J has the form

$$S \longleftarrow S(-1,-1)^4 \longleftarrow \frac{S(-2,-1)^2}{\bigoplus} \longleftarrow S(-2,-2) \longleftarrow 0.$$

Although the codimension of every associated prime of J is 2, there is no virtual resolution of S/J of length 2. If we had such a free complex $F = [F_0 \longleftarrow F_1 \longleftarrow F_2 \longleftarrow 0]$, then Corollary 2.5 would imply that F is a minimal free resolution of S/J, which would be a contradiction.

3. CASTELNUOVO-MUMFORD REGULARITY AND VIRTUAL RESOLUTIONS

In this section, we refine our results about short virtual resolutions. By exploiting multigraded Castelnuovo–Mumford regularity, we produce short virtual resolutions where the degrees of the generators of the free modules satisfy explicit bounds. Better yet, we obtain a converse, by showing that the set of virtual resolutions of a module determine its regularity.

Following [MS04, Definition 1.1], a finitely-generated \mathbb{Z}^r -graded B-saturated S-module M is m-regular, for some $m \in \mathbb{Z}^r$, if $H_B^i(M)_p = 0$ for all $i \ge 1$ and all $p \in \bigcup (m - q + \mathbb{N}^r)$, where the union is over all $q \in \mathbb{N}^r$ such that |q| = i - 1. The (*multigraded Castelnuovo–Mumford*) *regularity of* M is reg $M := \{p \in \mathbb{Z}^r : M \text{ is } p\text{-regular}\}$. Let $\Delta_i \subset \mathbb{Z}^r$ denote the set of twists of the summands in the i-th step of the minimal free resolution of the irrelevant ideal B.

Theorem 3.1. Let M be a finitely-generated \mathbb{Z}^r -graded B-saturated S-module. We have $\mathbf{d} \in \operatorname{reg} M$ if and only if the module $M(\mathbf{d})$ has a virtual resolution $F_0 \longleftarrow F_1 \longleftarrow \ldots \longleftarrow F_{|\mathbf{n}|} \longleftarrow 0$ such that, for all $0 \leq i \leq |\mathbf{n}|$, the degree of each generator of F_i belongs to $\Delta_i + \mathbb{N}^r$.

When r=1, we have $\Delta_i=\{-i\}$ and this theorem specializes to the existence of linear resolutions on projective space; see [Laz04, Proposition 1.8.8]. Since the minimal free resolution of S/B is a cellular resolution described explicitly in [BS98, Corollary 2.13], it follows that $\Delta_0:=\{0\}$ and that for $i\geqslant 1$, we have $\Delta_i:=\{-a\in\mathbb{Z}^r:0\leqslant a-1\leqslant n \text{ and } |a|=r+i-1\}$. We first illustrate Theorem 3.1 in the case of a hypersurface.

Example 3.2. Given a homogeneous polynomial $f \in S$ of degree d, the regularity of $S/\langle f \rangle$ has a unique minimal element e, where $e_j := \max\{0, d_j - 1\}$. As a consequence, it follows that $0 \in \operatorname{reg}(S/\langle f \rangle)$ if and only if $d_j \leq 1$ for all j.

Before proving Theorem 3.1, we need two technical lemmas.

Lemma 3.3. For $0 \le i \le |n|$, $b \in \Delta_i + \mathbb{N}^r$, and $a \in \mathbb{N}^r \setminus \{0\}$, we have $H^{|a|+i}(\mathbb{P}^n, \mathscr{O}_{\mathbb{P}^n}(b-a)) = 0$.

Proof. We induct on r. For the base case r=1, we can have nonzero higher cohomology for the given line bundle on projective space \mathbb{P}^{n_1} only if $|a|+i=a_1+i=n_1$. Since $b\in\Delta_i+\mathbb{N}=\{-i\}+\mathbb{N}$, or $b_1\geqslant -i$, we have $b_1-a_1\geqslant -i-(n_1-i)=-n_1>-n_1-1$, so the line bundle $\mathscr{O}_{\mathbb{P}^n}(b_1-a_1)$ has no higher cohomology.

For the induction step, we first consider the case where at least one entry of b-a is nonnegative, and we assume for contradiction that $H^{|a|+i}(\mathbb{P}^n, \mathscr{O}_{\mathbb{P}^n}(b-a)) \neq 0$. Since |a|+i>0, we may also assume, by reordering the factors, that the first entry of b-a is strictly negative and the last entry is nonnegative. We write $b-a=(b'-a',b_r-a_r)$ and $n=(n',n_r)$ in $\mathbb{Z}^{r-1}\oplus\mathbb{Z}$. Since $H^{|a|+i}(\mathbb{P}^n,\mathscr{O}_{\mathbb{P}^n}(b-a))\neq 0$, the Künneth formula implies that $H^{|a|+i}(\mathbb{P}^{n'},\mathscr{O}_{\mathbb{P}^{n'}}(b'-a'))\neq 0$. Decreasing the first entry of b'-a' will not alter this nonvanishing. Setting $a'':=a'+(a_r,0,0,\ldots,0)$, we obtain $a''\in\mathbb{N}^{r-1}\setminus\{0\},\ |a''|=|a'|+a_r=|a|,\ \text{and}\ H^{|a''|+i}(\mathbb{P}^{n'},\mathscr{O}_{\mathbb{P}^{n'}}(b'-a''))\neq 0$, which contradicts the induction hypothesis.

It remains to consider the case in which all entries of b-a are strictly negative. Hence, we can assume that |a|+i=|n|. The hypothesis $b \in \Delta_i + \mathbb{N}^r$ implies $|b| \geqslant -r-i+1$. Combining these yields $|b-a| = \sum_i b_i - a_i = |b| - |a| \geqslant (-r-i+1) - (|n|-i) = -|n|-r+1$. But the most positive line bundle with top-dimensional cohomology is the canonical bundle, and this inequality shows that $\mathcal{O}_{\mathbb{P}^n}(b-a)$ cannot have top dimensional cohomology.

Lemma 3.4. Let \mathscr{F} be a 0-regular $\mathscr{O}_{\mathbb{P}^n}$ -module and let $0 \leqslant a \leqslant n$. If $H^p(\mathbb{P}^n, \mathscr{F} \otimes \Omega^a(a)) \neq 0$, then we have $-a \in \Delta_{|a|-p} + \mathbb{N}^r$.

Proof. If a=0, then we have $\Omega^a(a)=\mathscr{O}_{\mathbb{P}^n}$, and the statement follows immediately from the 0-regularity of \mathscr{F} . Thus, we assume that $a\neq 0$. After possibly reordering the factors of \mathbb{P}^n , we may write $a=(a',0)\in \mathbb{Z}^{r'}\oplus \mathbb{Z}^{r-r'}$ where every entry of a' is strictly positive. For any $k\geqslant 1$, we have $(-a',0)\in \Delta_k+\mathbb{N}^r\iff (-a',-1)\in \Delta_k+\mathbb{N}^r\iff |a'|+(r-r')\leqslant k+r-1$. Setting k=|a|-p=|a'|-p establishes that $-a=(-a',0)\in \Delta_{|a|-p}+\mathbb{N}^r$ is equivalent to p< r'.

We next use truncated Koszul complexes to build a locally free resolution of $\Omega^{a}_{\mathbb{P}^{n}}(a)$. For j > r', we have $a_{j} = 0$ and $\Omega^{a_{j}}_{\mathbb{P}^{n_{j}}}(a_{j}) \cong \mathscr{O}_{\mathbb{P}^{n_{j}}}$. For $1 \leqslant j \leqslant r'$, the truncated Koszul complex twisted by $\mathscr{O}_{\mathbb{P}^{n_{j}}}(a_{j})$, namely

$$\mathscr{O}_{\mathbb{P}^{n_j}}(-1)^{\binom{n_j+1}{a_j+1}}\longleftarrow \mathscr{O}_{\mathbb{P}^{n_j}}(-2)^{\binom{n_j+1}{a_j+2}}\longleftarrow \cdots \longleftarrow \mathscr{O}_{\mathbb{P}^{n_j}}(-n_j-1+a_j)^{\binom{n_j+1}{n_j+1}}\longleftarrow 0,$$

resolves $\Omega^{a_j}_{\mathbb{P}^{n_j}}(a_j)$. Taking external tensor products gives a locally-free resolution \mathscr{G} of $\Omega^{\boldsymbol{a}}_{\mathbb{P}^n}(\boldsymbol{a})$. Any summand $\mathscr{O}_{\mathbb{P}^n}(\boldsymbol{c})$ in \mathscr{G}_i has the form $\boldsymbol{c}=(\boldsymbol{c}',0)\in\mathbb{Z}^{r'}\oplus\mathbb{Z}^{r-r'}$, where $|\boldsymbol{c}'|=-r'-i$. Tensoring the locally-free resolution \mathscr{G} with \mathscr{F} is a resolution of $\mathscr{F}\otimes\Omega^{\boldsymbol{a}}_{\mathbb{P}^n}(\boldsymbol{a})$.

Since $H^p(\mathscr{F} \otimes \Omega^a_{\mathbb{P}^n}(a)) \neq 0$, breaking the resolution $\mathscr{F} \otimes \mathscr{G}$ into short exact sequences implies that, for some index i, we have $H^{p+i}(\mathscr{F} \otimes \mathscr{G}_i) \neq 0$. Hence, there exists c = (c', 0) with |c'| = -r' - i

such that $H^{p+i}ig(\mathbb{P}^n,\mathscr{F}(c)ig)\neq 0$. Since \mathscr{F} is 0-regular, we have that |c|=|c'|<-(p+i). Therefore, we conclude that p<-|c'|-i=(r'+i)-i=r' and $a\in\Delta_{|a|-p}+\mathbb{N}^r$.

Proof of Theorem 3.1. Assume that M(d) has a virtual resolution F of the specified form. Since M is B-saturated, it suffices to show that $H^{|a|}(\mathbb{P}^n, \widetilde{M}(d-a)) = 0$ for all $a \in \mathbb{N}^r - \{0\}$. By splitting up F into short exact sequences, it suffices to show that $H^{|a|+i}(\mathbb{P}^n, \widetilde{F}_i(-a)) = 0$ for all $a \in \mathbb{N}^r \setminus \{0\}$. This is the content of Lemma 3.3.

For the converse, let \mathscr{K} denote the locally-free resolution of the diagonal $\mathbb{P}^n \hookrightarrow \mathbb{P}^n \times \mathbb{P}^n$ described in Lemma 2.1. Let p_1 and p_2 be the projections onto the first and second factors of $\mathbb{P}^n \times \mathbb{P}^n$ respectively. The sheaf $\widetilde{M}(d)$ is quasi-isomorphic to the complex $\mathscr{F} = \mathbf{R} \, p_{1*}(p_2^* \widetilde{M}(d) \otimes \mathscr{K})$, where

$$\mathscr{F}_j = igoplus_{|m{a}|-p=j} H^pig(\mathbb{P}^{m{n}}, \widetilde{M}(m{d}) \otimes \Omega^{m{a}}_{\mathbb{P}^{m{n}}}(m{a})ig) \otimes \mathscr{O}_{\mathbb{P}^{m{n}}}(-m{a}).$$

Lemma 3.4 says that $H^p(\mathbb{P}^n, \widetilde{M}(d) \otimes \Omega^a_{\mathbb{P}^n}(a)) \neq 0$ only if $-a \in \Delta_{|a|-p} + \mathbb{N}^r = \Delta_j + \mathbb{N}^r$. Since each \mathscr{F}_j is a sum of line bundles, the corresponding *S*-module F_j is free. It follows that the complex $F := [F_0 \longleftarrow F_1 \longleftarrow \cdots \longleftarrow F_{|n|} \longleftarrow 0]$ is a virtual resolution of M(d) with the desired form. \square

4. SIMPLER VIRTUAL RESOLUTIONS

We describe, in this section, an effective method for producing interesting virtual resolutions of a given S-module. Our construction depends on a B-saturated module M as well as an element $d \in \text{reg} M$. Although Theorem 4.1 defines the corresponding virtual resolution as a subcomplex of a minimal free resolution of M, Algorithm 4.4 shows that they can be calculated without first computing the entire minimal free resolution.

Theorem 4.1. For a finitely generated \mathbb{Z}^r -graded S-module M, consider a minimal free resolution F of M. For a degree $\mathbf{d} \in \mathbb{Z}^r$ and each i, let G_i be the direct sum of all free summands of F_i whose generator is in degree at most $\mathbf{d} + \mathbf{n}$, and let φ_i be the restriction of the i-th differential of F to G_i .

- (i) For all i, we have $\varphi_i(G_i) \subseteq G_{i-1}$ and $\varphi_i \circ \varphi_{i+1} = 0$, so G forms a free complex.
- (ii) Up to isomorphism, G depends only on M and d.
- (iii) If M is B-saturated and $d \in \text{reg} M$, then G is a virtual resolution of M.

When *M* is *B*-saturated and $d \in \text{reg} M$, the complex *G* is the *virtual resolution of the pair* (M, d).

Proof of Theorem 1.3. This theorem is simply a restatement of part (iii) of Theorem 4.1. \Box

To illustrate the basic idea behind the proof of Theorem 4.1, we revisit our first example.

Example 4.2. Let C be the hyperelliptic curve in $\mathbb{P}^1 \times \mathbb{P}^2$ defined by the ideal I in Example 1.4. The free complex in (1.4.2) is the virtual resolution of the pair (S/I, (2,1)) and it is naturally a subcomplex of the minimal free resolution (1.4.1) of S/I. The corresponding quotient complex E is

$$S(-1,-5)^{3} \oplus S(-2,-5)^{6} \qquad S(-3,-5)^{3} \oplus S(-3,-5)^{3} \oplus S(-3,-7)^{1} \longleftrightarrow S(-2,-7)^{2} \longleftrightarrow S(-3,-7)^{1} \longleftrightarrow 0.$$

$$S(0,-8)^{1} \oplus G(-1,-8)^{2} \qquad S(-2,-8)^{1}$$

Restricting attention to the terms of degree (*, -8), we have

$$S(0,-8)^1 \longleftarrow S(-1,-8)^2 \longleftarrow S(-2,-8)^1 \longleftarrow 0,$$

which looks like a twist of the Koszul complex on $x_{1,0}$ and $x_{1,1}$. In fact, the (*,-8), (*,-7), and (*,-5) strands each appear to have homology supported on the irrelevant ideal. This suggests that the complex \widetilde{E} is quasi-isomorphic to zero, and that is what we show in the proof of Theorem 4.1. \diamond

Lemma 4.3. Let \mathscr{E} be a bounded complex of coherent $\mathscr{O}_{\mathbb{P}^n}$ -modules. If $\mathscr{E} \otimes \mathscr{O}_{\mathbb{P}^n}(b)$ has no hypercohomology for all $0 \leq b \leq n$, then \mathscr{E} is quasi-isomorphic to 0.

Proof. By [EES15, Theorem 1.1], any bounded complex of coherent $\mathcal{O}_{\mathbb{P}^n}$ -modules is quasi-isomorphic to a Beilinson monad whose terms involve the hypercohomology evaluated at the line bundles of the form $0 \le b \le n$. The hypothesis on vanishing hypercohomology ensures that this Beilinson monad of \mathscr{E} is the 0 monad, and hence \mathscr{E} is quasi-isomorphic to 0. While [EES15, Theorem 1.1] is stated for a sheaf, the authors remark in Equation (1) on page 8 that a similar statement holds for bounded complexes of coherent sheaves.

Proof of Theorem 4.1. For part (i), write $F_i = G_i \oplus E_i$ for each i. Each generating degree e of E_i satisfies $e \not \leq d + n$. It follows that, for degree reasons, there are no nonzero maps from G_i to E_{i-1} . The i-th differential $\partial_i \colon F_i \to F_{i-1}$ has a block decomposition

$$\partial_i = egin{array}{cc} G_i & E_i \ G_{i-1} \left[egin{array}{cc} arphi_i & * \ 0 & * \end{array}
ight],$$

so $\varphi_i(G_i) \subseteq G_{i-1}$ and $\partial_i \circ \partial_{i+1} = 0$ implies that $\varphi_i \circ \varphi_{i+1} = 0$. As G only depends on F and G, part (ii) follows from the fact that the minimal free resolution of G is unique up to isomorphism. For part (iii), we may without loss of generality replace G by G and G by G. Let G be the virtual resolution of the pair G and consider the short exact sequence of complexes $G \to G \to F \to E \to 0$. It suffices to show that the complex G of sheaves is quasi-isomorphic to zero.

Fix some b, where $0 \le b \le n$. If $a \le n$, then the line bundle $\mathscr{O}_{\mathbb{P}^n}(-a+b)$ has no global sections. It follows that each summand of $\widetilde{F}(b)$ with global sections belongs to $\widetilde{G}(b)$. If $H^i(\mathbb{P}^n, \widetilde{F}(b))$ is the complex obtained by applying the functor $\mathscr{F} \mapsto H^i(\mathbb{P}^n, \mathscr{F})$ to the complex $\widetilde{F}(b)$, then we have $H^0(\mathbb{P}^n, \widetilde{F}(b)) = H^0(\mathbb{P}^n, \widetilde{G}(b))$. The notation $H^i(\mathbb{P}^n, \widetilde{F}(b))$ should not be confused with the hypercohomology group $\mathbb{H}^i(\mathbb{P}^n, \widetilde{F}(b))$, which equals $H^i(\mathbb{P}^n, \widetilde{M}(b))$ because \widetilde{F} is a locally-free resolution of the sheaf \widetilde{M} . Since $0 \in \operatorname{reg} M$ and $b \ge 0$, the Hilbert polynomial and Hilbert function of M agree in degree b. Because F is a minimal free resolution of M, it follows that the strand $[F]_b := [(F_0)_b \longleftarrow (F_1)_b \longleftarrow \cdots]$ is quasi-isomorphic to M_b , and hence

$$\mathbb{H}^0\big(\mathbb{P}^n,\widetilde{F}(\boldsymbol{b})\big)=M_{\boldsymbol{b}}\cong [F]_{\boldsymbol{b}}=H^0\big(\mathbb{P}^n,\widetilde{F}(\boldsymbol{b})\big)\cong H^0\big(\mathbb{P}^n,\widetilde{G}(\boldsymbol{b})\big).$$

If the line bundle $\mathscr{O}_{\mathbb{P}^n}(-a+n)$ has global sections, then we see that $\mathscr{O}_{\mathbb{P}^n}(-a+b)$ has no higher cohomology. Therefore, the only summands in \widetilde{F} that can potentially have higher cohomology are those that also appear in \widetilde{E} . Thus, for all i>0, we have $H^i(\mathbb{P}^n,\widetilde{F}(b))=H^i(\mathbb{P}^n,\widetilde{E}(b))$ and

 $H^i(\mathbb{P}^n, \widetilde{G}(b)) = 0$. It follows that $\mathbb{H}^0(\mathbb{P}^n, \widetilde{G}(b)) \cong H^0(\mathbb{P}^n, \widetilde{G}(b))$ and $\mathbb{H}^i(\mathbb{P}^n, \widetilde{G}(b)) = 0$ for all i > 0. Hence, the long exact sequence in hypercohomology yields

$$\mathbb{H}^iig(\mathbb{P}^n,\widetilde{E}(oldsymbol{b})ig) = egin{cases} 0 & ext{if } i=0, \ \mathbb{H}^iig(\mathbb{P}^n,\widetilde{F}(oldsymbol{b})ig) & ext{if } i>0. \end{cases}$$

Since $b \in \operatorname{reg} M$, the sheaf $\widetilde{M}(b)$ has no higher cohomology and $\widetilde{F}(b)$ has no higher hypercohomology. By Lemma 4.3, we conclude that \widetilde{E} is quasi-isomorphic to 0.

Although Theorem 4.1 presents the virtual resolution of the pair (M, d) as a subcomplex of a minimal free resolution, the following algorithm shows that we can compute a virtual resolution of the pair (M, d) without first computing an entire minimal free resolution. For a module M and a degree $d \in \mathbb{Z}^r$, let $M_{\geq d}$ denote the module $\bigoplus_{a \geq d} M_a$.

Algorithm 4.4 (Computing Virtual Resolutions of a Pair).

Input: A finitely generated \mathbb{Z}^r -graded B-saturated S-module M and

a vector $d \in \mathbb{Z}^r$ such that $d \in \operatorname{reg} M$.

Output: The virtual resolution G of the pair (M, d).

Initialize K := M and i := 0;

While $K \neq 0$ do

Choose a homogeneous minimal set \mathcal{G} of generators for K;

Initialize $G_i := \bigoplus_{g \in G} S(-\deg(g))$ and $\varphi_i : G_i \to K$ to be the corresponding surjection;

Set $K := (\operatorname{Ker} \varphi_i)_{\geqslant d+n}$;

Set i := i + 1;

Return $G := [G_0 \stackrel{\varphi_1}{\longleftarrow} G_1 \stackrel{\varphi_2}{\longleftarrow} G_2 \longleftarrow \cdots].$

Proof of Correctness. Let G be the complex produced by the algorithm, let F be the minimal free resolution of M, and let G' be the virtual resolution of (M, d). Let φ , ∂ , and ψ be the differentials of G, F, and G' respectively. We have $G_0 = F_0 = G'_0$, as $d \in \operatorname{reg} M$ implies that M is generated in degree at most d by [MS04, Theorem 1.3].

The definition of G implies that $(\operatorname{Im} \partial_i)_{\leq d+n}$ equals $\operatorname{Im} \psi_i$. We use induction on i to prove that $\operatorname{Im} \phi_i = (\operatorname{Im} \partial_i)_{\leq d+n}$. When i = 1, we have $\operatorname{Im} \phi_1 = (\operatorname{Im} \partial_1)_{\leq d+n}$. For i > 1, the key observation is

$$(\operatorname{Im} \partial_i)_{\leqslant d+n} = (\operatorname{Ker} \partial_{i-1})_{\leqslant d+n} = (\operatorname{Ker} \partial_{i-1}|_{G_{i-1}})_{\leqslant d+n},$$

where the righthand equality holds because any element in $(\text{Ker } \partial_{i-1})_{\leqslant d+n}$ only depends on the restriction of ∂_{i-1} to G_{i-1} . By induction, we have $\text{Im } \varphi_{i-1} = (\text{Im } \partial_{i-1})_{\leqslant d+n}$, so

$$\left(\operatorname{Ker} \partial_{i-1}|_{G_{i-1}}\right)_{\leq d+n} = \left(\operatorname{Ker} \varphi_{i-1}\right)_{\leq d+n} = \operatorname{Im} \varphi_i.$$

Therefore, we conclude that $\operatorname{Im} \varphi_i = \operatorname{Im} \psi_i$ for all i and $G \cong G'$.

Example 4.5. Let $Z \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^2$ be the subscheme consisting of 6 general points and let I be the corresponding B-saturated S-ideal. Macaulay2 [M2] shows that the minimal free resolution of S/I has the form $S^1 \longleftarrow S^{37} \longleftarrow S^{120} \longleftarrow S^{166} \longleftarrow S^{120} \longleftarrow S^{45} \longleftarrow S^7 \longleftarrow 0$, where for brevity we have omitted the twists. Using [MS04, Proposition 6.7], it follows that, up to symmetry in the

first two factors, the minimal elements in the regularity of S/J are (5,0,0),(2,1,0),(1,0,1), and (0,0,2). Table 4.5.1 compares some basic numerical invariants for the minimal free resolution and the corresponding virtual resolutions. The total Betti numbers of a free complex F are the ranks of the terms F_i ignoring the twists. Since Z has codimension 4, part (i) of Proposition 2.4 implies that

1	1	
Type of Free Complex	Total Betti Numbers	Number of Twists
minimal free resolution of S/I	(1,37,120,166,120,45,7)	78
virtual resolution of the pair $(S/J, (5,0,0))$	(1,24,50,33,6)	18
virtual resolution of the pair $(S/J, (2, 1, 0))$	(1,29,73,66,21)	22
virtual resolution of the pair $(S/J, (1,0,1))$	(1,25,63,57,18)	15
virtual resolution of the pair $(S/J, (0,0,2))$	(1,22,51,42,12)	13

TABLE 4.5.1. Comparison of various free complexes associated to Z

any virtual resolution for S/I must have length at least 4, so the minimum is achieved by all of these virtual resolutions. All four virtual resolutions also have a nonzero first homology module, which is supported on the irrelevant ideal. The first three virtual resolutions also have nonzero second homology modules. By examining the twists, we see that no pair of these virtual resolutions are comparable. This corresponds to the fact that the reg(S/I) has several distinct minimal elements. \diamond

5. VIRTUAL RESOLUTIONS FOR PUNCTUAL SCHEMES

This section formulates and proves an extension of a property of points in projective space. While every punctual scheme in projective space is arithmetically Cohen–Macaulay, this fails when the ambient space is a product of projective spaces; the minimal free resolution is nearly always too long. However, by using virtual resolutions, we obtain a unexpected variant for points in \mathbb{P}^n .

To state this analogue, we introduce some notation. The irrelevant ideal B on \mathbb{P}^n can be written as $B = \bigcap_{i=1}^r \langle x_{i,0}, x_{i,1}, \dots, x_{i,n_i} \rangle^{a_i}$. For a vector $a \in \mathbb{N}^r$, we set $B^a := \bigcap_{i=1}^r \langle x_{i,0}, x_{i,1}, \dots, x_{i,n_i} \rangle^{a_i}$.

Theorem 5.1. If $Z \subset \mathbb{P}^n$ is a zero-dimensional scheme and I is the corresponding B-saturated S-ideal, then there exists $\mathbf{a} \in \mathbb{N}^r$ with $a_r = 0$ such that the minimal free resolution of $S/(I \cap B^a)$ has length equal $|\mathbf{n}| = \dim \mathbb{P}^n$. Moreover, any $\mathbf{a} \in \mathbb{N}^r$ with $a_r = 0$ and other entries sufficiently positive yields such a virtual resolution of S/I.

Proof of Theorem 1.5. Applying Theorem 5.1, it suffices to choose $Q = B^a$ for any $a \in \mathbb{N}^r$ with $a_r = 0$ and other entries sufficiently positive.

While Theorem 5.1 establishes that, for appropriate $a \in \mathbb{N}^r$, the projective dimension of $S/(I \cap B^a)$ equals the codimension of Z, this does not mean that the algebra $S/(I \cap B^a)$ is Cohen–Macaulay; the ideal $I \cap B^a$ will often fail to be unmixed. For instance, on $\mathbb{P}^2 \times \mathbb{P}^2$, the ideals $\langle x_{i,0}, x_{i,1}, x_{i,2} \rangle$ for $1 \le i \le 2$ have codimension 3 whereas a zero-dimensional scheme Z would have codimension 4. Nevertheless, we do get Cohen–Macaulayness in one case.

Corollary 5.2. If $Z \subset \mathbb{P}^1 \times \mathbb{P}^1$ is a zero-dimensional subscheme and I is the corresponding B-saturated S-ideal, then there exists an ideal Q whose radical is $\langle x_{1,0}, x_{1,1} \rangle$ such that

(i) the algebra $S/(I \cap Q)$ is Cohen–Macaulay, and

(ii) there exists an $(m+1) \times m$ matrix over S whose maximal minors generate $I \cap Q$.

Proof. Theorem 5.1 yields an $a \in \mathbb{N}^r$ such that $I \cap B^a$ has projective dimension 2. On $\mathbb{P}^1 \times \mathbb{P}^1$, the irrelevant ideal B also has codimension 2, so $S/(I \cap B^a)$ has codimension 2. Thus, the algebra $S/(I \cap B^a)$ is Cohen–Macaulay. The second statement is an immediate consequence of the Hilbert–Burch Theorem [Eis95, Theorem 20.15].

Remark 5.3. Although this paper focuses on products of projective spaces, our proofs for both Theorem 5.1 and Corollary 5.2 can be adapted to hold in the more general context of iterated projective bundles. For instance, let X be the Hirzebruch surface with Cox ring $S = \mathbb{k}[y_0, y_1, y_2, y_3]$ where the variables have degrees (1,0), (1,0), (-2,1), and (0,1) respectively. Let $Z \subset X$ be the scheme-theoretic intersection of $y_0^5 y_2^2 + y_1 y_3^2$ and $y_0 y_1 + y_2 y_1^3$. If I is the B-saturated S-ideal of Z, then S/I has projective dimension 3 and $S/(I \cap \langle y_0, y_1 \rangle^a)$ has projective dimension 2 for any $a \ge 4$.

As in Section 3, we collect two lemmas before proving the main result of this section.

Lemma 5.4. If $Z \subset \mathbb{P}^n$ is a zero-dimensional scheme and I is the corresponding B-saturated S-ideal, then there exists $\mathbf{a} \in \mathbb{Z}^r$ with $a_r = 0$ such that the depth of $(S/I)_{\geqslant \mathbf{a}}$ is r. Moreover, this holds for any $\mathbf{a} \in \mathbb{Z}^r$ with $a_r = 0$ and other entries sufficiently positive.

Proof. Since extending the ground field does not change the depth of a module, we assume that \mathbb{k} is an infinite field. We have $\operatorname{codim}(S/I)_{\geqslant a} = \operatorname{codim}(S/I) = r$, so the depth of S/I is bounded above by r. For each $1 \leqslant i \leqslant r$, choose a general linear element ℓ_i in $\langle x_{i,0}, x_{i,1}, \ldots, x_{i,n_i} \rangle$. We claim that the elements $\ell_1, \ell_2, \ldots, \ell_r$ form a regular sequence on $(S/I)_{\geqslant a}$.

Let $M := \bigoplus_{b \in \mathbb{N}^r} H^0(Z, \mathcal{O}_Z(b))$. By construction, the elements $\ell_1, \ell_2, \dots, \ell_r$ form a regular sequence on M. Since I is B-saturated, it follows that $H^0_B(S/I) = 0$. The exact sequence relating local cohomology and sheaf cohomology [CLS11, Theorem 9.5.7] gives

$$0 \longleftarrow (H_B^1(S/I))_{\geqslant 0} \longleftarrow M \longleftarrow S/I \longleftarrow 0$$

$$\downarrow \cdot \ell_i \qquad \qquad \downarrow \cdot \ell_i \qquad \qquad \downarrow \cdot \ell_i$$

$$0 \longleftarrow (H_B^1(S/I))_{\geqslant 0} \longleftarrow M \longleftarrow S/I \longleftarrow 0.$$

The middle vertical arrow is an isomorphism because ℓ_i does not vanish on any point in Z. Hence, the Snake Lemma [Eis95, Exercise A3.10] implies that the right vertical arrow is injective.

Focusing on the last component of \mathbb{Z}^r , we identify the Cox ring $R:=\Bbbk[x_{r,0},x_{r,1},\ldots,x_{r,n_r}]$ of the factor \mathbb{P}^{n_r} with the subring $(S)_{(0,*)}:=\bigoplus_{\alpha\in\mathbb{N}}(S)_{(0,\alpha)}$ of S. For any $c'\in\mathbb{Z}^{r-1}$, consider the R-module $(S/I)_{(c',*)}:=\bigoplus_{\alpha\in\mathbb{N}}(S)_{(c',\alpha)}$. These modules form a directed set: for $c',c''\in\mathbb{Z}^{s-1}$ with $c''\geqslant c'$, multiplication by the form $\ell_1^{c''_1-c'_1}\ell_2^{c''_2-c'_2}\cdots\ell_{s-1}^{c''_{s-1}-c'_{s-1}}$ gives the inclusion $(S/I)_{(c',*)}\subseteq (S/I)_{(c',*)}$. Each R-module $(S/I)_{(c',*)}$ is a submodule of $(M)_{(c',*)}$ and $(M)_{(c',*)}\cong (M)_{(0,*)}$. It follows that the $(S/I)_{(c',*)}$ form an increasing sequence of finitely-generated R-submodules of $(M)_{(0,*)}$, so this sequence stabilizes. In particular, if $a'\in\mathbb{Z}^{r-1}$ is sufficiently positive, then the inclusion $(S/I)_{(c',*)}\subseteq (S/I)_{(c'+e_i,*)}$ is an isomorphism for each $c'\geqslant a'$ and each $1\leqslant i\leqslant r-1$. Hence,

 $\ell_1, \ell_2, \dots, \ell_{r-1}$ form a regular sequence on $(S/I)_{\geqslant (\boldsymbol{a}',0)}$ and

$$\frac{(S/I)_{\geqslant (\boldsymbol{a}',0)}}{\langle \ell_1, \ell_2, \dots, \ell_{r-1} \rangle} \cong (S/I)_{(\boldsymbol{a}',*)}.$$

Since *I* is *B*-saturated, ℓ_r is regular on S/I, so it is also regular on the *R*-module $(S/I)_{(a',*)}$. Setting a = (a', 0), the Auslander–Buchsbaum Formula [Eis95, Theorem 19.9] completes the proof.

Remark 5.5. The proof of Lemma 5.4 shows that $(S/I)_{\geqslant a}$ has a multigraded regular sequence of length r, but neither $S/(I \cap B^a)$ nor S/B^a generally has a multigraded regular sequence of length r.

Lemma 5.6. If $\mathbf{a} := (a_1, a_2, \dots, a_i, 0, \dots, 0) \in \mathbb{Z}^r$ for some $1 \le i \le r$, then the projective dimension of S/B^a is at most $n_1 + n_2 + \dots + n_i + 1$. Moreover, this holds for any $\mathbf{a} \in \mathbb{Z}^r$ with $a_j = 0$ for $i+1 \le j \le r$ and other entries sufficiently positive.

Proof. We proceed by induction on *i*. The base case i=1 is just the Hilbert Syzygy Theorem [Eis95, Theorem 1.3] applied to the Cox ring of \mathbb{P}^{n_1} . When i>1, set $\mathbf{a}':=(a_1,a_2,\ldots,a_{i-1},0,\ldots,0)$ and $\mathbf{a}'':=(0,\ldots,0,a_i,0,\ldots,0)$, so that $\mathbf{a}=\mathbf{a}'+\mathbf{a}''$ and $\mathbf{a}^a=\mathbf{a}^a'\cap\mathbf{a}^a$. The short exact sequence $0 \longleftarrow S/(B^{a'}+B^{a''}) \longleftarrow S/B^{a'} \longleftarrow S/B^a \longleftarrow 0$ yields

$$(5.6.4) pdim(S/B^{a}) \leq \max\{pdim(S/B^{a'}), pdim(S/B^{a''}), pdim(S/(B^{a'}+B^{a''})) - 1\}.$$

By induction, the projective dimension of $S/B^{a'}$ is at most $n_1 + n_2 + \cdots + n_{i-1} + 1$. By the Hilbert Syzygy Theorem on \mathbb{P}^{n_i} , the projective dimension of $S/B^{a''}$ is at most $n_i + 1$. Since $B^{a'}$ and $B^{a''}$ are supported on disjoint sets of variables, $\operatorname{pdim}(S/(B^{a'} + B^{a''})) = \operatorname{pdim}(S/B^{a'}) + \operatorname{pdim}(S/B^{a''})$ which is at most $n_1 + n_2 + \cdots + n_i + 2$. Applying (5.6.4), we obtain the result.

Proof of Theorem 5.1. Let $a \in \mathbb{Z}^r$ with $a_r = 0$ and other entries sufficiently positive. There is a short exact sequence $0 \longleftarrow S/B^a \longleftarrow S/(I \cap B^a) \longleftarrow (S/I)_{\geqslant a} \longleftarrow 0$. By Proposition 2.4, it suffices to prove that the projective dimension of $S/(I \cap B^a)$ is at most $n_1 + n_2 + \cdots + n_r$. Lemma 5.4 shows that $(S/I)_{\geqslant a}$ has projective dimension $n_1 + n_2 + \cdots + n_r$ and Lemma 5.6 shows that S/B^a has projective dimension at most $n_1 + n_2 + \cdots + n_{r-1} + 1$. It follows that the projective dimension of $S/(I \cap B^a)$ is at most $n_1 + n_2 + \cdots + n_r$ as well.

The next example contrasts the virtual resolutions produced by Theorem 1.3 and Theorem 1.5.

Example 5.7. As in Example 4.5, let $Z \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^2$ be the subscheme consisting of 6 general points and let I be the corresponding B-saturated S-ideal. Table 5.7.2 compares some basic numerical invariants for virtual resolutions arising from Theorem 5.1. Since the virtual resolutions

TABLE 5.7.2. Comparison of various free complexes associated to Z

Type of Free Complex	Total Betti Numbers	Number of Twists
minimal free resolution of S/I	(1,37,120,166,120,45,7)	78
virtual resolution from $a = (2, 1, 0)$	(1,17,34,24,6)	12
virtual resolution from $a = (3,3,0)$	(1,22,42,27,6)	13

in Table 5.7.2 involve non-minimal generators for I, they are different than those in Table 4.5.1.

Conversely, the virtual resolutions appearing in Table 4.5.1 cannot be obtained from Theorem 1.5 because those free complexes are not acyclic.

We end this section by extending Corollary 5.2 to any smooth projective toric surface.

Proposition 5.8. Fix a smooth projective toric surface X. Let $Z \subset X$ be the subscheme consisting of m general points and let I be the corresponding B-saturated S-ideal. There exists a virtual resolution $F := [S \longleftarrow F_1 \xleftarrow{\varphi} F_2 \longleftarrow 0]$ of S/I such that $\operatorname{rank}(F_1) = \operatorname{rank}(F_2) + 1$, the maximal minors of φ generate an ideal J with $I = (J : B^{\infty})$, and S/J is a Cohen–Macaulay algebra.

Proof. Any smooth projective toric surface can be realized as a blowup of $\pi: X \to Y$, where Y is \mathbb{P}^2 or a Hirzebruch surface [CLS11, Theorem 10.4.3]. Since $\pi(Z) \subset Y$ is a punctual scheme, we can apply the Hilbert–Burch Theorem when Y is \mathbb{P}^2 , or Corollary 5.2 and Remark 5.3 when Y is a Hirzebruch surface, to obtain a resolution of $\mathscr{O}_{\pi(Z)}$ of the form $\mathscr{O}_Y \longleftarrow \mathscr{E}_1 \longleftarrow \mathscr{E}_2 \longleftarrow 0$, where \mathscr{E}_1 and \mathscr{E}_2 are sums of line bundles on Y. Our genericity hypothesis implies that Z does not intersect the exceptional locus of π , so $\pi^*\mathscr{O}_Y \longleftarrow \pi^*\mathscr{E}_1 \longleftarrow \pi^*\mathscr{E}_2 \longleftarrow 0$ is a locally-free resolution of \mathscr{O}_Z . The corresponding complex of S-modules is a virtual resolution for S/I of the appropriate form. \square

Example 5.9. Consider the del Pezzo surface X of degree 7 or equivalently the smooth Fano toric surface obtained by blowing-up the projective plane at two torus-fixed points. The Cox ring of X is $S := \mathbb{k}[y_0, y_1, \dots, y_4]$ equipped with the \mathbb{Z}^3 -grading induced by

$$\deg(y_0) := \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad \deg(y_1) := \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \quad \deg(y_2) := \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \quad \deg(y_3) := \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}, \quad \deg(y_4) := \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

With this choice of basis for Pic(X), the nef cone equals the positive orthant. Let $Z \subset X$ be the subscheme consisting of the three points [1:1:1:1:1], [2:1:3:1:5], and [7:1:11:1:13] (expressed in Cox coordinates) and let I be the corresponding B-saturated S-ideal. Macaulay2 [M2] shows that the minimal free resolution of S/I has the form

However, there is a virtual resolution of S/I having the form

$$S^1 \leftarrow S(0,-2,0)^3 \leftarrow S(0,-3,0)^2 \leftarrow 0.$$

Example 5.10. Let $Z \subset \mathbb{P}^1 \times \mathbb{P}^1$ be the subscheme consisting of m general points and let I be the corresponding B-saturated S-ideal. Not only is there a Hilbert–Burch-type virtual resolution of S/I, it can be chosen to be a Koszul complex. Since $\dim H^0\left(\mathbb{P}^1 \times \mathbb{P}^1, \mathscr{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(i,j)\right) = (i+1)(j+1)$, the generality of the points implies that $\dim H^0\left(\mathbb{P}^1 \times \mathbb{P}^1, \mathscr{O}_Z(i,j)\right) = \min\{(i+1)(j+1), m\}$. Hence, if m = 2k for some $k \in \mathbb{N}$, then two independent global sections of $\mathscr{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1,k)$ vanish on Z. Using

this pair, we obtain a virtual resolution of S/I of the form $S \longleftarrow S(-1,-k)^2 \longleftarrow S(-2,-2k) \longleftarrow 0$. On the other hand, if m=2k+1, then there are independent global sections of $\mathscr{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1,k)$ and $\mathscr{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1,k+1)$ that vanish on Z, so we obtain a virtual resolution of S/I having the form

$$S \longleftarrow \frac{S(-1,-k)}{S(-1,-k-1)} \longleftarrow S(-2,-2k-1) \longleftarrow 0.$$

Question 5.11. Does Proposition 5.8 hold for any punctual scheme Z in a smooth toric surface X?

6. GEOMETRIC APPLICATIONS

In this section, we showcase four geometric applications of virtual resolutions. In particular, each of these support our overarching thesis that replacing minimal free resolutions by virtual resolutions yields the best geometric results for subschemes of \mathbb{P}^n .

Unmixedness. Given a subscheme that has a virtual resolution whose length equals its codimension, we prove an unmixedness result. This extends the classical unmixedness result for arithmetically Cohen–Macaulay subschemes, see [Eis95, Corollary 18.14].

Proposition 6.1 (Unmixedness). Let $Z \subset \mathbb{P}^n$ be a closed subscheme of codimension c and let I be the corresponding B-saturated S-ideal. If S/I has a virtual resolution of length c, then every associated prime of I has codimension c.

Proof. Let Q be an associated prime of I and let F denote a virtual resolution of S/I having length c. Our hypothesis on Z implies that $\operatorname{codim} Q \geqslant c$. Since Q does not contain the irrelevant ideal B, localizing at Q annihilates the homology of F that is supported at B. Thus, the complex F_Q is a free S_Q -resolution of $(S/I)_Q$. Since the projective dimension of a module is at least its codimension [Eis95, Proposition 18.2], it follows that $c \geqslant \operatorname{codim}_{S_Q}(S/I)_Q = \operatorname{codim} Q \geqslant c$.

Deformation Theory. Using virtual resolutions, we generalize results about unobstructed deformations for arithmetically Cohen–Macaulay subschemes of codimension two, arithmetically Gorenstein subschemes of codimension three, and complete intersections. To accomplish this, we first extend Piene and Schlessinger's Comparison Theorem [PS85] by relating the deformations of a closed subscheme $Y \subseteq \mathbb{P}^n$ with deformations of a corresponding graded module over the Cox ring.

Theorem 6.2 (Comparison Theorem). Let $Y \subset \mathbb{P}^n$ be a closed subscheme and let I be a homogeneous S-ideal defining Y scheme-theoretically and generated in degrees d_1, d_2, \ldots, d_s . If the natural map $(S/I)_{d_i} \to H^0(Y, \mathcal{O}_Y(d_i))$ is an isomorphism for all $1 \le i \le s$, then the embedded deformation theory of $Y \subset \mathbb{P}^n$ is equivalent to the degree zero embedded deformation theory of $V(I) \subset \operatorname{Spec}(S)$.

Proof. Piene and Schlessinger's proof of the Comparison Theorem [PS85] goes through essentially verbatim by replacing projective space and its coordinate ring with \mathbb{P}^n and its Cox ring S.

Proof of Theorem 1.6. If $e \in \operatorname{reg}(S/I)$ and F is the virtual resolution of the pair (S/I, e), then we have $H_0(F) = S/J$ for some ideal J whose B-saturation equals I. By Theorem 1.3, the generating degrees for J are a subset of those for I. It follows that $(S/J)_d = H^0(Y, \mathcal{O}_Y(d))$ for each degree d of a generator for J. Therefore, Theorem 6.2 implies that the embedded deformation theory of Y is equivalent to the degree zero embedded deformation theory of the subscheme $V(J) \subset \operatorname{Spec}(S)$.

- (i) The virtual resolution F has length 2, so Proposition 2.4 implies that F is the minimal free resolution of S/J. Thus, S/J is Cohen–Macaulay of codimension 2, and [Art76, §5] or [Sch77] implies that its embedded deformations are unobstructed.
- (ii) The virtual resolution F has length 3 and $\min\{n_i + 1 : 1 \le i \le r\} \ge 3$, so Proposition 2.4 implies that F is the minimal free resolution of S/J. Thus, S/J is Gorenstein of codimension 3, and [MR92, Theorem 2.1] implies that its embedded deformations are unobstructed.
- (iii) Let $c := \operatorname{codim} Y$. As F is a Koszul complex, we have $F_1 = \bigoplus_{i=1}^c S(-d_i)$. Since F is a virtual resolution of S/I, we also see that \widetilde{J} equals the ideal sheaf \mathscr{I}_Y for $Y \subset \mathbb{P}^n$. The complex $\widetilde{F}_1 \longleftarrow \widetilde{F}_2 \longleftarrow \widetilde{F}_3 \longleftarrow \cdots$ is a locally-free resolution of \mathscr{I}_Y , so the normal bundle of Y in \mathbb{P}^n is $\mathscr{N}_{Y/\mathbb{P}^n} := \mathscr{H}om(\mathscr{I}_Y/\mathscr{I}_Y^2, \mathscr{O}_Y) \cong \bigoplus_{i=1}^c \mathscr{O}_Y(d_i)$. For a fixed deformation of Y, the obstruction is a Čech cocycle in $H^1(Y, \mathscr{N}_{Y/\mathbb{P}^n})$ determined by local lifts of the syzygies; see [Har10, Theorem 6.2]. However, since Y is a scheme-theoretic complete intersection, its syzygies are all Koszul, so we can defined this cocycle by lifting those Koszul syzygies globally on Y. Hence, the Čech cocycle in $H^1(Y, \mathscr{N}_{Y/\mathbb{P}^n})$ is actually a coboundary and the obstruction vanishes.

Remark 6.3. In part (ii) of Theorem 1.6, we suspect that the hypothesis min $\{n_i\} \ge 2$ is unnecessary.

Example 6.4. Consider the hyperelliptic curve $C \subset \mathbb{P}^1 \times \mathbb{P}^2$ defined in Example 1.4. Applying part (i) of Theorem 1.6, the virtual resolution from (1.4.2) implies that C has unobstructed embedded deformations. Alternatively, this curve has a virtual resolution

$$S \xleftarrow{[f \ g]} \underbrace{S(-2,-2)}_{S(-3,-1)} \xleftarrow{\begin{bmatrix} -g \\ f \end{bmatrix}} S(-5,-3) \longleftarrow 0,$$

where $f = x_{1,0}^2 x_{2,0}^2 + x_{1,1}^2 x_{2,1}^2 + x_{1,0} x_{1,1} x_{2,2}^2$ and $g = x_{1,0}^3 x_{2,2} + x_{1,1}^3 (x_{2,0} + x_{2,1})$, so part (iii) of Theorem 1.6 provides another proof that this curve has unobstructed embedded deformations. \diamond

Regularity of Tensor Products. Using virtual resolutions, we can prove bounds for the regularity of a tensor product, similar to the bounds obtained for projective space; see [Laz04, Proposition 1.8.8]. Let e_i denote the i-th standard basis vector in $\mathbb{Z}^r = \text{Pic}(\mathbb{P}^n)$.

Proposition 6.5. Let $\mathscr E$ and $\mathscr F$ be coherent $\mathscr O_{\mathbb P^n}$ -modules such that $\mathscr T$ or $^j(\mathscr E,\mathscr F)=0$ for all j>0. If $a\in\operatorname{reg}\mathscr E$ and $b\in\operatorname{reg}\mathscr F$, then we have $a+b+1-e_i\in\operatorname{reg}(\mathscr E\otimes\mathscr F)$ for each $1\leqslant i\leqslant r$.

Proof. Let M and N be the B-saturated S-modules corresponding to $\mathscr E$ and $\mathscr F$. Since M(a) is 0-regular, Theorem 3.1 implies that it has a virtual resolution $F_0 \longleftarrow F_1 \longleftarrow \cdots$, where the degree of each generator of F_i belongs to $\Delta_i + \mathbb{N}^r$. Similarly, N(b) has a virtual resolution $G_0 \longleftarrow G_1 \longleftarrow \cdots$ satisfying the same conditions. The vanishing of Tor-groups implies that $H := F \otimes G$ is a virtual resolution of $M(a) \otimes N(b)$. Since $\Delta_i + \Delta_j + 1 - e_i \subseteq \Delta_{i+j} + \mathbb{N}^r$, it follows that the degree of each generator of the free module $H(1-e_i)_k$ belongs to $\Delta_k + \mathbb{N}^r$, for each k. Hence Theorem 3.1 implies that $(M(a) \otimes N(b))(1-e_i)$ is 0-regular.

Remark 6.6. Proposition 6.5 is sharp. When r = 1, it recovers Proposition 1.8.8 in [Laz04], as the higher Tor-groups vanish whenever one of the two sheaves is locally free. If r > 1, then it is possible to have 0-regular sheaves whose tensor product is not 0-regular. For instance, if $D, D' \subset \mathbb{P}^1 \times \mathbb{P}^2$

are degree (1,1)-hypersurfaces, then the product $\mathscr{O}_D \otimes \mathscr{O}_{D'}$ is isomorphic to the structure sheaf \mathscr{O}_C for a curve C with $H^1(C,\mathscr{O}_C(0,-1)) \neq 0$.

Remark 6.7. Definition 1.8 in [EES15] describes an asymptotic variant of Proposition 6.5.

Vanishing of Higher Direct Images. A relative notion of Castelnuovo–Mumford regularity with respect to a given morphism is defined in terms of the vanishing of derived pushforwards; see [Laz04, Example 1.8.24]. Just as virtual resolutions yield sharper bounds on multigraded Castelnuovo–Mumford regularity, they also provide sharper bounds for the vanishing of derived pushforwards. For some $1 \le s \le r$, fix a subset $\{i_1, i_2, \dots, i_s\} \subseteq \{1, 2, \dots, r\}$ and let Y be the corresponding product $\mathbb{P}^{n_{i_1}} \times \mathbb{P}^{n_{i_2}} \times \dots \times \mathbb{P}^{n_{i_r}}$ of projective spaces. The canonical projection $\pi \colon \mathbb{P}^n \to Y$ induces an inclusion $\pi^* \colon \operatorname{Pic}(Y) \to \operatorname{Pic}(\mathbb{P}^n)$ and we write $\rho \colon \operatorname{Pic}(\mathbb{P}^n) \to \operatorname{Coker}(\pi^*) = \operatorname{Pic}(\mathbb{P}^n)/\operatorname{Pic}(Y)$.

Proposition 6.8. Let M be a finitely-generated \mathbb{Z}^r -graded S-module and consider $a \in \mathbb{Z}^r$. If we have $\rho(a) \in \rho(\operatorname{reg} M)$, then it follows that $\mathbf{R}^i \pi_* \widetilde{M}(a) = 0$ for all i > 0.

Proof. Since $\rho(a) \in \rho(\text{reg } M)$, we can choose $b \in \text{Pic}(Y)$ such that $a + \pi^*b \in \text{reg } M$. The Projection Formula [Har77, Exercise III.8.3] gives $\mathbf{R}^i \pi_* \widetilde{M}(a) \otimes \mathscr{O}_Y(b) = \mathbf{R}^i \pi_* \widetilde{M}(a + \pi^*b)$ for all $b \in \text{Pic}(Y)$. Hence, by replacing a with $a + \pi^*b$, we assume that a itself lies in reg M.

Let G be the virtual resolution of the pair (M, a), and consider a summand S(-c) of G. By definition, we have $c \le a + n$. It follows that $-n \le -c + a$ and $\mathbf{R}^i \pi_* \mathscr{O}_{\mathbb{P}^n}(-c + a) = 0$ for all i > 0. From the hypercohomology spectral sequence $E_2^{p,q} := \mathbf{R}^p \widetilde{G_{-q}} \Longrightarrow \mathbf{R}^{p+q} \widetilde{M}$, we conclude that the higher direct images of $\widetilde{M}(a)$ also vanish.




FIGURE 6.8.1. Representation of reg(S/I) and its image under ρ .

Example 6.9. Let $Y \subset \mathbb{P}^1 \times \mathbb{P}^3$ be the surface defined by the *B*-saturated ideal

$$I = \left\langle \begin{array}{c} x_{1,1}x_{2,1}^2 + x_{1,0}x_{2,0}x_{2,2} + x_{1,1}x_{2,1}x_{2,3}, x_{1,0}^2x_{2,1}^2 + x_{1,0}x_{1,1}x_{2,2}^2 + x_{1,1}^2x_{2,0}x_{2,3}, x_{1,0}x_{2,1}^4 - x_{1,0}x_{2,0}x_{2,2}^3 + \\ x_{1,0}x_{2,1}^3x_{2,3} - x_{1,1}x_{2,0}^2x_{2,2}x_{2,3}, x_{2,1}^6 - x_{2,0}x_{2,1}^2x_{2,2}^3 + 2x_{2,1}^5x_{2,3} + x_{2,0}^3x_{2,2}^2x_{2,3} - x_{2,0}x_{2,1}x_{2,2}^3x_{2,3} + x_{2,1}^4x_{2,3}^2 \right\rangle,$$

and let $\pi: \mathbb{P}^1 \times \mathbb{P}^3 \to \mathbb{P}^1$ be the projection onto the first factor. To understand the vanishing of the higher direct images of \mathcal{O}_Y , we consider the minimal free resolution of S/I which has the form

If we tensor the corresponding locally-free resolution with the line bundle $\mathscr{O}_Y(0,3)$, then none of the terms in the resulting complex have nonzero higher direct images, so $\mathbf{R}^1\pi_*\mathscr{O}_Y(0,c)=0$ for $c\geqslant 3$. However, Proposition 6.8 yields a sharper vanishing result. Since Macaulay2 [M2] shows that $(1,1)\in \operatorname{reg}(S/I)$, we have $\mathbf{R}^1\pi_*\mathscr{O}_Y(0,c)=0$ for all $c\geqslant 1$. This bound is sharp because a general fiber of π is a curve of genus 1.

7. QUESTIONS

We expect that virtual resolutions will produce further analogues of theorems involving minimal free resolutions on projective space. We close by highlighting several promising directions.

The first question is to find a notion of depth that controls the minimal length of a virtual resolution and provides an analogue of the Auslander–Buchsbaum Theorem.

Question 7.1. Given an S-module M, what invariants of M determine the length of the shortest possible virtual resolution of M?

Even understanding this question for curves in $\mathbb{P}^1 \times \mathbb{P}^2$ would be compelling. In light of Theorem 1.6, this case would produce unirationality results for certain parameter spaces of curves.

Question 7.2. For which values of d, e, and g, does there exist a smooth curve in $\mathbb{P}^1 \times \mathbb{P}^2$ of bidegree (d,e) and genus g with a virtual resolution of the form $S \longleftarrow F_1 \longleftarrow F_2 \longleftarrow 0$?

Proposition 6.1 and Theorem 1.6 suggest that having a virtual resolution whose length equals the codimension of the underlying variety can have significant geometric implications. As these results parallel the arithmetic Cohen–Macaulay property over projective space, it would be interesting to seek out analogues of being arithmetically Gorenstein.

Question 7.3. Consider a positive-dimensional subscheme $Z \subseteq \mathbb{P}^n$ such that $\omega_Z = \mathscr{O}_{\mathbb{P}^n}(d)|_Z$ for some $d \in \mathbb{Z}^r$. Is there a self-dual virtual resolution of Z?

It would also be interesting to better understand scheme-theoretic complete intersections.

Question 7.4. Develop an algorithm to determine if a subvariety $Z \subseteq \mathbb{P}^n$ has a virtual resolution that is a Koszul complex. This is already interesting in the case of points on $\mathbb{P}^1 \times \mathbb{P}^1$.

Finally, we believe that many of these results should hold for more general toric varieties.

Question 7.5. Prove an analogue of Proposition 1.2 for an arbitrary smooth toric variety.

REFERENCES

- [Art76] Michael Artin, On deformations of singularities, Tata Institute of Fundamental Research, Bombay, 1976.
- [Aud91] Michèle Audin, *The topology of torus actions on symplectic manifolds*, Progress in Mathematics 93, Birkhäuser Verlag, Basel, 1991.
- [BS98] Dave Bayer and Bernd Sturmfels, *Cellular resolutions of monomial modules*, J. Reine Angew. Math. **502** (1998), 123–140.
- [BC17] Nicolás Botbol and Marc Chardin, *Castelnuovo Mumford regularity with respect to multigraded ideals*, J. Algebra **474** (2017), 361–392.
- [BE77] David A. Buchsbaum and David Eisenbud, *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension* 3, Amer. J. Math. **99** (1977), no. 3, 447–485.

- [Cox95] David A. Cox, *The homogeneous coordinate ring of a toric variety*, J. Algebraic Geom. **4** (1995), no. 1, 17–50.
- [Căl05] Andrei Căldăraru, *Derived categories of sheaves: a skimming*, Snowbird lectures in algebraic geometry, Contemp. Math. 388, Amer. Math. Soc., Providence, RI, 2005, pp. 43–75.
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics 124, Amer. Math. Soc., Providence, RI, 2011.
- [DS00] Wolfram Decker and Frank-Olaf Schreyer, *Non-general type surfaces in* **P**⁴: *some remarks on bounds and constructions*, J. Symbolic Comput. **29** (2000), no. 4-5, 545–582.
- [DFS16] Anand Deopurkar, Maksym Fedorchuk, and David Swinarski, *Toward GIT stability of syzygies of canonical curves*, Algebr. Geom. **3** (2016), no. 1, 1–22.
- [EGA3I] Alexander Grothendieck and Jean A. Dieudonné, Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents, Première partie, Inst. Hautes Études Sci. Publ. Math. 11 (1961), 5–167.
- [EGA3II] ______, Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents, Seconde partie, Inst. Hautes Études Sci. Publ. Math. 17 (1963), 5–91.
 - [EL15] Lawrence Ein and Robert Lazarsfeld, *The gonality conjecture on syzygies of algebraic curves of large degree*, Publ. Math. Inst. Hautes Études Sci. **122** (2015), 301–313.
 - [EL93] ______, Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension, Invent. Math. 111 (1993), no. 1, 51–67.
 - [Eis95] David Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995.
- [EES15] David Eisenbud, Daniel Erman, and Frank-Olaf Schreyer, *Tate resolutions for products of projective spaces*, Acta Math. Vietnam. **40** (2015), no. 1, 5–36.
- [EP99] David Eisenbud and Sorin Popescu, *Gale duality and free resolutions of ideals of points*, Invent. Math. **136** (1999), no. 2, 419–449.
- [Far09] Gavril Farkas, Koszul divisors on moduli spaces of curves, Amer. J. Math. 131 (2009), no. 3, 819-867.
- [Fuj83] Takao Fujita, *Vanishing theorems for semipositive line bundles*, Algebraic geometry (Tokyo/Kyoto, 1982), Lecture Notes in Math. 1016, Springer, Berlin, 1983, pp. 519–528.
- [GR03] Ofer Gabber and Lorenzo Ramero, *Almost ring theory*, Lecture Notes in Math. 1800, Springer-Verlag, Berlin Heidelberg, 2003.
- [GP99] Francisco J. Gallego and Bangere P. Purnaprajna, *Projective normality and syzygies of algebraic surfaces*, J. Reine Angew. Math. **506** (1999), 145–180.
- [GGP95] Anthony V. Geramita, Alessandro Gimigliano, and Yves Pitteloud, *Graded Betti numbers of some embedded rational n-folds*, Math. Ann. **301** (1995), no. 2, 363–380.
- [GL86] Mark Green and Robert Lazarsfeld, *On the projective normality of complete linear series on an algebraic curve*, Invent. Math. **83** (1986), no. 1, 73–90.
- [GVT15] Elena Guardo and Adam Van Tuyl, *Arithmetically Cohen-Macaulay sets of points in* $\mathbb{P}^1 \times \mathbb{P}^1$, SpringerBriefs in Mathematics, Springer, Cham, 2015.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics 52, Springer-Verlag, New York-Heidelberg, 1977.
- [Har10] ______, *Deformation theory*, Graduate Texts in Mathematics 257, Springer, New York, 2010.
- [HH93] Melvin Hochster and Craig Huneke, *Phantom homology*, Mem. Amer. Math. Soc. **103** (1993), no. 490, vi+91.
- [Huy06] Daniel Huybrechts, *Fourier-Mukai transforms in algebraic geometry*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, Oxford, 2006.
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Series of Modern Surveys in Mathematics 48, Springer-Verlag, Berlin, 2004.
- [MS04] Diane Maclagan and Gregory G. Smith, *Multigraded Castelnuovo-Mumford regularity*, J. Reine Angew. Math. **571** (2004), 179–212.

- [M2] Daniel R. Grayson and Michael E. Stillman, *Macaulay2*, a software system for research in algebraic geometry, available at http://www.math.uiuc.edu/Macaulay2/.
- [MR92] Rosa M. Miró-Roig, *Nonobstructedness of Gorenstein subschemes of codimension* 3 in \mathbf{P}^n , Beiträge Algebra Geom. **33** (1992), 131–138.
- [PS85] Ragni Piene and Michael Schlessinger, *On the Hilbert scheme compactification of the space of twisted cubics*, Amer. J. Math. **107** (1985), no. 4, 761–774.
- [Rob87] Paul Roberts, Le théorème d'intersection, C. R. Acad. Sci. Paris Sér. I Math. 304 (1987), no. 7, 177–180.
- [Sch77] Mary Schaps, *Deformations of Cohen–Macaulay schemes of codimension 2 and non-singular deformations of space curves*, Amer. J. Math. **99** (1977), no. 4, 669–685.
- [Voi02] Claire Voisin, *Green's generic syzygy conjecture for curves of even genus lying on a K3 surface*, J. Eur. Math. Soc. (JEMS) **4** (2002), no. 4, 363–404.

CHRISTINE BERKESCH ZAMAERE: SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA, 55455, UNITED STATES OF AMERICA; cberkesc@math.umn.edu

DANIEL ERMAN: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN, 53706, UNITED STATES OF AMERICA; derman@math.wisc.edu

GREGORY G. SMITH: DEPARTMENT OF MATHEMATICS & STATISTICS, QUEEN'S UNIVERSITY, KINGSTON, ONTARIO, K7L 3N6, CANADA; ggsmith@mast.queensu.ca