

Gauge fields for ultracold atoms in optical superlattices

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2010 New J. Phys. 12 033007

(<http://iopscience.iop.org/1367-2630/12/3/033007>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 202.78.175.199

This content was downloaded on 25/09/2015 at 20:50

Please note that [terms and conditions apply](#).

Gauge fields for ultracold atoms in optical superlattices

Fabrice Gerbier¹ and Jean Dalibard¹

Laboratoire Kastler Brossel, ENS, UPMC-Paris 6, CNRS, 24 rue Lhomond,
75005 Paris, France

E-mail: fabrice.gerbier@lkb.ens.fr and jean.dalibard@lkb.ens.fr

New Journal of Physics **12** (2010) 033007 (16pp)

Received 21 October 2009

Published 9 March 2010

Online at <http://www.njp.org/>

doi:10.1088/1367-2630/12/3/033007

Abstract. We present a scheme that produces a strong U(1)-like gauge field on cold atoms confined in a two-dimensional square optical lattice. Our proposal relies on two essential features, a long-lived metastable excited state that exists for alkaline-earth or ytterbium atoms and an optical superlattice. As in the proposal by Jaksch and Zoller (2003 *New J. Phys.* **5** 56), laser-assisted tunneling between adjacent sites creates an effective magnetic field. In the tight-binding approximation, atomic motion is described by the Harper Hamiltonian, with a flux across each lattice plaquette that can realistically take any value between 0 and π . We show how one can take advantage of the superlattice to ensure that each plaquette acquires the same phase, thus simulating a uniform magnetic field. We discuss the observable consequences of the artificial gauge field on non-interacting bosonic and fermionic gases. We also outline how the scheme can be generalized to non-Abelian gauge fields.

¹ Authors to whom any correspondence should be addressed.

Contents

1. Introduction	2
2. Two-electron atoms in state-dependent optical lattices	4
3. State-dependent lattice: Harper Hamiltonian for $\alpha = 1/2$	6
4. State-dependent superlattice: Harper Hamiltonian for arbitrary α	7
4.1. Flux rectification	7
4.2. Optical superlattice	8
4.3. Conditions of validity	9
4.4. Comparison with the JZ proposal	9
5. Role of an external trap potential	10
5.1. Density of states in the presence of an external trap	10
5.2. Spatial distribution for non-interacting fermions	12
5.3. Momentum distribution for non-interacting bosons	12
6. Conclusion and perspectives	13
Acknowledgments	13
References	14

1. Introduction

The fractional quantum Hall phases realized by two-dimensional (2D) electron gases in very large magnetic fields are among the most intriguing states of matter (see, for instance, [1]). In such systems, electrons ‘bind’ to magnetic vortices, forming strongly correlated phases with striking properties, such as exotic excitations (‘anyons’), which obey fractional statistics [2]. Analogous quantum Hall phases should also arise in cold atomic gases when they are set into fast rotation (see [3] and references therein). Owing to the mathematical similarity between Coriolis and Lorentz forces, rotating neutral gases are indeed the exact analogue of an assembly of charged particles plunged in a magnetic field. Observing these highly correlated phases is one of the major goals in the field of trapped quantum gases [3, 4]. This goal has, however, not yet been reached, because of the difficulty of communicating the required amount of angular momentum (on the order of $N\hbar$ per atom, with N the number of particles) to the system [5, 6]. In practice, residual static trap anisotropy limits the total angular momentum to much smaller values, for which the rotating gas is well described by a mean field approach [7].

Recently, alternative schemes of ‘simulating’ artificial gauge fields for neutral atoms have been explored using 2D optical lattices [8]–[14]. As they do not involve any mechanical rotation of the system, they should be less sensitive to the imperfection of the trapping potential and thus easier to implement. The guideline for these proposals is the celebrated Harper model [15, 16], defined by the 2D single-particle Hamiltonian,

$$H_{\text{Harper}} = -J \sum_{n,m,\pm} e^{\pm i2\pi\alpha m} \hat{c}_{n\pm 1,m}^\dagger \hat{c}_{n,m} + \hat{c}_{n,m\pm 1}^\dagger \hat{c}_{n,m} + \text{h.c.} \quad (1)$$

This tight-binding model was initially introduced [15] to describe electrons hopping on a square lattice perpendicular to a constant magnetic field B . The operator $\hat{c}_{n,m}^\dagger$ creates a particle at position $(x = nd, y = md)$, d is the lattice spacing and J is the tunnel energy to nearest

neighbors in the absence of a magnetic field. The effect of a vector potential (here in the Landau gauge, $\mathbf{A} = -By\mathbf{e}_x$) is included in the so-called Peierls phase factor, $2\pi\alpha m$, where $\alpha = eBd^2/h$ is the flux per unit cell expressed in units of the flux quantum $\phi_0 = h/e$. The Harper Hamiltonian in equation (1) has been studied extensively in the literature (see e.g. [15]–[18]). For zero flux, the Harper model reduces to a standard tight-binding model on a square lattice, with a single Bloch band of width $8J$. For rational fluxes $\alpha = p/q$ (where p and q are integers), the model is still periodic but with larger unit cells of size $qd \times d$. The Bloch band splits into q magnetic sub-bands, and the ground state becomes q -fold degenerate. This results in the peculiar self-similar structure in an energy–magnetic flux diagram known as Hofstadter’s butterfly [16]. This structure exists in a strict sense for infinitely extended systems. For a finite system of size L , the details of this structure are washed out on fine scales such that $q > L/d$.

In their seminal proposition for an implementation of (1) [8], Jaksch and Zoller (JZ) suggested that the phase $2\pi\alpha$ can be imprinted by a laser beam inducing hopping between adjacent sites (see also [9]–[14] for alternative proposals). A major advantage of this approach is that large flux $\alpha \sim 1$ can be reached under experimentally realistic conditions. For a filling factor with ~ 1 atom per site, the quantum Hall regime then becomes reachable in the presence of moderate atomic interactions [11], [19]–[22]. Theoretical studies have found that some of these phases are essentially similar to those encountered in the bulk [11, 19, 20, 22], whereas others are without counterparts in the bulk phase [23, 24].

Although the existing proposals for implementing the Harper Hamiltonian with atomic quantum gases open many avenues, they are still challenging from an experimental point of view (for reasons discussed in detail below). In this paper, we extend the JZ proposal and discuss a new scheme based on an optical superlattice [25, 26] to generate a gauge potential leading to (1). At variance with earlier works where alkali atoms were considered, here we propose using atoms with a long-lived metastable excited state, such as alkaline-earth [27]–[29] or ytterbium [30, 31] atoms. For concreteness we will discuss the case of ytterbium, for which degenerate gases have already been produced for both bosonic and fermionic isotopes [30, 31]. We take advantage of this level structure to alleviate many practical difficulties encountered with alkalis. Importantly, our scheme uses only building blocks that have already been individually demonstrated.

This paper is organized as follows. We first introduce the system, consisting of atoms with a ground state g and a long-lived metastable excited state e trapped in a 2D lattice. The optical lattice is tuned near an ‘anti-magic’ wavelength where the polarizabilities of e and g are opposite [32]. The two internal states thus live in spatially separated sublattices. We show that the case $\alpha = 1/2$, leading to Dirac points analogous to those appearing in graphene [33], can be achieved rather simply with a single laser beam coupling e and g . We then extend our scheme to generate arbitrary values of α using an optical superlattice, at the expense of several lasers operating at different frequencies. Subsequently, we address the influence of the additional potential that must be added to confine the particles. We investigate its effect by looking at (i) the density of states that determines the density profile of a trapped ideal Fermi gas and (ii) the ground state of the system, relevant for non-interacting bosons. In both cases, we obtain a clear signature for a nonzero value of α . We conclude by indicating how the method can be generalized to generate non-Abelian gauge fields [34].




Figure 1. (a) Sketch of the level scheme showing the two internal states g and e coherently coupled by a laser beam. (b) Dynamic polarizability $\alpha(\lambda)$ of atoms in the ground state g (solid line) or excited state e (dashed line). The light shift experienced by the atoms is $\alpha(\lambda)|\mathbf{E}|^2/2$, with $|\mathbf{E}|$ being the electric field strength at wavelength λ . Two particular wavelengths are of special interest, a ‘magic’ wavelength $\lambda_m \approx 760$ nm and an ‘anti-magic’ wavelength $\lambda_{am} \approx 1.12 \mu\text{m}$. (c) Sketch of the atomic configuration in a state-dependent optical lattice, showing the two interlaced sublattices for g (gray dots) and e (black dots) atoms. The y lattice is formed by a standing wave at the ‘magic’ wavelength and confines both states identically. The x lattice is tuned near the ‘anti-magic’ wavelength and confines the two states in two distinct sublattices shifted by $\lambda_{am}/4$.

2. Two-electron atoms in state-dependent optical lattices

Two-electron atoms, such as ytterbium and alkaline-earth atoms, generally have a spin-singlet ($^1S_0 \equiv g$) ground state and an extremely long-lived spin triplet ($^3P_0 \equiv e$) excited state (see figure 1(a)). The very long lifetime of e (around 20 s for Yb [35]) allows one to operate optical atomic clocks at the resonance wavelength $\lambda_0 \approx 578$ nm between g and e [36]–[38]. It has also inspired novel proposals for quantum information processing [39]–[41] or quantum simulation [42, 43]. States g and e usually react differently to far-off resonance laser light, as they couple optically to many different states. The polarizabilities of the two states calculated using the data from [44] are shown in figure 1(b). For laser light tuned at the so-called *magic wavelength* λ_m (~ 760 nm for Yb), both states have the same polarizability and feel the same optical trapping potential [45]. Central to our proposal is the existence of an ‘anti-magic’ wavelength $\lambda_{am} \approx 1.12 \mu\text{m}$ [32], for which the polarizabilities of g and e are opposite^{2,3}. As we will see, this allows one to create easily internal-state-dependent potentials.

² There exists another anti-magic wavelength near 620 nm, which is unfortunately rather close to a transition connecting to the 3P_0 state (649 nm). A lattice at this wavelength is thus likely to suffer from excessive spontaneous emission.

³ A many-electron calculation [46] using more accurate parameters for the positions and widths of the optical transitions predicts the anti-magic wavelengths near 1120.3 nm and 618.7 nm (Derevianko A 2009 Private communication).

The first building block of our proposal consists in a laser beam at the magic wavelength, forming a horizontal light sheet that freezes the vertical motion of the atoms irrespective of their internal state e or g . From now on, we shall focus on the atom dynamics in the horizontal xy plane. The second ingredient consists in an optical standing wave along the y -axis, also at the magic wavelength, which generates the same potential $V(y) = -V_0 \cos^2(\pi y/d_y)$ for both states g and e ($V_0 > 0$). For the third building block, we apply another optical standing wave at the ‘anti-magic’ wavelength λ_{am} along the x -axis, generating the potential $V'(x) = \pm V_1 \cos^2(\pi x/2d_x)$, where the $+$ and $-$ signs refer to e and g , respectively. The resulting potential $V'(x) + V(y)$ creates two shifted sublattices (see figure 1(c)). Atoms in state g sit at positions $\mathbf{r}_g = (2n, m)$ and atoms in state e at positions $\mathbf{r}_e = (2n+1, m)$, where n, m are integers. Here we take as length scales $d_x = \lambda_{\text{am}}/4$ along x , and $d_y = \lambda_m/2$ along y . We write the Wannier functions for the lowest Bloch band as $w_g(\mathbf{r} - \mathbf{r}_g)$ and $w_e(\mathbf{r} - \mathbf{r}_e)$. The potential V_0 is chosen such that the tunneling energy $J^{(y)}$ along the vertical lines is significant. On the contrary, we choose V_1 large enough to neglect direct tunneling along the x -axis.

Tunneling along the x -axis is induced by a coherent optical coupling between g and e . Since atoms in e and g are trapped in separate optical lattices, this coupling induces hopping from one sublattice to the other. Following Jaksch and Zoller [8], we calculate the effective $g-e$ hopping matrix element between two neighboring sites located at $\mathbf{r}_g = (2n, m)$ and $\mathbf{r}_e = \mathbf{r}_g + \mathbf{b} = (2n+1, m)$ as

$$J_{eg}^{(x)} e^{i\mathbf{q} \cdot \mathbf{r}_g} = \frac{\hbar \Omega}{2} e^{i\mathbf{q} \cdot \mathbf{r}_g} \int w_e^*(\mathbf{r} - \mathbf{b}) e^{i\mathbf{q} \cdot \mathbf{r}} w_g(\mathbf{r}) d^2\mathbf{r}, \quad (2)$$

with Ω and \mathbf{q} the laser Rabi frequency and wavevector. We assume that the coupling laser propagates in the yz plane, so that $\mathbf{q} \cdot \mathbf{r}_g = 2\pi\alpha m$, where $|\alpha|$ can take any value between 0 and $\lambda_m/2\lambda_0$ ($\simeq 0.66$ for Yb) by adjusting the angle between the coupling laser beam and the z -axis.

Before proceeding with further analysis, let us give some numerical estimates for the different physical quantities involved in this experimental configuration. The anti-magic wavelength λ_{am} corresponds, for Yb, to a recoil energy $E_R/h = \hbar/(2m\lambda_{\text{am}}^2) \approx 900$ Hz. Realistic values up to several tens of kHz can be achieved for the depth of the optical lattice potential V_1/h . In order to avoid excitations to higher-lying Bloch bands, the Rabi frequency Ω must be small compared to Δ/\hbar , where Δ is the energy gap between the ground and first excited bands. To give numerical estimates, we take $V_1 = 20E_R$, corresponding to a gap $\Delta \approx 8E_R$. Assuming a moderate Rabi frequency $\Omega = E_R/\hbar$, we find a laser-induced tunneling energy $J_{eg}^{(x)} \approx 0.05E_R$, much larger than the residual tunneling energy in each sublattice without change in internal state, $J_{gg}^{(x)} = J_{ee}^{(x)} \approx 2.5 \times 10^{-3}E_R$. In principle, besides resonant transitions between the fundamental bands of the two sublattices, the coupling laser also induces second-order tunneling processes, where an atom tunnels to a neighboring site in a given sublattice, via a virtual transition to an intermediate excited band in the other sublattice. We have verified that the corresponding tunneling rates were negligible compared to $J_{gg}^{(x)}$ for the range of parameters investigated here. Finally, one can calculate the spontaneous emission and associated heating rates for atoms in each sublattice [47]. For lattice depths $V_0 = V_1 \sim 30E_R$, we find the total heating rates $\gamma_h^{(g)} \sim 0.4 \text{ nK s}^{-1}$ and $\gamma_h^{(e)} \sim 0.9 \text{ nK s}^{-1}$, respectively, or $\gamma_h^{(g)}/E_R \sim 0.01 \text{ s}^{-1}$ and $\gamma_h^{(e)}/E_R \sim 0.02 \text{ s}^{-1}$. The heating rates (dominated by the vertical arm at λ_m) are small enough to maintain atoms in the ground band for several seconds.




Figure 2. Laser schemes for realizing staggered (a, b) and uniform (c, d) magnetic fields. (a) The figure shows the different on-site energies along x , in the case of a simple-period lattice. All transition frequencies are degenerate, so that a single laser beam at frequency ω_L is sufficient to couple all sites across the lattice. (b) Staggered effective magnetic field for a single coupling laser propagating along y with degenerate transition frequencies as shown in (a). (c) In a configuration where an additional, doubly periodic shift of the on-site energies is introduced, the degeneracy between the transition frequencies is lifted. (d) With several dedicated laser beams (ω_+ propagating along y , ω_- and ω'_- propagating along $-y$), one obtains a ‘rectified’ uniform magnetic field.

3. State-dependent lattice: Harper Hamiltonian for $\alpha = 1/2$

We first consider the simplest experimental arrangement, with a spin-dependent lattice as described above and a single coupling laser (figure 2(a)). The two-component $g - e$ quantum gas, which is assumed to populate the lowest Bloch band only, is described by the Hamiltonian

$$\begin{aligned}
 H = & -J \sum_{n,m,n \text{ even}} \left(e^{i2\pi\alpha m} \hat{c}_{2n+1,m}^\dagger \hat{c}_{2n,m} + e^{i2\pi\alpha m} \hat{c}_{2n-1,m}^\dagger \hat{c}_{2n,m} + \text{h.c.} \right) \\
 & -J \sum_{n,m,\pm} \left(\hat{c}_{n,m\pm 1}^\dagger \hat{c}_{n,m} + \text{h.c.} \right),
 \end{aligned} \tag{3}$$

when the lattice amplitudes are adjusted so that $J^{(y)} = J_{eg}^{(x)} = J$. As in [8], the operator $\hat{c}_{n,m}^\dagger$ creates an atom at site (n, m) , the parity of n identifying the internal state unambiguously. Alternatively, the internal degrees of freedom can be eliminated by going to a dressed state basis, and the spatial wavefunctions associated with each dressed state obey the *same* Harper equation [15, 16] as the one derived from (1). Hence, at least in the absence of interactions

(which we assume throughout this paper), the internal state is merely a label that can be ignored in the analysis.

In the original equation (1), the phase picked up across each link has the same sign for a given link direction in *real* space: $e^{+i2\pi\alpha m}$ in the $+x$ -direction and $e^{-i2\pi\alpha m}$ in the $-x$ -direction. Here, we find in contrast that the phase is tied to the direction in *internal* space: $e^{+i2\pi\alpha m}$ for $g \rightarrow e$ transitions and $e^{-i2\pi\alpha m}$ for $e \rightarrow g$ transitions. In real space, given the sublattice geometry, this corresponds to a phase that alternates in sign from an arbitrary unit cell (or ‘plaquette’) to the neighboring one along the x -axis. More precisely, consider a particle going clockwise around an elementary cell (figure 1(c)). The phase picked up along the path $(2n, m-1)_g \rightarrow (2n, m)_g \rightarrow (2n+1, m)_e \rightarrow (2n+1, m-1)_e \rightarrow (2n, m-1)_g$ is the product of the two factors $e^{i2\pi\alpha m} e^{-i2\pi\alpha(m-1)} = e^{i2\pi\alpha}$. Thanks to the phase factor $\propto e^{i\mathbf{q} \cdot \mathbf{r}_g}$, a particle making a loop around a plaquette acquires a nonzero global phase factor, reflecting the nonzero ‘magnetic flux’ through the cell. However, for the adjacent plaquette, the path (also with clockwise orientation) $(2n+1, m-1)_e \rightarrow (2n+1, m)_e \rightarrow (2n+2, m)_g \rightarrow (2n+2, m-1)_g \rightarrow (2n+1, m-1)_e$ leads to a phase factor $e^{-i2\pi\alpha}$ opposite to the previous one. One then achieves in this way a *staggered* magnetic field, with a phase per plaquette $\pm 2\pi\alpha$ constant on a given column and changing sign from one column to the next (figure 2(b)).

The case $\alpha = \frac{1}{2}$ is peculiar, since in this case a staggered or a uniform flux both correspond to a tunneling phase $e^{\pm i2\pi\alpha m} = (-1)^m$ alternating between adjacent links depending on the parity of m . The Hamiltonian remains periodic, with a unit cell $d_x \times 2d_y$ and two sites per unit cell. The energy spectrum is given by

$$E(k_x, k_y) = \pm 2J \sqrt{\cos^2(k_x d_x) + \cos^2(k_y d_y)}, \quad (4)$$

where the Bloch vector (k_x, k_y) sits in the first Brillouin zone $]-\pi/d_x, \pi/d_x] \times]-\pi/2d_y, \pi/2d_y]$. We note that this dispersion relation gives rise to two ‘Dirac points’ for $(k_x, k_y) = (\pm\pi/2d_x, \pi/2d_y)$ around which the dispersion relation is linear [13, 48] (see also [49]–[53] for discussions of the Dirac points occurring in different settings, such as hexagonal lattices or non-Abelian gauge fields). The physical consequences of the existence of these Dirac points are similar to those found in the case of graphene [33]. However, their topological robustness [54] might be affected by small deviations from $\alpha = 1/2$ that break the two-site periodicity.

For $\alpha \neq \frac{1}{2}$, the staggered value of the phase per plaquette leads to a Hamiltonian that is different from (1), and that, in fact, does not exhibit any structure reminiscent of Hofstadter’s butterfly [55]. This can be linked to a general argument due to Haldane [56], emphasizing that breaking time-reversal invariance symmetry is the key factor to generating systems with properties similar to (integer) quantum Hall phases. The situation considered up to this point, where the same laser beam drives all $g \rightarrow e$ transitions, is invariant with respect to time reversal. We describe in the next section a means to break this invariance and achieve a flux with a constant sign on all plaquettes.

4. State-dependent superlattice: Harper Hamiltonian for arbitrary α

4.1. Flux rectification

To ensure an identical phase $2\pi\alpha$ per plaquette, instead of the staggered values $\pm 2\pi\alpha$, we consider a situation where the on-site energies are modulated spatially along x with a period

equal to twice the lattice spacing, as depicted in figure 2(c) (the y lattice remains identical for both states and is ignored in the following). Assume that the on-site energies are modulated according to

$$\begin{aligned} E_g - V_1 & \text{ for } x/d_x = 4n, \\ E_e + V_2 & \text{ for } x/d_x = 4n + 1, \\ E_g - V_1 - V_2 & \text{ for } x/d_x = 4n + 2, \\ E_e & \text{ for } x/d_x = 4n + 3. \end{aligned}$$

Here the E_α ($\alpha = e, g$) denote the internal energies in free space, where V_1 is the amplitude of the fundamental lattice with period d_x and where $V_2 \ll V_1$ is the amplitude of the modulation with period $2d_x$. In this potential landscape, the resonance frequencies for transitions linking neighboring sites become non-degenerate (figure 2(c)),

$$\begin{aligned} \omega_+ &= \omega_0 + (V_1 + V_2)/\hbar \quad \text{for } |g; 4n, m\rangle \rightarrow |e; 4n + 1, m\rangle, \\ \omega_- &= \omega_0 + (V_1 + 2V_2)/\hbar \quad \text{for } |e; 4n + 1, m\rangle \rightarrow |g; 4n + 2, m\rangle, \\ \omega_+ &= \omega_0 + (V_1 + V_2)/\hbar \quad \text{for } |g; 4n + 2, m\rangle \rightarrow |e; 4n + 3, m\rangle, \\ \omega'_- &= \omega_0 + V_1/\hbar \quad \text{for } |e; 4n + 3, m\rangle \rightarrow |g; 4n + 4, m\rangle, \end{aligned}$$

with $\hbar\omega_0 = E_e - E_g$.

To the state-dependent lattice, one thus applies three coupling lasers propagating along y with frequencies ω_+ , ω_- and ω'_- . The laser at frequency ω_+ is chosen with a wavevector \mathbf{k}_L , and the lasers at frequencies ω_- and ω'_- with the opposite wavevector $-\mathbf{k}_L$. The alternance of the wavevectors compensates the alternance of the plaquette phases, thereby ‘rectifying’ the staggered flux obtained in a single-frequency configuration. If we neglect off-resonant transitions, an analysis similar to that made in the previous section confirms that each plaquette now acquires the same phase factor $e^{i2\pi\alpha}$, and that the system is indeed described by the Harper Hamiltonian (1).

4.2. Optical superlattice

To realize such a modulation in practice, we propose superimposing along the x -direction an additional lattice potential with a double spatial period $2\lambda_{\text{am}}$, which can be written as $V'(x) = V_{g/e} \cos^2(\pi x/4d_x + \varphi)$, where φ is the relative phase between the fundamental and double period lattices⁴. Such superlattice potentials have been demonstrated experimentally for quantum gases [25, 26]. This potential can be realized in practice by adding a second laser with frequency $c/2\lambda_{\text{am}}$ phase locked to the fundamental lattice to control the relative phase φ [26]. The amplitudes $V_{g/e}$ depend on the internal state and are fixed by the initial choice of wavelength for the fundamental lattice. The analysis in section 4.1 (where $V_g = V_e$) can be generalized straightforwardly to this less symmetric potential, with four non-degenerate transition frequencies in the general case. Moreover, it is possible to reduce the number of required laser frequencies to three by using a particular value of the relative phase φ that reproduces exactly the scheme described previously, with $V_2 = V_e \cos \varphi - V_g \sin \varphi$.

⁴ Practically, the fluctuations $\delta\varphi$ of φ must be small enough to ensure that fluctuations of V_2 remain much smaller than the width of the $g - e$ resonance, set by the Rabi frequency Ω , or $\delta\varphi \ll \hbar\Omega/V_2 \lesssim 0.1$. This is well within present technological capabilities.

Table 1. Characteristic tunnel energies (along the x -axis) achievable for realistic experimental parameters in the superlattice configuration. Here $J_{eg}^{(x)}$ denotes the laser-assisted coupling from one sublattice to the other. We used a Rabi frequency $\hbar\Omega = V_2/10 = \Delta/30$ (see text).

Depth V_1/E_R	Energy gap Δ/E_R	Laser-assisted tunneling $J_{eg}^{(x)}/E_R$
10	4.7	16×10^{-3}
20	7.8	13×10^{-3}
30	9.8	9×10^{-3}

4.3. Conditions of validity

We now discuss the domain of validity of our model.

- Firstly, as in the previous sections, we demand that laser-assisted tunneling dominates over regular tunneling without a change in internal state. For a superlattice potential in the tight-binding regime, regular tunneling is suppressed when $V_2 \geq E_R$, since $J_{gg}^{(x)}, J_{ee}^{(x)} \ll E_R$.
- Secondly, the ratio $\hbar\Omega/V_2$, with Ω the Rabi frequency characterizing atom–laser coupling, must remain small to neglect off-resonant transitions and describe the system with an effective Harper model [8]. In practice, one can use a conservative value $\hbar\Omega = V_2/10$.
- Finally, the modulation V_2 must be small enough to avoid excitations to higher-lying Bloch bands by one of the coupling lasers. A reasonable choice is $V_2 = \Delta/3$, where Δ is the energy gap between the ground and first excited bands for $V_2 = 0$. This guarantees that transitions mixing different bands are detuned by at least $\Delta/3$. For example, if we focus on the $|e; 4n+3, m\rangle \rightarrow |g; 4n+4, m\rangle$ transition, transitions linking the ground and first excited bands occur at frequencies $\omega'_\pm \pm \Delta/\hbar$. One can check that the detunings for the different coupling lasers are $|V_2 \pm \Delta|$, $|2V_2 \pm \Delta|$, Δ , which are indeed $\geq \Delta/3$ for the choice $V_2 = \Delta/3$.

We note that in the tight-binding regime, Δ is notably larger than E_R , which allows one to fulfill simultaneously all these conditions while maintaining a substantial value for $J_{eg}^{(x)}$. We summarize our considerations in table 1, where we give typical values for lattice parameters and laser-assisted tunneling matrix elements in the superlattice configuration. We note that the values for $J_{eg}^{(x)}$ remain sizeable, being on the same order as the tunnel energies $J_{gg}^{(x)}$ in the g sublattice ($\hbar\Omega = V_2 = 0$) at moderate depths ($\sim 10E_R$).

4.4. Comparison with the JZ proposal

The scheme as described in this paper is inspired by the earlier proposal by Jaksch and Zoller [8]. That proposal was designed for alkali atoms in spin-dependent optical lattices. In this context, g and e denoted two different hyperfine states in the ground state manifold, and the spin-dependent optical potential was obtained by exploiting the vector light shift arising in a laser field with suitable polarization (see [57, 58], for example).

The first difference between our method and the JZ proposal comes from the choice of two-electron atoms versus ‘one-electron’ alkalis. To obtain a useful spin-dependent lattice for

alkalis demands, in practice, a lattice laser tuned between the D_1 and D_2 lines, relatively close to resonance. For light alkalis, such as ^6Li or ^{40}K , the fine splitting between the D_1 and D_2 transitions is rather small (less than 1nm), and spontaneous emission rates are too large to maintain the cloud in the nK regime. The most practical candidate seems to be ^{87}Rb , which has only stable bosonic isotopes and would probably still suffer from spurious spontaneous emission due to the coupling (Raman) lasers or the lattice lasers. The choice of Yb removes all these drawbacks, by using only far-off resonant lasers and a dissipation-free coupling transition, and by allowing the study of bosonic and fermionic systems in the same setup.

The use of a superlattice to ensure the same flux for all plaquettes constitutes the second main difference of our proposal with respect to JZ's proposal. There, a strong 'tilting' potential independent of the internal state was introduced in order to distinguish $|g; 2n, m\rangle \rightarrow |e; 2n \pm 1, m\rangle$. Although this could be done, in principle, using a large static electric field or the light shift from an intense far-off resonant laser beam, it turns out to be rather challenging from an experimental point of view⁵. In addition, this tilting potential has to be almost perfectly linear over the extent of the cloud. Otherwise, the various transition frequencies would depend on space, spoiling the flux uniformity and the scalability of the method. We believe that the use of a superlattice optical potential will be a more flexible and more reliable tool that can be readily implemented experimentally.

5. Role of an external trap potential

We now discuss the properties of a system of non-interacting particles described by equation (1) in order to identify observable consequences of the laser-induced gauge potential. For a uniform system without trapping potential, and for $\alpha = 0$, the model reduces to a standard tight-binding Hamiltonian. The spectrum consists in an allowed energy band of width $8J$ centered at zero energy. For the uniform Harper model, without trapping potential but with $\alpha = p/q \neq 0$ (p and q are integers), the initial Bloch band splits into q magnetic sub-bands separated by well-defined energy gaps. This results in q distinct peaks in the density of state (DoS—see figures 3(a) and (b)). In a practical experiment, an additional trapping potential is inevitably present in addition to the optical lattice. Hence the physical system is both finite and inhomogeneous. This can strongly affect the properties of the uniform Harper model given in equation (1), but also provide novel experimental signatures, as we discuss in this section.

5.1. Density of states in the presence of an external trap

We consider here the Harper model in equation (1) with an additional isotropic trapping potential $V_{\text{trap}} = \kappa(r/d)^2$. Let us recall that, with $\alpha = 0$, one finds a quasi-continuum of allowed states in the combined lattice plus external trap, in striking contrast with the uniform case [59]. States with energy lower than $4J$ are qualitatively similar to the Bloch eigenstates of the uniform problem. These states are approximately contained in a disk of radius $R_{\text{eff}} \sim \sqrt{J/\kappa}$. States with higher energy (which would lie in the bandgap for the uniform system) are well-localized states

⁵ The Stark shift from the electric field is proportional to $\alpha|E|^2$, with α the static polarizability. Using a configuration where the cloud is offset from the center of the electrodes creating the field, one can create a linear potential of the form $\eta x/d_x$, with $\eta \sim \alpha U^2 d_x / D^3$ (D being the distance between the electrodes and U the applied electrostatic potential difference). For ^{87}Rb and $D \sim 1$ cm, we find that a site-to-site energy offset on the order of the energy gap Δ , $\eta \sim \Delta/3 \sim 10$ kHz requires $U \sim 50$ kV, which is rather challenging to produce.




Figure 3. (a) Energy spectrum and (b) density of states (DoS) for the uniform model with $\alpha = 1/3$. (c) DoS for the trapped model with $\alpha = 1/3$ and confinement strength $\kappa/J = 0.01$. The dashed line shows the expected DoS for $E \gg J$, irrespective of α . (d) Density profile for non-interacting fermions in a harmonic trap with $\kappa/J = 0.01$, for a flux $\alpha = 1/3$ (solid lines) and Fermi energies $E_F = 0$ (half-filled band) and $4J$ (filled band). The density profiles for $\alpha = 0$ (dashed lines) are also shown for comparison. The logarithmic divergence around $E = 0$ is smeared out by the finite size of the lattice used in our calculation (63×63 space grid).

offset from the cloud center [60], with an energy mostly determined by the local potential energy near the turning point. In two dimensions, the density of states saturates at an asymptotic value of π/κ .

We now address the case with $\alpha = p/q \neq 0$. The finiteness of the central region where tunneling is significant limits the relevant values of q to $q < q_{\max} = R_{\text{eff}}/d$. We consider here the case $\alpha = 1/3$ for concreteness. Solving the full model numerically, we find that the trapping potential has a dominant effect on the DoS and little remains of the initial peak structure (cf figures 3(b) and (c)). This indicates that global observables (such as thermodynamic quantities) will not be very sensitive to the presence of the gauge potential. We can understand this behavior from a semiclassical argument. Indeed, in a local density approximation, the density of states in a generic power-law potential $\kappa(r/d)^\gamma$ can be approximated as

$$\rho_{\text{trap}}(E) = \sum_n \int \frac{d^{(2)}\mathbf{r} d^{(2)}\mathbf{k}}{(2\pi)^2} \delta\left(E - \epsilon_n(\mathbf{k}) - \kappa \left(\frac{r}{d}\right)^\gamma\right), \quad (5)$$

where $\epsilon_n(\mathbf{k}) = 4J \chi_n(\mathbf{k})$ is the relation dispersion for the n th magnetic sub-band of the untrapped system. This can be simplified to

$$\rho_{\text{trap}}(E) = \frac{2\pi(4J)^{(2/\gamma)-1}}{\gamma \kappa^{2/\gamma}} \sum_n \int_{(E/4J) > \chi_n(\mathbf{k})} \frac{d^{(2)}\mathbf{k}}{S_{\text{BZ}}} \left(\frac{E}{4J} - \chi_n(\mathbf{k})\right)^{(2/\gamma)-1}. \quad (6)$$

Here $S_{\text{BZ}} = (2\pi/d)^2$ is the surface of the Brillouin zone. For a harmonic potential ($\gamma = 2$), the dependence on the reduced dispersion function χ is mild since it only arises from the upper bound of the integral: the peculiarities of the spectrum are essentially washed out by the spatial integration. This can be improved by using a steeper trap with $\gamma > 2$, e.g. a quartic potential as demonstrated in [6].

5.2. Spatial distribution for non-interacting fermions

In the case of non-interacting fermions in a gauge potential, a more sensitive observable is given by the particle density [22], assuming it can be measured *in situ*. Indeed, still within a semiclassical approximation, the spatial density can be written as

$$n(\mathbf{r}) = \sum_n \int \frac{d^{(2)}\mathbf{k}}{(2\pi)^2} f_{\text{FD}} [\epsilon_n(\mathbf{k}) + V_{\text{trap}}(r) - E_{\text{F}}],$$

where f_{FD} is the Fermi–Dirac distribution at $T = 0$ and E_{F} is the Fermi energy. We rewrite this as

$$\begin{aligned} n(\mathbf{r}) &= \sum_n \int \frac{d^{(2)}\mathbf{k}}{(2\pi)^2} \int_{-4J}^{\infty} dE f_{\text{FD}} [V_{\text{trap}} + E - E_{\text{F}}] \delta [\epsilon_n(\mathbf{k}) - E] \\ &= \int_{-4J}^{\infty} dE' f_{\text{FD}} [V_{\text{trap}} + E' - E_{\text{F}}] \rho_{\text{hom}}[E'], \end{aligned} \quad (7)$$

where

$$\rho_{\text{hom}}[E] = \sum_n \int \frac{d^{(2)}\mathbf{k}}{(2\pi)^2} \delta [\epsilon_n(\mathbf{k}) - E] \quad (8)$$

is the density of state for the *homogeneous* system. For low enough temperatures, the clustering of energy levels in sub-bands in the presence of a gauge potential will result in well-defined plateaux in the spatial density of the trapped cloud. These plateaux correspond to ‘band-insulating’ regions, similar to those appearing for other incompressible systems in traps such as Mott insulators [61, 62]. We have verified this fact in our numerical calculations, as shown in figure 3(d). We conclude that the in-trap density profile will provide a clear signature of the gauge potential in the case of fermionic atoms.

5.3. Momentum distribution for non-interacting bosons

With non-interacting bosons, such a signature will be absent, as the atoms will condense into the lowest single-particle states (for the purpose of this paper, we do not consider interactions). However, signatures of the gauge potential will appear in reciprocal momentum space, which can be probed using standard time-of-flight imaging. The dephasing term $e^{i2\pi\alpha m}$ generates quasi-momentum components at the various harmonics of the fundamental period $2\pi\alpha$. As a result, clear satellite peaks can be observed in the momentum distribution (see figure 4(a)), which will provide a direct experimental signature for bosonic species. We note, in addition, that the density distributions corresponding to these eigenstates show periodic structures with period q (see figure 4(b)), although this signature appears on a much finer scale than in the fermionic case (thus being more difficult to observe).




Figure 4. Typical momentum (a) and spatial (b) distributions in the ground state for non-interacting bosons and a flux $\alpha = 1/3$. The presence of a gauge potential reveals itself in momentum components at $\pm 2\pi/3d$ in the former plot, and in the fine structures with period $3d$ in the latter. The confinement strength is $\kappa/J = 0.005$.

6. Conclusion and perspectives

In this article, we have discussed how one can implement Abelian artificial gauge potentials for Yb (or alkaline-earth) atoms, a scheme valid for bosons and fermions alike. In contrast to bosonic isotopes with zero total spin, fermionic isotopes have nuclear spin. This potentially enables one to simulate non-Abelian gauge fields, along the lines proposed by Osterloh *et al* [34]. Let us focus for concreteness on the ^{171}Yb isotope, with nuclear spin $I = \frac{1}{2}$. In the presence of a moderate magnetic field of a few tens of gauss, the degeneracy of the $m_I = \pm \frac{1}{2}$ states within the e and g manifolds is lifted, giving in general four different internal transitions, $\pm \frac{1}{2} \rightarrow \pm \frac{1}{2}$ and $\pm \frac{1}{2} \rightarrow \mp \frac{1}{2}$, with different frequencies. At the expense of using more laser frequencies and state-dependent lattices along both the x - and y -directions, one can then arrange for the lasers to imprint different phases depending on the internal state or generate arbitrary rotations in internal space. As shown in [34], a straightforward generalization of the Abelian scheme then allows one to engineer *non-Abelian* gauge potentials. Instead of a simple phase, hopping from sublattice g to e then corresponds to a rotation in internal space generated by non-commuting matrices. Combined with interatomic interactions, an entirely new class of superfluid or strongly correlated systems becomes accessible with ultracold atoms [34, 53], [63]–[65]. In a wider context, this technique can be used to emulate spin–orbit coupling, similar to the Rashba interaction $\propto \hat{\sigma}_x \cdot k_y - \hat{\sigma}_y \cdot k_x$ in planar semiconductors, or for the study of strongly interacting, ultracold fermionic matter in a non-Abelian background field, possibly relevant to high-energy physics.

Acknowledgments

We thank the members of the ‘Bose–Einstein condensates’ and ‘Fermi gases’ groups at LKB, G Montambaux and G Juzeliunas for helpful discussions, and A Derevianko for useful correspondence. This work was supported by ANR (Grant ANR-08-BLAN-65BOFL), IFRAF, the European Union (MIDAS project) and DARPA (OLE project). Laboratoire Kastler Brossel is unité mixte de recherche (UMR) number 8552 of the CNRS.

References

- [1] Girvin S M 1999 The quantum Hall effect: novel excitations and broken symmetries *Topological Aspects of Low Dimensional Systems* ed A Comtet, T Jolicoeur, S Ouvry and F David (Berlin: Springer) (arXiv:cond-mat/9907002)
- [2] Stern A 2008 Anyons and the quantum Hall effect—a pedagogical review *Ann. Phys.* **323** 204–49
- [3] Cooper N R 2008 Rapidly rotating atomic gases *Adv. Phys.* **57** 539
- [4] Bloch I, Dalibard J and Zwerger W 2008 Many-body physics with ultracold gases *Rev. Mod. Phys.* **80** 885
- [5] Schweikhard V, Coddington I, Engels P, Mogendorff V P and Cornell E A 2004 Rapidly rotating Bose–Einstein condensates in and near the lowest Landau level *Phys. Rev. Lett.* **92** 040404
- [6] Bretin V, Stock S, Seurin Y and Dalibard J 2004 Fast rotation of a Bose–Einstein condensate *Phys. Rev. Lett.* **92** 050403
- [7] Ho T-L 2001 Bose–Einstein condensates with large number of vortices *Phys. Rev. Lett.* **87** 060403
- [8] Jaksch D and Zoller P 2003 Creation of effective magnetic fields in optical lattices: the Hofstadter butterfly for cold neutral atoms *New J. Phys.* **5** 56
- [9] Ruostekoski J, Dunne G V and Javanainen J 2002 Particle number fractionalization of an atomic Fermi–Dirac gas in an optical lattice *Phys. Rev. Lett.* **88** 180401
- [10] Mueller E J 2004 Artificial electromagnetism for neutral atoms: Escher staircase and Laughlin liquids *Phys. Rev. A* **70** 041603
- [11] Sorensen A S, Demler E and Lukin M D 2005 Fractional quantum Hall states of atoms in optical lattices *Phys. Rev. Lett.* **94** 086803
- [12] Dudarev A M, Diener R B, Carusotto I and Niu Q 2004 Spin–orbit coupling and Berry phase with ultracold atoms in 2d optical lattices *Phys. Rev. Lett.* **92** 153005
- [13] Lim L-K, Morais C and Smith Hemmerich A 2008 Staggered-vortex superfluid of ultracold bosons in an optical lattice *Phys. Rev. Lett.* **100** 130402
- [14] Klein A and Jaksch D 2009 Phonon-induced artificial magnetic fields in optical lattices *Europhys. Lett.* **85** 13001
- [15] Harper P G 1955 Single band motion of conduction electrons in a uniform magnetic field *Proc. Phys. Soc. A* **68** 874–8
- [16] Hofstadter D R 1976 Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields *Phys. Rev. B* **14** 2239–49
- [17] Thouless D J, Kohmoto M, Nightingale M P and den Nijs M 1982 Quantized Hall conductance in a two-dimensional periodic potential *Phys. Rev. Lett.* **49** 405–8
- [18] Kohmoto M 1989 Zero modes and the quantized Hall conductance of the two-dimensional lattice in a magnetic field *Phys. Rev. B* **39** 11943–9
- [19] Palmer R N and Jaksch D 2006 High-field fractional quantum Hall effect in optical lattices *Phys. Rev. Lett.* **96** 180407
- [20] Palmer R N, Klein A and Jaksch D 2008 Optical lattice quantum Hall effect *Phys. Rev. A* **78** 013609
- [21] Hafezi M, Sorensen A S, Demler E and Lukin M D 2007 Fractional quantum Hall effect in optical lattices *Phys. Rev. A* **76** 023613
- [22] Umucalilar R O, Zhai H and Oktel M O 2008 Trapped Fermi gases in rotating optical lattices: realization and detection of the topological Hofstadter insulator *Phys. Rev. Lett.* **100** 070402
- [23] Hafezi M, Sorensen A S, Lukin M D and Demler E 2008 Characterization of topological states on a lattice with Chern number *Europhys. Lett.* **81** 10005
- [24] Möller G and Cooper N R 2009 Composite fermion theory for bosonic quantum Hall states on lattices *Phys. Rev. Lett.* **103** 105303
- [25] Sebby-Strabley J, Anderlini M, Jessen P S and Porto J V 2006 Lattice of double wells for manipulating pairs of cold atoms *Phys. Rev. A* **73** 033605

- [26] Fölling S, Trotzky S, Cheinet P, Feld M, Saers R, Widera A, Müller T and Bloch I 2007 Direct observation of second-order atom tunnelling *Nature* **448** 1029–32
- [27] Kraft S, Vogt F, Appel O, Riehle F and Sterr U 2009 Bose–Einstein condensation of alkaline earth atoms: ^{40}Ca *Phys. Rev. Lett.* **103** 130401
- [28] Stellmer S, Khoon M Tey, Huang B, Grimm R and Schreck F 2009 Bose–Einstein condensation of strontium *Phys. Rev. Lett.* **103** 200401
- [29] Martínez de Escobar Y N, Mickelson P G, Yan M, DeSalvo B J, Nagel S B and Killian T C 2009 Bose–Einstein condensation of ^{84}Sr *Phys. Rev. Lett.* **103** 200402
- [30] Takasu Y, Maki K, Komori K, Takano T, Honda K, Kumakura M, Yabuzaki T and Takahashi Y 2003 Spin-singlet Bose–Einstein condensation of two-electron atoms *Phys. Rev. Lett.* **91** 040404
- [31] Fukuhara T, Takasu Y, Kumakura M and Takahashi Y 2007 Degenerate Fermi gases of ytterbium *Phys. Rev. Lett.* **98** 030401
- [32] Yi W, Daley A J, Pupillo G and Zoller P 2008 State-dependent, addressable subwavelength lattices with cold atoms *New J. Phys.* **10** 073015
- [33] Castro A H, Neto Guinea F, Peres N M R, Novoselov K S and Geim A K 2009 The electronic properties of graphene *Rev. Mod. Phys.* **81** 109
- [34] Osterloh K, Baig M, Santos L, Zoller P and Lewenstein M 2005 Cold atoms in non-Abelian gauge potentials: from the Hofstadter ‘moth’ to lattice gauge theory *Phys. Rev. Lett.* **95** 010403
- [35] Porsev S G, Derevianko A and Fortson E N 2004 Possibility of an optical clock using the $6^1\text{S}_0 \rightarrow 6^3\text{P}_0$ transition in $^{171,173}\text{Yb}$ atoms held in an optical lattice *Phys. Rev. A* **69** 021403
- [36] Hong T, Cramer C, Cook E, Nagourney W and Fortson E N 2005 Observation of the $^1\text{S}_0\text{--}^3\text{P}_0$ transition in atomic ytterbium for optical clocks and qubit arrays *Opt. Lett.* **30** 2644–6
- [37] Hoyt C W, Barber Z W, Oates C W, Fortier T M, Diddams S A and Hollberg L 2005 Observation and absolute frequency measurements of the $\text{S}_1\text{--P}_3$ optical clock transition in neutral ytterbium *Phys. Rev. Lett.* **95** 083003
- [38] Barber Z W, Hoyt C W, Oates C W, Hollberg L, Taichenachev A V and Yudin V I 2006 Direct excitation of the forbidden clock transition in neutral ^{174}Yb atoms confined to an optical lattice *Phys. Rev. Lett.* **96** 083002
- [39] Reichenbach I and Deutsch I H 2007 Sideband cooling while preserving coherences in the nuclear spin state in group-II-like atoms *Phys. Rev. Lett.* **99** 123001
- [40] Stock R, Babcock N S, Raizen M G and Sanders B C 2008 Entanglement of group-II-like atoms with fast measurement for quantum information processing *Phys. Rev. A* **78** 022301
- [41] Gorshkov A V, Rey A M, Daley A J, Boyd M M, Ye J, Zoller P and Lukin M D 2009 Alkaline-earth-metal atoms as few-qubit quantum registers *Phys. Rev. Lett.* **102** 110503
- [42] Sato M, Takahashi Y and Fujimoto S 2009 Non-Abelian topological order in s-wave superfluids of ultracold fermionic atoms *Phys. Rev. Lett.* **103** 020401
- [43] Gorshkov A V, Hermele M, Gurarie V, Xu C, Julienne P S, Ye J, Zoller P, Demler E, Lukin M D and Rey A M 2009 Two-orbital $\text{SU}(N)$ magnetism with ultracold alkaline-earth atoms arXiv:0905.2610
- [44] Barber Z 2007 Ytterbium optical lattice clock *PhD Thesis* University of Colorado
- [45] Ye J, Kimble H J and Katori H 2008 Quantum state engineering and precision metrology using state-insensitive light traps *Science* **320** 1734–8
- [46] Dzuba V A and Derevianko A 2009 Dynamic polarizabilities and related properties of clock states of ytterbium atom arXiv:0908.2278v1
- [47] Gordon P J and Ashkin A 1980 Motion of atoms in a radiation trap *Phys. Rev. A* **21** 1606–17
- [48] Hou J-M, Yang W-X and Liu X-J 2009 Massless Dirac fermions in a square optical lattice *Phys. Rev. A* **79** 043621
- [49] Zhao E and Paramekanti A 2006 BCS–BEC crossover on the two-dimensional honeycomb lattice *Phys. Rev. Lett.* **97** 230404

- [50] Zhu S-L, Wang B and Duan L-M 2007 Simulation and detection of Dirac fermions with cold atoms in an optical lattice *Phys. Rev. Lett.* **98** 260402
- [51] Wu C and Das Sarma S 2008 $p^{x,y}$ -orbital counterpart of graphene: cold atoms in the honeycomb optical lattice *Phys. Rev. B* **77** 235107
- [52] Loon Lee K, Grémaud B, Han R, Englert B-G and Miniatura C 2009 Ultracold fermions in a graphene-type optical lattice *Phys. Rev. A* **80** 043411
- [53] Goldman N, Kubasiak A, Bermudez A, Gaspard P, Lewenstein M and Martin-Delgado M A 2009 Non-Abelian optical lattices: anomalous quantum Hall effect and Dirac fermions *Phys. Rev. Lett.* **103** 035301
- [54] Montambaux G, Piéchon F, Fuchs J-N and Goerbig M O 2009 Merging of Dirac points in a two-dimensional crystal *Phys. Rev. B* **80** 153412
- [55] Wang Y-F and Gong C-D 2006 Hofstadter butterflies in macroscopic time-reversal invariant systems: tight-binding electrons under $n\sqrt{2} \times n\sqrt{2}$ staggered magnetic fields *Phys. Rev. B* **74** 193301
- [56] Haldane F D M 1988 Model for a quantum Hall effect without Landau levels: condensed-matter realization of the ‘parity anomaly’ *Phys. Rev. Lett.* **61** 2015–8
- [57] Cohen-Tannoudji C and Dupont-Roc J 1972 Experimental study of Zeeman light shifts in weak magnetic fields *Phys. Rev. A* **5** 968–84
- [58] Mandel O, Greiner M, Widera A, Rom T, Hänsch T W and Bloch I 2003 Coherent transport of neutral atoms in spin-dependent optical lattice potentials *Phys. Rev. Lett.* **91** 010407
- [59] Hooley C and Quintanilla J 2004 Single-atom density of states of an optical lattice *Phys. Rev. Lett.* **93** 080404
- [60] Ott H, de Mirandes E, Ferlaino F, Roati G, Türeci V, Modugno G and Inguscio M 2004 Radio frequency selective addressing of localized atoms in a periodic potential *Phys. Rev. Lett.* **93** 120407
- [61] Folling S, Widera A, Müller T, Gerbier F and Bloch I 2006 Formation of spatial shell structure in the superfluid to Mott insulator transition *Phys. Rev. Lett.* **97** 060403
- [62] Campbell G K, Mun J, Boyd M, Medley P, Leanhardt A E, Marcassa L G, Pritchard D E and Ketterle W 2006 Imaging the Mott insulator shells by using atomic clock shifts *Science* **313** 649–52
- [63] Goldman N 2007 Spatial patterns in optical lattices submitted to gauge potentials *Europhys. Lett.* **80** 20001
- [64] Satija I I, Dakin D C, Vaishnav J Y and Clark C W 2008 Physics of a two-dimensional electron gas with cold atoms in non-Abelian gauge potentials *Phys. Rev. A* **77** 043410
- [65] Goldman N, Kubasiak A, Gaspard P and Lewenstein M 2009 Ultracold atomic gases in non-Abelian gauge potentials: the case of constant Wilson loop *Phys. Rev. A* **79** 023624