

The even Orlicz Minkowski problem

Christoph Haberl, Erwin Lutwak^{*,1}, Deane Yang¹, Gaoyong Zhang¹

Polytechnic Institute of NYU, Six MetroTech Center, Brooklyn, NY 11201, United States

Received 14 September 2009; accepted 8 February 2010

Available online 26 February 2010

Communicated by the Managing Editors of AIM

Abstract

The classical Minkowski problem leads to the L_p Minkowski problem and now to the Orlicz Minkowski problem. Existence is demonstrated for the even Orlicz Minkowski problem. A byproduct is a new approach to the solution of the classical Minkowski problem.

© 2010 Elsevier Inc. All rights reserved.

Keywords: Aleksandrov body; Minkowski problem; Orlicz norm; Petty projection inequality; Support function; Surface area measure

1. Introduction

The Minkowski problem concerns the existence, uniqueness, and stability of convex hypersurfaces whose Gauss curvature (possibly in a generalized sense) is prescribed as a function of the outer unit normals. The Minkowski problem is one of the centerpieces of the classical Brunn–Minkowski theory. The complete solution to the Minkowski problem (for arbitrary “data” — with no smoothness assumptions) goes back exactly three-quarters of a century to the work of Aleksandrov and Fenchel and Jessen (see Schneider [59] for references). Of course, when the Minkowski “data” is discrete then the solution goes back to Minkowski’s work at the turn of the 19th into the 20th Century.

Almost a century after Minkowski’s original work, an L_p version of the Brunn–Minkowski theory began to emerge, beginning largely with [33,34], and expanding rapidly thereafter (see,

* Corresponding author.

E-mail address: lutwak@poly.edu (E. Lutwak).

¹ Research supported, in part, by NSF Grant DMS-0706859.

e.g., [1–5, 7–11, 13, 15–22, 25–40, 42–45, 49, 50, 52–55, 58, 60–63, 65–68]). The L_p Minkowski problem is also of central importance in this new L_p -Brunn–Minkowski theory.

For a given continuous function $g : S^{n-1} \rightarrow \mathbb{R}$ (called the “data”), the *regular L_p Minkowski problem* seeks solutions $h : S^{n-1} \rightarrow (0, \infty)$ to the partial differential equation

$$h^{1-p} \det(h_{ij} + h\delta_{ij}) = g \quad \text{on } S^{n-1}, \quad (1)$$

where (h_{ij}) is the Hessian matrix of h with respect to an orthonormal frame on S^{n-1} . The case $p = 1$ is classical with landmark contributions such as Cheng and Yau [6] and Pogorelov [56].

The *general L_p Minkowski problem* asks: Given a real p , what are the necessary and sufficient conditions on a Borel measure μ on the unit sphere, S^{n-1} , such that there exists a convex body K in \mathbb{R}^n with support function h_K and surface area measure S_K (see Section 2 for definitions) so that

$$h_K^{1-p} dS_K = d\mu. \quad (2)$$

The solutions to the general L_p Minkowski problem (2) may be viewed as weak solutions to the regular L_p Minkowski problem (1). While regularity is a central focus in the fields of partial differential equations and differential geometry, the existence and the uniqueness of weak solutions of the Minkowski problem are of paramount interest to those working in convex geometric and functional analysis. This is because it is precisely these solutions that correspond to support functions of polytopes or to Banach norms (which often have limited smoothness).

The *even L_p Minkowski problem* seeks solutions under the assumption that the data measure μ is an even Borel measure on S^{n-1} (i.e. assumes the same values on antipodal Borel sets) or the data function g is an even function (i.e. assumes the same values on antipodal points of S^{n-1}). The L_p Minkowski problem with even data is not only natural but of practical interest because its solutions correspond to norms of Banach spaces.

In this paper the “ L_p Minkowski problem” will always refer to the general L_p Minkowski problem (2). The case $p = 1$ of the L_p Minkowski problem (2) is of course the classical Minkowski problem. For $p > 1$, a solution to the even L_p Minkowski problem was given in [33] under the assumption that $p \neq n$. In [41], it was shown that, for $p \neq n$, the L_p Minkowski problem (2) has an equivalent volume-normalized formulation and a solution of the even volume-normalized L_p Minkowski problem was given for all $p > 1$. The regular even L_p Minkowski problem was studied in [35].

In the plane ($n = 2$), the L_p Minkowski problem was treated by Stancu [61–63], Umanskiy [65], Chen [5], and most recently by Jiang [24].

The L_p Minkowski problem (without the assumption that the data is even) was treated by Guan and Lin [15] and later by Chou and Wang [7]. Hug et al. [23] gave an alternate approach to some of the results of Chou and Wang [7].

The solution of the even L_p Minkowski problem was a critical ingredient that allowed the authors of [40] to extend the affine Sobolev inequality [69] and obtain the L_p affine Sobolev inequality and later enabled Cianchi et al. [8] to establish the affine Moser–Trudinger and the affine Morrey–Sobolev inequalities.

Recent work of Haberl and Schuster [19,20], Haberl, Schuster, and Xiao [21], Ludwig and Reitzner [32], and Ludwig [31], shows the need to take the next step in the evolution of the Brunn–Minkowski theory towards an Orlicz Brunn–Minkowski theory. This is the third paper in

a series [46,47] that attempts to develop some of the elements of an Orlicz Brunn–Minkowski theory.

Suppose $\varphi : (0, \infty) \rightarrow (0, \infty)$ is a fixed continuous function. The aim of this paper is to study the even L_φ Minkowski problem: If μ is an even finite Borel measure on S^{n-1} which is not concentrated on a great subsphere of S^{n-1} , then does there exist an origin symmetric convex body K in \mathbb{R}^n such that

$$c\varphi(h_K) dS_K = d\mu,$$

for some positive number c ? We will show that under some mild assumptions on φ , the even L_φ Minkowski problem does have a solution. For example we shall establish:

Theorem. *Suppose $\varphi : (0, \infty) \rightarrow (0, \infty)$ is a continuous decreasing function. If μ is an even finite Borel measure on S^{n-1} which is not concentrated on a great subsphere of S^{n-1} , then there exists an origin symmetric convex body K in \mathbb{R}^n such that*

$$c\varphi(h_K) dS_K = d\mu,$$

where c is a power of the volume of K — specifically, $c = V(K)^{\frac{1}{2n}-1}$.

When φ and the data μ are assumed to be sufficiently smooth, and under certain other restrictions, the existence of solutions to the L_φ Minkowski problem was already demonstrated by Chou and Wang [7]. For applications in analysis (such as in obtaining analytic inequalities of the type presented in [69] and [40]) and for applications in geometry (such as in obtaining affine isoperimetric inequalities), various Banach norms and convex bodies must be constructed. These constructions amount to solving an L_φ Minkowski problem — but usually with minimal restrictions on φ and the measure μ . (Note that a polytopal solution of the L_φ Minkowski problem corresponds to a measure μ whose support is a finite set.) In this paper we will establish the existence of solutions to the L_φ Minkowski problem with such minimal restrictions on φ and μ .

One interesting feature of our work is that it presents a new approach to the classical Minkowski problem as well as the even L_p Minkowski problem for $p > 0$.

2. Preliminaries

For quick later reference we develop some notation and basic facts about convex bodies. Good general references for the theory of convex bodies are provided by the books of Gardner [12], Gruber [14], Schneider [59], and Thompson [64].

Our setting will be Euclidean n -space \mathbb{R}^n where $n \geq 2$. The standard inner product of the vectors $x, y \in \mathbb{R}^n$ is denoted by $x \cdot y$. We write $S^{n-1} = \{x \in \mathbb{R}^n : x \cdot x = 1\}$ for the boundary of the Euclidean unit ball B in \mathbb{R}^n .

The set of continuous functions on the sphere S^{n-1} will be denoted by $C(S^{n-1})$ and will always be viewed as equipped with the max-norm metric:

$$\|f - g\|_\infty = \max_{u \in S^{n-1}} |f(u) - g(u)|,$$

for $f, g \in C(S^{n-1})$. The subspace of positive continuous functions will be denoted by $C^+(S^{n-1})$ and $C_e^+(S^{n-1})$ will denote the subspace of $C^+(S^{n-1})$ consisting of only the even functions.

Write V for n -dimensional Lebesgue measure and \mathcal{H}^{n-1} for $(n-1)$ -dimensional Hausdorff measure. The letter μ will be used exclusively to denote a finite Borel measure on S^{n-1} . For such a measure μ , we denote by $|\mu|$ its total mass, i.e. $|\mu| = \mu(S^{n-1})$. The letter c (possibly with subscripts or other distinguishing features) will be used exclusively to denote a positive real number.

A *convex body* is a compact convex subset of \mathbb{R}^n with non-empty interior. The set of convex bodies in \mathbb{R}^n containing the origin in their interiors is denoted by \mathcal{K}_o^n . The set of convex bodies in \mathbb{R}^n that are symmetric about the origin will be denoted by \mathcal{K}_e^n .

A compact, convex set $K \subset \mathbb{R}^n$ is uniquely determined by its *support function* $h_K : \mathbb{R}^n \rightarrow \mathbb{R}$, where $h_K(x) = \max\{x \cdot y : y \in K\}$, for each $x \in \mathbb{R}^n$. For example, the support function of the line segment \bar{v} joining the points $\pm v \in \mathbb{R}^n$ is given by

$$h_{\bar{v}}(x) = |x \cdot v|, \quad x \in \mathbb{R}^n.$$

We will need the trivial fact that for the support function of the dilate $cK = \{cx : x \in K\}$ of a compact, convex K we have

$$h_{cK} = ch_K, \quad c > 0. \quad (3)$$

Note that support functions are positively homogeneous of degree 1 and subadditive. It follows immediately from the definition of support functions that for compact, convex $K, L \subset \mathbb{R}^n$,

$$K \subseteq L \iff h_K \leq h_L. \quad (4)$$

Consequently, the support function of a body $K \in \mathcal{K}_o^n$ is bounded from above and below by positive reals.

The set \mathcal{K}_o^n will be viewed as equipped with the Hausdorff metric and thus for a sequence $\{K_i\}$ of bodies in \mathcal{K}_o^n and a body $K \in \mathcal{K}_o^n$, we have $\lim_{i \rightarrow \infty} K_i = K$ provided that

$$\|h_{K_i} - h_K\|_\infty \rightarrow 0.$$

A boundary point $x \in \partial K$ is said to have $u \in S^{n-1}$ as an outer normal provided $x \cdot u = h_K(u)$. A boundary point is said to be *singular* if it has more than one unit normal vector. It is well known (see, e.g., [59]) that the set of singular boundary points of a convex body has \mathcal{H}^{n-1} -measure equal to 0.

For a convex body K and each Borel set $\omega \subset S^{n-1}$, the *inverse spherical image*, $\tau(K, \omega)$, of K at ω is the set of all boundary points of K which have an outer unit normal belonging to the set ω . Associated with each convex body $K \in \mathcal{K}_o^n$ is a Borel measure, S_K , on S^{n-1} called the Aleksandrov–Fenchel–Jessen *surface area measure* of K , defined by

$$S_K(\omega) = \mathcal{H}^{n-1}(\tau(K, \omega)), \quad (5)$$

for each Borel set $\omega \subseteq S^{n-1}$. Observe that for the surface area measure of the dilate cK of K we have

$$S_{cK} = c^{n-1} S_K, \quad c > 0. \quad (6)$$

We will use the weak continuity of surface area measures; i.e., if $\{K_i\}$ is a sequence of bodies in \mathcal{K}_o^n then

$$\lim_{i \rightarrow \infty} K_i = K \in \mathcal{K}_o^n \implies \lim_{i \rightarrow \infty} S_{K_i} = S_K, \quad \text{weakly.} \quad (7)$$

The *mixed volume* $V_1(K, L)$ of the convex bodies $K, L \in \mathcal{K}_o^n$ may be defined by

$$V_1(K, L) = \frac{1}{n} \int_{S^{n-1}} h_L dS_K. \quad (8)$$

The fact that

$$V(K) = V_1(K, K),$$

or equivalently,

$$V(K) = \frac{1}{n} \int_{S^{n-1}} h_K dS_K, \quad (9)$$

is of critical importance. The fundamental inequality for mixed volumes is *Minkowski's mixed volume inequality*: For $K, L \in \mathcal{K}_o^n$,

$$V_1(K, L)^n \geq V(K)^{n-1} V(L) \quad (10)$$

with equality if and only if K and L are homothetic.

3. Aleksandrov bodies

A function $h \in C^+(S^{n-1})$ defines a family $\{H_u\}_{u \in S^{n-1}}$ of hyperplanes

$$H_u = \{x \in \mathbb{R}^n : x \cdot u = h(u)\}.$$

This family gives rise to concepts such as envelopes in classical differential geometry, generalized envelopes in convex geometric analysis (see, e.g., [51]), and hedgehogs (see, e.g., [48]).

We shall be interested in the intersection of the halfspaces that are associated to h by the family $\{H_u\}_{u \in S^{n-1}}$. This gives rise to the convex body

$$K = \bigcap_{u \in S^{n-1}} \{x \in \mathbb{R}^n : x \cdot u \leq h(u)\}.$$

The body K is called the *Aleksandrov body associated with h* . Note that since h is both positive and continuous its Aleksandrov body, K , must be an element of \mathcal{K}_o^n . The Aleksandrov body associated with h can alternatively be defined as the unique maximal element, with respect to set inclusion, of the set

$$\{Q \in \mathcal{K}_o^n : h_Q \leq h\}.$$

For the Aleksandrov body K associated with h we now examine equality (in S^{n-1}) in the inequality

$$h_K \leq h$$

in some detail. Aleksandrov showed that each element of $\tau(K, \omega_h)$, the inverse spherical image of K of the set

$$\omega_h = \{u \in S^{n-1} : h_K(u) < h(u)\},$$

must be a singular boundary point of K . Since the set of singular boundary points of a convex body has \mathcal{H}^{n-1} -measure zero, and since by (5), we know that $S_K(\omega_h) = \mathcal{H}^{n-1}(\tau(K, \omega_h))$, it follows that $S_K(\omega_h) = 0$. Consequently, while $h_K \leq h$, in fact

$$h_K = h, \quad \text{a.e. with respect to } S_K. \quad (11)$$

We will make use of the following three basic properties of Aleksandrov bodies. First, if h is the support function of a convex body $K \in \mathcal{K}_o^n$, then K itself is the Aleksandrov body associated with h . Second, as is easily shown, if h is an even function, then the Aleksandrov body associated with h is origin symmetric. Third and critical, is *Aleksandrov's convergence lemma* (see, e.g., [59, Lemma 6.5.2]): If the functions $h_i \in C^+(S^{n-1})$ have associated Aleksandrov bodies $K_i \in \mathcal{K}_o^n$, then

$$h_i \rightarrow h \in C^+(S^{n-1}) \implies K_i \rightarrow K,$$

where K is the Aleksandrov body associated with h .

The volume $V(h)$ of a function $h \in C^+(S^{n-1})$ is defined as the volume of the Aleksandrov body associated with h . Since the Aleksandrov body associated with the support function h_K of a convex body $K \in \mathcal{K}_o^n$ is the body K itself, we have

$$V(h_K) = V(K). \quad (12)$$

From Aleksandrov's convergence lemma and the continuity of ordinary volume on \mathcal{K}_o^n we see that

$$V : C^+(S^{n-1}) \rightarrow \mathbb{R} \quad \text{is continuous.}$$

Let $I \subset \mathbb{R}$ be an interval containing 0 and suppose that $h_t(u) = h(t, u) : I \times S^{n-1} \rightarrow (0, \infty)$ is continuous. For fixed $t \in I$, let

$$K_t = \bigcap_{u \in S^{n-1}} \{x \in \mathbb{R}^n : x \cdot u \leq h(t, u)\}$$

be the Aleksandrov body associated with h_t . The family of bodies $\{K_t\}_{t \in I}$ will be called *the family of Aleksandrov bodies associated with h* . Obviously, from (11) we have

$$h_{K_t} \leq h_t \quad \text{and} \quad h_{K_t} = h_t, \quad \text{a.e. with respect to } S_{K_t}, \quad (13)$$

for each $t \in I$.

Lemma 1. Let $I \subset \mathbb{R}$ be an interval containing 0 and some positive number and the function $h(t, u) : I \times S^{n-1} \rightarrow (0, \infty)$ be continuous and such that the convergence in

$$h'_+(0, u) = \lim_{t \rightarrow 0^+} \frac{h(t, u) - h(0, u)}{t} \quad (14)$$

is uniform on S^{n-1} . If $\{K_t\}_{t \in I}$ is the family of Aleksandrov bodies associated with h , then

$$\lim_{t \rightarrow 0^+} \frac{V(K_t) - V(K_0)}{t} = \int_{S^{n-1}} h'_+(0, u) dS_{K_0}(u).$$

Proof. The uniform convergence of (14) implies that $h_t \rightarrow h_0$, uniformly on S^{n-1} . Therefore, the Aleksandrov convergence lemma (mentioned above) yields

$$\lim_{t \rightarrow 0^+} K_t = K_0. \quad (15)$$

By (7) we conclude that the S_{K_t} converge weakly to S_{K_0} as $t \rightarrow 0^+$. Since the measures S_{K_t} are finite, converge weakly to S_{K_0} and since the convergence in

$$\lim_{t \rightarrow 0^+} \frac{h(t, u) - h(0, u)}{t}$$

is uniform on S^{n-1} , we obtain

$$\lim_{t \rightarrow 0^+} \int_{S^{n-1}} \frac{h_t(u) - h_0(u)}{t} dS_{K_t}(u) = \int_{S^{n-1}} h'_+(u, 0) dS_{K_0}(u). \quad (16)$$

Formulas (9) and (13) imply

$$V(K_t) = \frac{1}{n} \int_{S^{n-1}} h_{K_t}(u) dS_{K_t}(u) = \frac{1}{n} \int_{S^{n-1}} h_t(u) dS_{K_t}(u). \quad (17)$$

From (17), (8), and inequality (13) at $t = 0$, we have

$$\begin{aligned} \liminf_{t \rightarrow 0^+} \frac{V(K_t) - V_1(K_t, K_0)}{t} &= \frac{1}{n} \liminf_{t \rightarrow 0^+} \int_{S^{n-1}} \frac{h_t(u) - h_{K_0}(u)}{t} dS_{K_t}(u) \\ &\geq \frac{1}{n} \liminf_{t \rightarrow 0^+} \int_{S^{n-1}} \frac{h_t(u) - h_0(u)}{t} dS_{K_t}(u), \end{aligned}$$

which when combined with (16) gives

$$\liminf_{t \rightarrow 0^+} \frac{V(K_t) - V_1(K_t, K_0)}{t} \geq \frac{1}{n} \int_{S^{n-1}} h'_+(u, 0) dS_{K_0}(u). \quad (18)$$

For the sake of brevity set

$$l = \frac{1}{n} \int_{S^{n-1}} h'_+(u, 0) dS_{K_0}(u).$$

Inequality (18) and Minkowski's mixed volume inequality (10) show

$$l \leq \liminf_{t \rightarrow 0^+} \frac{V(K_t) - V_1(K_t, K_0)}{t} \leq \liminf_{t \rightarrow 0^+} \frac{V(K_t) - V(K_t)^{1-\frac{1}{n}} V(K_0)^{\frac{1}{n}}}{t}.$$

But (15) gives $\lim_{t \rightarrow 0^+} V(K_t) = V(K_0)$ and hence

$$l \leq V(K_0)^{1-\frac{1}{n}} \liminf_{t \rightarrow 0^+} \frac{V(K_t)^{\frac{1}{n}} - V(K_0)^{\frac{1}{n}}}{t}. \quad (19)$$

Now (8), the inequality in (13), and the uniform convergence in (14) give

$$\begin{aligned} \limsup_{t \rightarrow 0^+} \frac{V_1(K_0, K_t) - V(K_0)}{t} &= \frac{1}{n} \limsup_{t \rightarrow 0^+} \int_{S^{n-1}} \frac{h_{K_t}(u) - h_0(u)}{t} dS_{K_0}(u) \\ &\leq \frac{1}{n} \limsup_{t \rightarrow 0^+} \int_{S^{n-1}} \frac{h_t(u) - h_0(u)}{t} dS_{K_0}(u) \\ &= \frac{1}{n} \int_{S^{n-1}} h'_+(u, 0) dS_{K_0}(u) \\ &= l. \end{aligned}$$

This, together with Minkowski's mixed volume inequality (10), yields

$$l \geq \limsup_{t \rightarrow 0^+} \frac{V_1(K_0, K_t) - V(K_0)}{t} \geq \limsup_{t \rightarrow 0^+} \frac{V(K_0)^{1-\frac{1}{n}} V(K_t)^{\frac{1}{n}} - V(K_0)}{t},$$

and hence

$$l \geq V(K_0)^{1-\frac{1}{n}} \limsup_{t \rightarrow 0^+} \frac{V(K_t)^{\frac{1}{n}} - V(K_0)^{\frac{1}{n}}}{t}. \quad (20)$$

Combining (19) and (20) gives

$$l = V(K_0)^{1-\frac{1}{n}} \lim_{t \rightarrow 0^+} \frac{V(K_t)^{\frac{1}{n}} - V(K_0)^{\frac{1}{n}}}{t}. \quad (21)$$

Define a function $g : I \rightarrow \mathbb{R}$ by $g(t) = V(K_t)^{\frac{1}{n}}$. Identity (21) shows that the right derivative of g exists at 0. But this implies that the right derivative of g^n exists at 0 and that

$$\lim_{t \rightarrow 0^+} \frac{g(t)^n - g(0)^n}{t} = n g(0)^{n-1} \lim_{t \rightarrow 0^+} \frac{g(t) - g(0)}{t}.$$

Thus the definition of g and (21) show that

$$\lim_{t \rightarrow 0^+} \frac{V(K_t) - V(K_0)}{t} = nl,$$

which completes the proof of the lemma. \square

We shall require the following corollary of Lemma 1.

Corollary 1. *Let $I \subset \mathbb{R}$ be an interval containing 0 in its interior and let the function $h(t, u) : I \times S^{n-1} \rightarrow (0, \infty)$ be continuous and such that the convergence in*

$$h'(0, u) = \lim_{t \rightarrow 0} \frac{h(t, u) - h(0, u)}{t}$$

is uniform on S^{n-1} . If $\{K_t\}_{t \in I}$ is the family of Aleksandrov bodies associated with h , then

$$\lim_{t \rightarrow 0} \frac{V(K_t) - V(K_0)}{t} = \int_{S^{n-1}} h'(0, u) dS_{K_0}(u).$$

Proof. From Lemma 1 we see that we only need to show that

$$\lim_{t \rightarrow 0^-} \frac{V(K_t) - V(K_0)}{t} = \int_{S^{n-1}} h'(0, u) dS_{K_0}(u). \quad (22)$$

To that end, define $\tilde{h}(t, u) : -I \times S^{n-1} \rightarrow (0, \infty)$ by $\tilde{h}(t, u) = h(-t, u)$. For the corresponding family $\{\tilde{K}_{-t}\}_{t \in I}$ of Aleksandrov bodies associated with \tilde{h} we have $\tilde{K}_{-t} = K_t$ and $\tilde{K}_0 = K_0$. Thus, by Lemma 1,

$$\lim_{t \rightarrow 0^-} \frac{V(K_t) - V(K_0)}{-t} = \lim_{t \rightarrow 0^+} \frac{V(\tilde{K}_t) - V(\tilde{K}_0)}{t} = \int_{S^{n-1}} \tilde{h}'(0, u) dS_{K_0}(u).$$

Obviously, $\tilde{h}'(0, u) = -h'(0, u)$, which immediately implies (22). \square

The following lemma is a slight variant of a standard result about differentiability under an integral sign. For the sake of completeness, we include a proof.

Lemma 2. *Let $\phi : (0, \infty) \rightarrow (0, \infty)$ be continuously differentiable, $I \subset \mathbb{R}$ be an open interval, and*

$$h : I \times S^{n-1} \rightarrow (0, \infty), \quad (t, u) \mapsto h(t, u)$$

be a continuous function such that the partial derivative $\frac{\partial h}{\partial t}(t, u)$ exists for all $(t, u) \in I \times S^{n-1}$. If $\frac{\partial h}{\partial t}$ is bounded and h is bounded from above and from below by positive numbers, then the function $H : I \rightarrow (0, \infty)$ defined by

$$H(t) = \int_{S^{n-1}} (\phi \circ h)(t, u) d\mu(u),$$

for $t \in I$, is differentiable on I and

$$H'(t) = \int_{S^{n-1}} \frac{\partial(\phi \circ h)}{\partial t}(t, u) d\mu(u). \quad (23)$$

Moreover, if $\partial(\phi \circ h)/\partial t$ is continuous with respect to t , then H' is continuous.

Proof. Since ϕ' , the derivative of ϕ , is assumed to be continuous and h is bounded from above and from below by positive numbers, there exists a $c_1 \in (0, \infty)$ such that

$$|(\phi' \circ h)(t, u)| \leq c_1$$

for all $(t, u) \in I \times S^{n-1}$. This, the chain rule, and the assumption that $\frac{\partial h}{\partial t}$ is bounded show that there exists a $c_2 \in (0, \infty)$ such that

$$\left| \frac{\partial(\phi \circ h)}{\partial t}(t, u) \right| = \left| \left((\phi' \circ h) \cdot \frac{\partial h}{\partial t} \right)(t, u) \right| \leq c_2, \quad (24)$$

for all $(t, u) \in I \times S^{n-1}$.

Let $\{t_i\}$ be a sequence such that $\lim_{i \rightarrow \infty} t_i = t \in I$ with $t_i \neq t$ for all i . Set

$$f_i = \frac{(\phi \circ h)(t_i, \cdot) - (\phi \circ h)(t, \cdot)}{t_i - t}.$$

The mean value theorem shows that for each i and $u \in S^{n-1}$ there exists a $t'_i = t'_i(u) \in I$ such that

$$|f_i(u)| = \left| \frac{\partial(\phi \circ h)}{\partial t}(t'_i, u) \right| \leq c_2,$$

where the last inequality follows from (24). Since the measure μ is assumed to be finite, we have shown that all the $|f_i|$ are bounded from above by the same integrable function. The dominated convergence theorem applied to the sequence $\{f_i\}$ gives the differentiability of H and (23).

The last assertion of the lemma is again an immediate consequence of (24) and the dominated convergence theorem. \square

4. Orlicz norms

Throughout this section, let $\phi : [0, \infty) \rightarrow [0, \infty)$ be strictly increasing, continuously differentiable on $(0, \infty)$ with positive derivative, and satisfy $\lim_{t \rightarrow \infty} \phi(t) = \infty$. Note that under these assumptions, ϕ has an inverse $\phi^{-1} : \phi([0, \infty)) \rightarrow [0, \infty)$ which is continuously differentiable on $\phi((0, \infty))$.

Let μ be a finite Borel measure on the sphere S^{n-1} . For a continuous $f : S^{n-1} \rightarrow [0, \infty)$, the Orlicz norm $\|f\|_\phi$ is defined by

$$\|f\|_\phi = \inf \left\{ \lambda > 0 : \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f}{\lambda}\right) d\mu \leq \phi(1) \right\}. \quad (25)$$

Observe that the Orlicz norm of a function depends not only on ϕ but also on μ although this is not reflected in our notation. Moreover, since ϕ may not be convex, the Orlicz norm defined above may not satisfy the triangle inequality. The usual L_p norm is obtained by taking $\phi(t) = t^p$. The reader is referred to [57] for background on Orlicz norms.

Observe that for continuous $f : S^{n-1} \rightarrow [0, \infty)$,

$$\|cf\|_\phi = c\|f\|_\phi, \quad c > 0. \quad (26)$$

In particular we have

$$\|c\|_\phi = c, \quad c > 0. \quad (27)$$

Moreover, it follows immediately from the monotonicity of the function ϕ that for continuous functions $f, g : S^{n-1} \rightarrow [0, \infty)$,

$$f \leq g \implies \|f\|_\phi \leq \|g\|_\phi. \quad (28)$$

The following simple fact will prove useful.

Lemma 3. Suppose μ is a finite Borel measure on S^{n-1} and the function $f : S^{n-1} \rightarrow [0, \infty)$ is continuous and such that $\mu(\{f \neq 0\}) > 0$. Then the Orlicz norm $\|f\|_\phi$ of f is positive and

$$\|f\|_\phi = \lambda_0 \iff \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f}{\lambda_0}\right) d\mu = \phi(1).$$

Proof. Define a function $\psi : (0, \infty) \rightarrow [0, \infty)$, for $\lambda > 0$, by

$$\psi(\lambda) = \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f}{\lambda}\right) d\mu.$$

Since ϕ is strictly increasing and $\mu(\{f \neq 0\}) > 0$, the function ψ is strictly decreasing. It therefore has an inverse $\psi^{-1} : \psi((0, \infty)) \rightarrow (0, \infty)$. The dominated convergence theorem and the continuity of ϕ on $(0, \infty)$, show that ψ is continuous, as well.

The non-negativity of ϕ , Fatou's lemma, and the fact that $\lim_{t \rightarrow \infty} \phi(t) = \infty$ give

$$\liminf_{\lambda \rightarrow 0} \psi(\lambda) \geq \liminf_{\lambda \rightarrow 0} \frac{1}{|\mu|} \int_{\{f > 0\}} \phi\left(\frac{f}{\lambda}\right) d\mu \geq \frac{1}{|\mu|} \int_{\{f > 0\}} \liminf_{\lambda \rightarrow 0} \phi\left(\frac{f}{\lambda}\right) d\mu = \infty,$$

and hence

$$\lim_{\lambda \rightarrow 0} \psi(\lambda) = \infty. \quad (29)$$

Next, we show that

$$\phi(1) \in \psi((0, \infty)). \quad (30)$$

Since f is continuous on S^{n-1} , there exists a $c \in (0, \infty)$ with $f(u) < c$ for every $u \in S^{n-1}$. Thus, the monotonicity of ϕ yields

$$\psi(\lambda) = \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f}{\lambda}\right) d\mu < \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{c}{\lambda}\right) d\mu = \phi(c/\lambda),$$

for every positive λ . In particular, $\psi(2c) < \phi(1/2) < \phi(1)$. From (29) and the continuity of ψ we therefore deduce (30) by the intermediate value theorem.

Finally (30) and the strict monotonicity of ψ yield

$$\|f\|_{\phi} = \inf\{\lambda > 0: \psi(\lambda) \leq \phi(1)\} = \psi^{-1}(\phi(1)).$$

This shows that $\|f\|_{\phi} > 0$ and establishes the desired equivalence. \square

The continuity of the Orlicz norm is contained in the following lemma.

Lemma 4. Suppose $f \in C(S^{n-1})$ with $\mu(\{f \neq 0\}) > 0$ and $\{f_i\}$ is a sequence of non-negative functions in $C(S^{n-1})$. If

$$f_i \rightarrow f \quad \text{in } C(S^{n-1}),$$

then

$$\|f_i\|_{\phi} \rightarrow \|f\|_{\phi}.$$

Proof. From the uniform convergence, it follows that there exists a real $c > 0$ such that $f_i(u) \leq c$ for all i and all $u \in S^{n-1}$. From (27) and (28) we thus obtain

$$0 \leq \|f_i\|_{\phi} \leq \|c\|_{\phi} = c,$$

for every i . Thus the sequence $\{\|f_i\|_{\phi}\}$ is bounded. To show that the sequence converges to $\|f\|_{\phi}$, we prove that every convergent subsequence converges to $\|f\|_{\phi}$. Denote an arbitrary convergent subsequence of $\{\|f_i\|_{\phi}\}$ by $\{\|f_i\|_{\phi}\}$ as well.

To see that $\lim_{i \rightarrow \infty} \|f_i\|_\phi > 0$, suppose the contrary; i.e., $\|f_i\|_\phi \rightarrow 0$. Then Lemma 3, the non-negativity of ϕ , Fatou's lemma, and the fact that $\lim_{t \rightarrow \infty} \phi(t) = \infty$ would produce the desired contradiction:

$$\begin{aligned} \phi(1) &= \lim_{i \rightarrow \infty} \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f_i}{\|f_i\|_\phi}\right) d\mu \\ &\geq \liminf_{i \rightarrow \infty} \frac{1}{|\mu|} \int_{\{f>0\}} \phi\left(\frac{f_i}{\|f_i\|_\phi}\right) d\mu \\ &\geq \frac{1}{|\mu|} \int_{\{f>0\}} \liminf_{i \rightarrow \infty} \phi\left(\frac{f_i}{\|f_i\|_\phi}\right) d\mu \\ &= \infty. \end{aligned}$$

Thus $\lim_{i \rightarrow \infty} \|f_i\|_\phi > 0$. Note that since $\mu(\{f \neq 0\}) > 0$ and $f_i \rightarrow f$ in $C(S^{n-1})$ we have $\mu(\{f_i \neq 0\}) > 0$ for sufficiently large i . From the continuity of ϕ and Lemma 3 we therefore deduce

$$\frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f}{\lim_{i \rightarrow \infty} \|f_i\|_\phi}\right) d\mu = \lim_{i \rightarrow \infty} \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{f_i}{\|f_i\|_\phi}\right) d\mu = \phi(1).$$

Lemma 3 again shows $\lim_{i \rightarrow \infty} \|f_i\|_\phi = \|f\|_\phi$. \square

We will need the following basic fact.

Lemma 5. *If μ is a finite Borel measure on the sphere S^{n-1} which is not concentrated on a great subsphere of S^{n-1} , then there exists a real $c > 0$ such that $\|h_{\bar{v}}\|_\phi > c$ for every $v \in S^{n-1}$.*

Proof. Note that since μ is not concentrated on a great subsphere of S^{n-1} , we have, for every $v \in S^{n-1}$,

$$\mu(\{h_{\bar{v}} > 0\}) = \mu(S^{n-1} \setminus v^\perp) > 0,$$

where v^\perp denotes the codimension 1 subspace orthogonal to v . Hence Lemma 3 shows that $\|h_{\bar{v}}\|_\phi > 0$ for every unit vector v . In order to establish the assertion of the lemma, it suffices to show that the function $v \mapsto \|h_{\bar{v}}\|_\phi$ is continuous.

Suppose $v_i \in S^{n-1}$ and $v_i \rightarrow v$. That $h_{\bar{v}_i} \rightarrow h_{\bar{v}}$, uniformly on S^{n-1} is easily seen. Thus $\|h_{\bar{v}_i}\|_\phi$ converges to $\|h_{\bar{v}}\|_\phi$ by Lemma 4. This establishes the desired continuity of $v \mapsto \|h_{\bar{v}}\|_\phi$. \square

For a continuous $f : S^{n-1} \rightarrow [0, \infty)$ with $\|f\|_\phi > 0$, define

$$\langle f \rangle = \frac{f}{\|f\|_\phi}. \quad (31)$$

From (26) we see that we always have

$$\|\langle f \rangle\|_\phi = 1.$$

We will use the fact that $\langle h_K \rangle$ is homogeneous of degree 0 with respect to dilation of K . Indeed, from the definition of $\langle \cdot \rangle$, together with (3) and (26), we see that for every $c > 0$,

$$\langle h_K \rangle = \frac{h_K}{\|h_K\|_\phi} = \frac{h_{cK}}{\|h_{cK}\|_\phi} = \langle h_{cK} \rangle. \quad (32)$$

Lemma 3 shows that if μ is such that $\mu(\{f > 0\}) > 0$, then

$$\frac{1}{|\mu|} \int_{S^{n-1}} \phi \circ \langle f \rangle d\mu = \phi(1). \quad (33)$$

5. The even Orlicz Minkowski problem

The first of the two main results shows existence for the even Orlicz Minkowski problem under some mild assumptions regarding φ .

Theorem 1. *Suppose $\varphi : (0, \infty) \rightarrow (0, \infty)$ is a continuous decreasing function. If μ is an even finite Borel measure on S^{n-1} which is not concentrated on a great subsphere of S^{n-1} , then for each $\alpha \in (0, 1)$ there exists an origin symmetric convex body K in \mathbb{R}^n such that*

$$c\varphi(h_K) dS_K = d\mu,$$

with $c = V(K)^{\frac{\alpha}{n}-1}$.

Note that for $\varphi \equiv 1$, Theorem 1 provides a solution to the even Minkowski problem — by appealing to (6).

Proof of Theorem 1. Define the function $\phi : [0, \infty) \rightarrow [0, \infty)$ by

$$\phi(t) = \int_0^t \frac{1}{\varphi(s)} ds.$$

Observe that since φ is decreasing and $0 < \alpha < 1$, it follows that for $c > 0$

$$\lim_{t \rightarrow \infty} \phi(ct)/t^\alpha = \infty, \quad (34)$$

and

$$\lim_{t \rightarrow 0^+} \phi(t)/t^\alpha = 0. \quad (35)$$

Since φ is decreasing, the derivative of ϕ is increasing and therefore the function ϕ is convex.

Furthermore, for $c > 0$ let $\psi_c : (0, \infty) \rightarrow \mathbb{R}$ be defined by

$$\psi_c(t) = \left(\frac{n}{\alpha} V(B)^{\frac{\alpha}{n}} - \frac{|\mu|\phi(ct)}{t^\alpha} \right) t^\alpha.$$

From (34) we conclude that $\lim_{t \rightarrow \infty} \psi_c(t) = -\infty$. In particular, for each c there exists a real $r_c > 0$ such that

$$t > r_c \implies \psi_c(t) < 0. \quad (36)$$

Motivated by the work of Chou and Wang [7], we define the functional $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(f) = \frac{n}{\alpha} V(f)^{\frac{\alpha}{n}} - \int_{S^{n-1}} \phi \circ f \, d\mu,$$

for $f \in C_e^+(S^{n-1})$. Since, as seen in Section 3, the volume $V : C^+(S^{n-1}) \rightarrow \mathbb{R}$ is continuous, with respect to the max-norm metric, Φ is continuous as well.

Note that by (12), for every $r > 0$,

$$\Phi(h_{rB}) = \frac{n}{\alpha} r^\alpha V(B)^{\frac{\alpha}{n}} - \phi(r)|\mu| = \left(\frac{n}{\alpha} V(B)^{\frac{\alpha}{n}} - \frac{|\mu|\phi(r)}{r^\alpha} \right) r^\alpha.$$

From this and (35) we see that $\Phi(h_{rB})$ is positive for small positive r . Hence

$$\text{there exists } K \in \mathcal{K}_e^n \text{ such that } \Phi(h_K) > 0. \quad (37)$$

We are searching for a function at which Φ attains a maximum. The search can be restricted to support functions of origin symmetric convex bodies. To see this, recall from Section 3 that the Aleksandrov body K associated with a given function $h \in C_e^+(S^{n-1})$ is origin symmetric and has a support function h_K which satisfies $0 < h_K \leq h$. Since ϕ is increasing and $V(h) = V(h_K)$ by (12) we deduce $\Phi(h) \leq \Phi(h_K)$.

Next, we will show that the search for a function at which Φ attains a maximum can be further restricted to support functions of origin symmetric convex bodies contained in some ball of fixed radius. To this end, first note that the continuous function on S^{n-1} ,

$$v \mapsto \int_{S^{n-1}} h_{\bar{v}} \, d\mu,$$

is positive since μ is not concentrated on a great subsphere. Thus, there exists a $c \in (0, \infty)$ such that

$$\frac{1}{|\mu|} \int_{S^{n-1}} h_{\bar{v}} \, d\mu \geq c, \quad (38)$$

for every $v \in S^{n-1}$. Let $K \in \mathcal{K}_e^n$ and choose $v_K \in S^{n-1}$ such that for a suitable real $r_K > 0$ the point $r_K v_K$ is an element of K with maximal distance from the origin. Since K is origin

symmetric, the line segment with endpoints $\pm r_K v_K$ is contained in K . From (3) and (4) we deduce $r_K h_{\bar{v}_K} \leq h_K$. The monotonicity of ϕ , Jensen's inequality, and (38) therefore yield

$$\int_{S^{n-1}} \phi(h_K) d\mu \geq \int_{S^{n-1}} \phi(r_K h_{\bar{v}_K}) d\mu \geq |\mu| \phi\left(\frac{1}{|\mu|} \int_{S^{n-1}} r_K h_{\bar{v}_K} d\mu\right) \geq |\mu| \phi(cr_K).$$

Now (12), the fact that $K \subset r_K B$, and the last inequality show that

$$\begin{aligned} \Phi(h_K) &= \frac{n}{\alpha} V(K)^{\frac{\alpha}{n}} - \int_{S^{n-1}} \phi(h_K) d\mu \\ &\leq \frac{n}{\alpha} r_K^\alpha V(B)^{\frac{\alpha}{n}} - |\mu| \phi(cr_K) \\ &= \left(\frac{n}{\alpha} V(B)^{\frac{\alpha}{n}} - \frac{|\mu| \phi(cr_K)}{r_K^\alpha} \right) r_K^\alpha \\ &= \psi_c(r_K). \end{aligned}$$

From (36) we therefore conclude that there exists a real $r = r_c > 0$ such that

$$r_K > r \implies \Phi(h_K) < 0. \quad (39)$$

It follows from (37) and (39) that in order to find a maximum of the functional Φ on $C_e^+(S^{n-1})$, it is sufficient to search among support functions of members of the set

$$\mathcal{F} = \{K \in \mathcal{K}_e^n : K \subset rB\}.$$

Let $\{K_i\}$ be a maximizing sequence in \mathcal{F} for Φ , i.e.

$$\lim_{i \rightarrow \infty} \Phi(h_{K_i}) = \sup\{\Phi(h_K) : K \in \mathcal{F}\}.$$

Obviously, the sequence $\{K_i\}$ is bounded. Blaschke's selection theorem (see, e.g., [59]) guarantees the existence of a convergent subsequence, which we also denote by $\{K_i\}$, with $\lim_{i \rightarrow \infty} K_i = K_0$. Since the $K_i \in \mathcal{K}_e^n$, the body K_0 is origin symmetric. Moreover, the continuity of volume, (12), and the positivity of $\lim_{i \rightarrow \infty} \Phi(h_{K_i})$ yield

$$\frac{n}{\alpha} V(K_0)^{\frac{\alpha}{n}} = \lim_{i \rightarrow \infty} \frac{n}{\alpha} V(K_i)^{\frac{\alpha}{n}} = \lim_{i \rightarrow \infty} \frac{n}{\alpha} V(h_{K_i})^{\frac{\alpha}{n}} \geq \lim_{i \rightarrow \infty} \Phi(h_{K_i}) > 0.$$

Consequently, the body K_0 has non-empty interior and thus $K_0 \in \mathcal{K}_e^n$. The continuity of Φ now shows that

$$\Phi(f) \leq \Phi(h_{K_0}),$$

for every $f \in C_e^+(S^{n-1})$.

Suppose $f \in C_c^+(S^{n-1})$ is arbitrary but fixed. For sufficiently small $\delta > 0$ we can define a function, $h : (-\delta, \delta) \times S^{n-1} \rightarrow (0, \infty)$, which is bounded from above and below by positive reals, by

$$h_t(u) := h(t, u) = h_{K_0}(u) + tf(u).$$

By Lemma 2 and Corollary 1, the function $t \mapsto \Phi \circ h_t$ is differentiable at 0. Since K_0 is a maximizer of the functional Φ and $h_0 = h_{K_0}$ we have

$$\left. \frac{d}{dt} (\Phi \circ h_t) \right|_{t=0} = 0. \quad (40)$$

Note that by Corollary 1,

$$\left. \frac{d}{dt} \frac{n}{\alpha} V(h_t)^{\frac{\alpha}{n}} \right|_{t=0} = V(K_0)^{\frac{\alpha}{n}-1} \int_{S^{n-1}} f dS_{K_0},$$

and by Lemma 2,

$$\left. \frac{d}{dt} \int_{S^{n-1}} \phi \circ h_t d\mu \right|_{t=0} = \int_{S^{n-1}} \frac{1}{\phi \circ h_{K_0}} f d\mu.$$

The definition of Φ together with (40) therefore give

$$V(K_0)^{\frac{\alpha}{n}-1} \int_{S^{n-1}} f dS_{K_0} - \int_{S^{n-1}} \frac{1}{\phi \circ h_{K_0}} f d\mu = 0.$$

Since this holds for all positive even continuous functions f on S^{n-1} , it holds for all even continuous f on S^{n-1} and hence

$$V(K_0)^{\frac{\alpha}{n}-1} \phi(h_{K_0}) dS_{K_0} = d\mu. \quad \square$$

We now establish the following variant of our first theorem.

Theorem 2. Suppose $\varphi : (0, \infty) \rightarrow (0, \infty)$ is a continuous function such that $\phi(t) = \int_0^t 1/\varphi(s) ds$ exists for every positive t and is unbounded as $t \rightarrow \infty$. If μ is an even finite Borel measure on S^{n-1} which is not concentrated on a great subsphere of S^{n-1} , then there exist an origin symmetric convex body K in \mathbb{R}^n and a $c > 0$ such that

$$c\varphi(h_K) dS_K = d\mu, \quad (41)$$

and

$$\|h_K\|_\phi = 1. \quad (42)$$

In order to establish this theorem we first establish four lemmas.

First extend the definition of ϕ to $[0, \infty)$ by

$$\phi(t) = \int_0^t \frac{1}{\varphi(s)} ds, \quad \text{for } t > 0, \quad \text{and} \quad \phi(0) = \lim_{t \rightarrow 0^+} \phi(t).$$

The function ϕ is strictly increasing and continuously differentiable on $(0, \infty)$, and $\phi' > 0$. Thus ϕ has an inverse $\phi^{-1} : \phi([0, \infty)) \rightarrow [0, \infty)$ which is continuously differentiable on $\phi((0, \infty))$. Observe that $\lim_{t \rightarrow \infty} \phi(t) = \infty$.

Suppose that $f \in C^+(S^{n-1})$ and $K \in \mathcal{K}_o^n$. Recall that the function ϕ is strictly increasing. There exists $\delta > 0$ so that

$$\phi(1) - tf(u) \in \phi\left(\left(\frac{1}{2}, 2\right)\right),$$

for all $(t, u) \in (-\delta, \delta) \times S^{n-1}$. Define the function $\tilde{h} : (-\delta, \delta) \times S^{n-1} \rightarrow (0, \infty)$, at (t, u) , by

$$\tilde{h}(t, u) = \frac{h_K(u)}{\phi^{-1}(\phi(1) - tf(u))}. \quad (43)$$

Clearly, \tilde{h} is bounded from above and below by positive reals (dependent on f and K). From (27) and (28) we therefore have that also $\|\tilde{h}_t\|_\phi$ is bounded from above and below by positive reals.

The following lemma follows directly from the definitions.

Lemma 6. *The function \tilde{h} is continuous and the derivative $\frac{\partial \tilde{h}}{\partial t}(t, u)$ exists on $(-\delta, \delta) \times S^{n-1}$ and is given by*

$$\frac{\partial \tilde{h}}{\partial t}(t, u) = \frac{h_K(u) f(u) (\phi^{-1})'(\phi(1) - tf(u))}{[\phi^{-1}(\phi(1) - tf(u))]^2}. \quad (44)$$

Moreover, it is continuous and bounded from above and below by positive reals.

We now establish:

Lemma 7. *The function $t \mapsto V(\tilde{h}_t)$ is differentiable at 0.*

Proof. By Lemma 6, the function \tilde{h} is continuous. So in order to apply Corollary 1 it remains to show that the convergence in

$$\lim_{t \rightarrow 0} \frac{\tilde{h}_t(u) - \tilde{h}_0(u)}{t} \quad (45)$$

is uniform on S^{n-1} . Let $\delta' \in (0, \delta)$. By the mean value theorem, for every $t \in [-\delta', \delta']$ we can find a $t' \in [-t, t]$ such that

$$\left| \frac{\tilde{h}_t(u) - \tilde{h}_0(u)}{t} - \tilde{h}'_0(u) \right| = \left| \frac{\partial \tilde{h}}{\partial t}(t', u) - \frac{\partial \tilde{h}}{\partial t}(0, u) \right|.$$

Since $[-\delta', \delta'] \times S^{n-1}$ is compact, Lemma 6 shows that the partial derivative $\frac{\partial \tilde{h}}{\partial t}(t, u)$ is uniformly continuous on $[-\delta', \delta'] \times S^{n-1}$. Hence, the convergence in (45) is uniform on S^{n-1} . \square

Also needed will be:

Lemma 8. *The function $t \mapsto \|\tilde{h}_t\|_\phi$ is differentiable on $(-\delta, \delta)$ with bounded derivative.*

Proof. As previously mentioned (after the definition of \tilde{h}) there exist $c_1, c_2 \in (0, \infty)$ such that

$$c_1 < \tilde{h}_t(u) < c_2, \quad \text{for all } (t, u) \in (-\delta, \delta) \times S^{n-1}.$$

Define the function $G : (-\delta, \delta) \times (c_1, c_2) \rightarrow \mathbb{R}$ at (t, λ) by

$$G(t, \lambda) = \frac{1}{|\mu|} \int_{S^{n-1}} \phi\left(\frac{\tilde{h}_t}{\lambda}\right) d\mu - \phi(1).$$

Furthermore, for fixed $\lambda \in (c_1, c_2)$, define $h_1 : (-\delta, \delta) \times S^{n-1} \rightarrow (0, \infty)$ by

$$h_1(t, u) = \tilde{h}(t, u)/\lambda,$$

and, for fixed $t \in (-\delta, \delta)$, define $h_2 : (c_1, c_2) \times S^{n-1} \rightarrow (0, \infty)$ by

$$h_2(\lambda, u) = \tilde{h}(t, u)/\lambda.$$

Clearly, the functions h_1 and h_2 are bounded from above and below by positive reals. Moreover, by Lemma 6, the derivatives $\frac{\partial h_1}{\partial t}$ and $\frac{\partial h_2}{\partial \lambda}$ exist, are continuous, and bounded on their domains. Thus, by applying Lemma 2 to h_1 and h_2 respectively, the partial derivatives $\partial G/\partial t$, $\partial G/\partial \lambda$ exist. Since $\partial(\phi \circ h_1)/\partial t$ and $\partial(\phi \circ h_2)/\partial \lambda$ are continuous by Lemma 6, another application of Lemma 2 shows that $\partial G/\partial t$ and $\partial G/\partial \lambda$ are in fact continuous. Since by Lemma 2 interchanging differentiation and integration are permitted, an elementary calculation shows that $\partial G/\partial t$ and $-\partial G/\partial \lambda$ are bounded from above and below by positive reals. In particular, $\partial G/\partial \lambda$ is always nonzero.

Let $t \in (-\delta, \delta)$. For sufficiently small ε , Lemma 3 and the mean value theorem give

$$0 = G(t + \varepsilon, \|\tilde{h}_{t+\varepsilon}\|_\phi) - G(t, \|\tilde{h}_t\|_\phi) = \varepsilon \frac{\partial G}{\partial t}(x_\varepsilon) + (\|\tilde{h}_{t+\varepsilon}\|_\phi - \|\tilde{h}_t\|_\phi) \frac{\partial G}{\partial \lambda}(x_\varepsilon), \quad (46)$$

where x_ε is a point on the line segment joining the points $(t, \|\tilde{h}_t\|_\phi)$ and $(t + \varepsilon, \|\tilde{h}_{t+\varepsilon}\|_\phi)$.

Note that from the continuity of \tilde{h} , and compactness, follows the uniform continuity of $\tilde{h} : [-\delta', \delta'] \times S^{n-1}$ for arbitrary $0 < \delta' < \delta$. Thus the convergence in $\lim_{\varepsilon \rightarrow 0} \tilde{h}_{t+\varepsilon} = \tilde{h}_t$ is uniform and hence, by Lemma 4, $\lim_{\varepsilon \rightarrow 0} \|\tilde{h}_{t+\varepsilon}\|_\phi = \|\tilde{h}_t\|_\phi$. Hence $x_\varepsilon \rightarrow (t, \|\tilde{h}_t\|_\phi)$ as $\varepsilon \rightarrow 0$.

Since, as shown above, the partial derivatives of G are continuous and $\partial G/\partial \lambda$ is always nonzero, we conclude from (46) that $\|\tilde{h}_t\|_\phi$ is differentiable with

$$\frac{d\|\tilde{h}_t\|_\phi}{dt} = - \left[\frac{\partial G}{\partial t}(t, \|\tilde{h}_t\|_\phi) \right] / \left[\frac{\partial G}{\partial \lambda}(t, \|\tilde{h}_t\|_\phi) \right].$$

The fact that $\partial G/\partial t$ and $-\partial G/\partial \lambda$ are bounded from above and below by positive reals, finally shows that $d\|\tilde{h}_t\|_\phi/dt$ is bounded. \square

In the final lemma needed for the proof of Theorem 2, we establish some differentiability properties of functions involving \tilde{h} . Recall that $\langle \tilde{h}_t \rangle = \tilde{h}_t / \|\tilde{h}_t\|_\phi$.

Lemma 9. *The functions*

$$t \mapsto \|\tilde{h}_t\|_\phi / V(\tilde{h}_t)^{1/n} \quad \text{and} \quad t \mapsto \int_{S^{n-1}} \phi \circ \langle \tilde{h}_t \rangle d\mu$$

are differentiable at 0 with derivatives

$$\frac{1}{V(\tilde{h}_0)^{\frac{1}{n}}} \left(\frac{d}{dt} \|\tilde{h}_t\|_\phi \Big|_{t=0} - \frac{\|\tilde{h}_0\|_\phi}{nV(\tilde{h}_0)} \int_{S^{n-1}} \frac{\tilde{h}_0 f}{\phi'(1)} dS_K \right) \quad (47)$$

and

$$\frac{1}{\|\tilde{h}_0\|_\phi} \int_{S^{n-1}} \frac{1}{\varphi(\langle \tilde{h}_0 \rangle)} \left(\frac{\tilde{h}_0 f}{\phi'(1)} - \langle \tilde{h}_0 \rangle \frac{d}{dt} \|\tilde{h}_t\|_\phi \Big|_{t=0} \right) d\mu, \quad (48)$$

respectively.

Proof. First observe that \tilde{h}_t is bounded from below by a positive real so its Aleksandrov body contains the origin in its interior. The differentiability, at 0, of the function

$$t \mapsto \|\tilde{h}_t\|_\phi / V(\tilde{h}_t)^{1/n}$$

is an immediate consequence of Lemmas 7 and 8. Formula (47) for the respective derivative follows directly from (44), Corollary 1, and

$$(\phi^{-1})'(\phi(1)) = \frac{1}{\phi'(1)}. \quad (49)$$

Since \tilde{h} and $\|\tilde{h}_t\|_\phi$ are bounded from below and above by positive reals, so is $\langle \tilde{h}_t \rangle$. From Lemmas 6 and 8 we infer that $\langle \tilde{h}_t \rangle$ is differentiable with respect to t . An elementary calculation shows that

$$\frac{\partial \langle \tilde{h}_t \rangle}{\partial t} = \frac{1}{\|\tilde{h}_t\|_\phi} \left(\frac{d\tilde{h}_t}{dt} - \langle \tilde{h}_t \rangle \frac{d\|\tilde{h}_t\|_\phi}{dt} \right). \quad (50)$$

Since $\|\tilde{h}_t\|_\phi$ is bounded from below and above by positive reals and $\langle \tilde{h}_t \rangle$, $d\tilde{h}_t/dt$, $d\|\tilde{h}_t\|_\phi/dt$ are bounded, we therefore conclude that $\partial \langle \tilde{h}_t \rangle / \partial t$ is bounded as well. Lemma 2 yields the differentiability, at 0, of

$$t \mapsto \int_{S^{n-1}} \phi \circ \langle \tilde{h}_t \rangle d\mu,$$

and together with (44), (49), and (50), the desired formula (48). \square

We are now in a position to establish Theorem 2.

Proof of Theorem 2. For $f \in C_e^+(S^{n-1})$, define the functional $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(f) = \|f\|_\phi / V(f)^{1/n}.$$

Lemma 4, together with the fact that $V : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ is continuous, shows that Φ is continuous as well.

We are searching for a function at which Φ attains a minimum. As before, the search can be restricted to support functions of origin symmetric convex bodies: Indeed, recall that the Aleksandrov body K associated with a given function $h \in C_e^+(S^{n-1})$ is origin symmetric and has a support function h_K which satisfies $0 < h_K \leq h$. The fact that $V(h) = V(h_K)$ together with (28) shows that $\Phi(h_K) \leq \Phi(h)$.

Since Φ is positively homogeneous of degree 0, the search can be further restricted to support functions of convex bodies of unit volume. Let $c_1 = \Phi(h_{B'})$, where B' is the dilate of B chosen so that $V(B') = 1$. It follows that in order to find a minimum of the functional Φ on $C_e^+(S^{n-1})$, it is sufficient to search among the support functions of the members of the set

$$\mathcal{F} = \{K \in \mathcal{K}_e^n : \Phi(h_K) \leq c_1 \text{ and } V(K) = 1\}.$$

Let $\{K_i\}$ be a minimizing sequence, of bodies in \mathcal{F} , for the functional Φ , i.e.,

$$\lim_{i \rightarrow \infty} \Phi(h_{K_i}) = \inf\{\Phi(h_K) : K \in \mathcal{F}\}.$$

We now show that the sequence $\{K_i\}$ is bounded. For each i , let $v_i \in S^{n-1}$ be chosen such that for suitable $r_i > 0$ the points $r_i v_i$ are elements of K_i with maximal distance from the origin. Since each K_i is origin symmetric, the segments with endpoints $\pm r_i v_i$ are contained in K_i . From (3) and (4) we deduce $r_i h_{\bar{v}_i} \leq h_{K_i}$. Hence (26), (28), and the fact that the K_i belong to \mathcal{F} imply

$$r_i \|h_{\bar{v}_i}\|_\phi = \|r_i h_{\bar{v}_i}\|_\phi \leq \|h_{K_i}\|_\phi = \Phi(h_{K_i}) \leq c_1. \quad (51)$$

By Lemma 5, there exists a $c_2 \in (0, \infty)$ such that

$$c_2 \leq \|h_{\bar{v}_i}\|_\phi \quad (52)$$

for all i . Combining (51) and (52) we see that the r_i are bounded from above and hence the sequence $\{K_i\}$ is bounded.

Now Blaschke's selection theorem guarantees the existence of a convergent subsequence of $\{K_i\}$, which we also denote by $\{K_i\}$, with $\lim_{i \rightarrow \infty} K_i = K_0$. Clearly, the body K_0 is again an origin symmetric compact, convex set. Since obviously $V(K_0) = 1$, we see that $K_0 \in \mathcal{K}_e^n$. The continuity of Φ now implies that $K_0 \in \mathcal{F}$ and thus that

$$\Phi(h_{K_0}) \leq \Phi(f),$$

for every $f \in C_e^+(S^{n-1})$.

Choose a fixed $f \in C_e^+(S^{n-1})$. As in (43), define $\tilde{h} : (-\delta, \delta) \times S^{n-1} \rightarrow (0, \infty)$ by

$$\tilde{h}(t, u) = \frac{h_{K_0}(u)}{\phi^{-1}(\phi(1) - tf(u))}.$$

Lemma 9 shows that the function $t \mapsto \Phi \circ \tilde{h}_t$ is differentiable at 0. Since h_{K_0} is a minimizer of the functional Φ and $\tilde{h}_0 = h_{K_0}$ we have

$$\left. \frac{d}{dt} (\Phi \circ \tilde{h}_t) \right|_{t=0} = 0.$$

The expression for the above derivative given in (47) implies

$$\left. \frac{d}{dt} \|\tilde{h}_t\|_\phi \right|_{t=0} = \frac{\|\tilde{h}_0\|_\phi}{nV(\tilde{h}_0)} \int_{S^{n-1}} \frac{\tilde{h}_0 f}{\phi'(1)} dS_{K_0}. \quad (53)$$

For each t such that $|t| < \delta$, we have from (33)

$$\frac{1}{|\mu|} \int_{S^{n-1}} \phi \circ \langle \tilde{h}_t \rangle d\mu = \phi(1).$$

Thus, the derivative (with respect to t) of the function on the left is 0. This fact, at $t = 0$, together with (48) now gives

$$\int_{S^{n-1}} \frac{1}{\phi \circ \langle \tilde{h}_0 \rangle} \left(\frac{\tilde{h}_0 f}{\phi'(1)} - \langle \tilde{h}_0 \rangle \left. \frac{d}{dt} \|\tilde{h}_t\|_\phi \right|_{t=0} \right) d\mu = 0.$$

In this substitute the value of the derivative given by (53), use (12), and the fact that $\tilde{h}_0 = h_{K_0}$, to get

$$\frac{1}{nV(K_0)} \int_{S^{n-1}} h_{K_0} f dS_{K_0} \int_{S^{n-1}} \frac{h_{K_0}}{\phi \circ \langle h_{K_0} \rangle} d\mu = \int_{S^{n-1}} \frac{h_{K_0} f}{\phi \circ \langle h_{K_0} \rangle} d\mu.$$

By (3), (6), (32), and the homogeneity of volume (of degree n), the last equation remains unchanged if we replace K_0 by a dilate of K_0 . In particular, if we choose a dilate K'_0 of K_0 such that

$$\|h_{K'_0}\|_\phi = 1,$$

then we obtain

$$c \int_{S^{n-1}} h_{K'_0} f dS_{K'_0} = \int_{S^{n-1}} \frac{h_{K'_0} f}{\varphi(h_{K'_0})} d\mu,$$

where

$$c = \frac{1}{nV(K'_0)} \int_{S^{n-1}} \frac{h_{K'_0}}{\varphi(h_{K'_0})} d\mu.$$

Since this holds for all positive even continuous f on S^{n-1} , it holds for all even continuous f on S^{n-1} , and hence

$$c dS_{K'_0} = \frac{1}{\varphi(h_{K'_0})} d\mu,$$

or equivalently

$$c\varphi(h_{K'_0}) dS_{K'_0} = d\mu. \quad \square$$

From Theorem 2, we obtain the solution to the even L_p Minkowski problem for all positive p , when $0 < p \neq n$, and the solution to the even volume-normalized L_p Minkowski problem for all positive p .

Corollary 2. *If μ is an even finite Borel measure on the sphere S^{n-1} which is not concentrated on a great subsphere of S^{n-1} , then*

(i) *for $p > 0$, there exists an origin symmetric convex body K in \mathbb{R}^n such that*

$$ch_K^{1-p} dS_K = d\mu, \tag{54}$$

where $c = 1/V(K)$;

(ii) *for $0 < p \neq n$, there exists an origin symmetric convex body K in \mathbb{R}^n such that*

$$h_K^{1-p} dS_K = d\mu.$$

Proof. Note that from (3) and (6) it follows that the maps

$$K \mapsto h_K^{1-p} dS_K \quad \text{and} \quad K \mapsto \frac{1}{V(K)} h_K^{1-p} dS_K$$

are homogeneous of degree $n - p$ and $-p$, respectively; i.e., for $K \in \mathcal{K}_e^n$ and $\lambda > 0$,

$$h_{\lambda K}^{1-p} dS_{\lambda K} = \lambda^{n-p} h_K^{1-p} dS_K, \tag{55}$$

and

$$\frac{1}{V(\lambda K)} h_{\lambda K}^{1-p} dS_{\lambda K} = \lambda^{-p} \frac{1}{V(K)} h_K^{1-p} dS_K. \quad (56)$$

We first prove (i). Take $\varphi(t) = t^{1-p}$. From (41) we have (54) and from (42) that h_K satisfies

$$\frac{1}{|\mu|} \int_{S^{n-1}} h_K^p d\mu = 1. \quad (57)$$

We now show that a dilation of K gives the desired constant. By using (9), integration in (41) gives

$$\int_{S^{n-1}} h_K^p d\mu = cnV(K). \quad (58)$$

From (57) and (58) it follows that $1/c = nV(K)/|\mu|$. To complete the proof, let $K' = \lambda K$, where $\lambda^p = n/|\mu|$ is chosen so that by (56)

$$\frac{|\mu|}{nV(K)} h_K^{1-p} dS_K = \frac{1}{V(K')} h_{K'}^{1-p} dS_{K'}.$$

To see that (ii) follows from (i), observe that by (55),

$$ch_K^{1-p} dS_K = h_{K'}^{1-p} dS_{K'},$$

where $K' = \lambda K$, where λ is chosen so that $\lambda^{n-p} = c$. \square

References

- [1] S. Campi, P. Gronchi, The L^p -Busemann–Petty centroid inequality, *Adv. Math.* 167 (2002) 128–141, MR1901248, Zbl 1002.52005.
- [2] S. Campi, P. Gronchi, On the reverse L^p -Busemann–Petty centroid inequality, *Mathematika* 49 (2002) 1–11, MR2059037, Zbl 1056.52005.
- [3] S. Campi, P. Gronchi, On volume product inequalities for convex sets, *Proc. Amer. Math. Soc.* 134 (2006) 2393–2402, MR2213713, Zbl 1095.52002.
- [4] S. Campi, P. Gronchi, Volume inequalities for L_p -zonotopes, *Mathematika* 53 (2006) 71–80, (2007), MR2304053, Zbl 1117.52011.
- [5] W. Chen, L_p Minkowski problem with not necessarily positive data, *Adv. Math.* 201 (2006) 77–89, MR2204749, Zbl 1102.34023.
- [6] S.-Y. Cheng, S.-T. Yau, On the regularity of the solutions of the n -dimensional Minkowski problem, *Comm. Pure Appl. Math.* 29 (1976) 495–516, MR423267, Zbl 0363.53030.
- [7] K.-S. Chou, X.-J. Wang, The L_p -Minkowski problem and the Minkowski problem in centroaffine geometry, *Adv. Math.* 205 (2006) 33–83, MR2254308, Zbl pre05054348.
- [8] A. Cianchi, E. Lutwak, D. Yang, G. Zhang, Affine Moser–Trudinger and Morrey–Sobolev inequalities, *Calc. Var. Partial Differential Equations* 36 (2009) 419–436.
- [9] N. Dafnis, G. Paouris, Small ball probability estimates, Ψ_2 -behavior and the hyperplane conjecture, *J. Funct. Anal.* 258 (2010) 1933–1964.
- [10] B. Fleury, O. Guédon, G.A. Paouris, A stability result for mean width of L_p -centroid bodies, *Adv. Math.* 214 (2007) 865–877, MR2349721, Zbl 1132.52012.

- [11] R.J. Gardner, The Brunn–Minkowski inequality, *Bull. Amer. Math. Soc. (N.S.)* 39 (2002) 355–405, MR1898210, Zbl 1019.26008.
- [12] R.J. Gardner, *Geometric Tomography*, 2nd edition, *Encyclopedia Math. Appl.*, vol. 58, Cambridge University Press, Cambridge, 2006.
- [13] A. Giannopoulos, A. Pajor, G. Paouris, A note on subgaussian estimates for linear functionals on convex bodies, *Proc. Amer. Math. Soc.* 135 (2007) 2599–2606, MR2302581, Zbl 1120.52003.
- [14] P.M. Gruber, *Convex and Discrete Geometry*, *Grundlehren Math. Wiss.*, vol. 336, Springer, Berlin, 2007, MR2335496, Zbl 1139.52001.
- [15] P. Guan, C.-S. Lin, On equation $\det(u_{ij} + \delta_{ij}u) = u^p f$ on S^n , preprint.
- [16] C. Haberl, L_p intersection bodies, *Adv. Math.* 217 (2008) 2599–2624, MR2397461, Zbl 1140.52003.
- [17] C. Haberl, Star body valued valuations, *Indiana Univ. Math. J.* 58 (2009) 2253–2276.
- [18] C. Haberl, M. Ludwig, A characterization of L_p intersection bodies, *Int. Math. Res. Not. IMRN* 17 (2006) 29, Article ID 10548, Zbl 1115.52006.
- [19] C. Haberl, F. Schuster, General L_p affine isoperimetric inequalities, *J. Differential Geom.* 83 (2009) 1–26.
- [20] C. Haberl, F. Schuster, Asymmetric affine L_p Sobolev inequalities, *J. Funct. Anal.* 257 (2009) 641–658.
- [21] C. Haberl, F. Schuster, J. Xiao, An asymmetric affine Pólya–Szegő principle, arXiv:0908.1557.
- [22] C. Hu, X.-N. Ma, C. Shen, On the Christoffel–Minkowski problem of Firey’s p -sum, *Calc. Var. Partial Differential Equations* 21 (2004) 137–155, MR2085300, Zbl pre02113880.
- [23] D. Hug, E. Lutwak, D. Yang, G. Zhang, On the L_p Minkowski problem for polytopes, *Discrete Comput. Geom.* 33 (2005) 699–715, MR2132298, Zbl 1078.52008.
- [24] M.-Y. Jiang, Remarks on the 2-dimensional L_p -Minkowski problem, preprint.
- [25] D. Klain, The Minkowski problem for polytopes, *Adv. Math.* 185 (2004) 270–288, MR2060470, Zbl 1053.52015.
- [26] M. Ludwig, Projection bodies and valuations, *Adv. Math.* 172 (2002) 158–168, MR1942402, Zbl 1019.52003.
- [27] M. Ludwig, Valuations on polytopes containing the origin in their interiors, *Adv. Math.* 170 (2002) 239–256, MR1932331, Zbl 1015.52012.
- [28] M. Ludwig, Ellipsoids and matrix-valued valuations, *Duke Math. J.* 119 (2003) 159–188, MR1991649, Zbl 1033.52012.
- [29] M. Ludwig, Minkowski valuations, *Trans. Amer. Math. Soc.* 357 (2005) 4191–4213, MR2159706, Zbl 1077.52005.
- [30] M. Ludwig, Intersection bodies and valuations, *Amer. J. Math.* 128 (2006) 1409–1428, MR2275906, Zbl 1115.52007.
- [31] M. Ludwig, General affine surface areas, *Adv. Math.*, in press.
- [32] M. Ludwig, M. Reitzner, A classification of $SL(n)$ invariant valuations, *Ann. of Math.*, in press.
- [33] E. Lutwak, The Brunn–Minkowski–Firey theory. I. Mixed volumes and the Minkowski problem, *J. Differential Geom.* 38 (1993) 131–150, MR1231704, Zbl 0788.52007.
- [34] E. Lutwak, The Brunn–Minkowski–Firey theory. II. Affine and geominimal surface areas, *Adv. Math.* 118 (1996) 244–294, MR1378681, Zbl 0853.52005.
- [35] E. Lutwak, V. Oliker, On the regularity of solutions to a generalization of the Minkowski problem, *J. Differential Geom.* 41 (1995) 227–246, MR1316557, Zbl 0867.52003.
- [36] E. Lutwak, G. Zhang, Blaschke–Santaló inequalities, *J. Differential Geom.* 47 (1997) 1–16, MR1601426, Zbl 0906.52003.
- [37] E. Lutwak, D. Yang, G. Zhang, L_p affine isoperimetric inequalities, *J. Differential Geom.* 56 (2000) 111–132, MR1863023, Zbl 1034.52009.
- [38] E. Lutwak, D. Yang, G. Zhang, A new ellipsoid associated with convex bodies, *Duke Math. J.* 104 (2000) 375–390, MR1781476, Zbl 0974.52008.
- [39] E. Lutwak, D. Yang, G. Zhang, The Cramer–Rao inequality for star bodies, *Duke Math. J.* 112 (2002) 59–81, MR1890647, Zbl 1021.52008.
- [40] E. Lutwak, D. Yang, G. Zhang, Sharp affine L_p Sobolev inequalities, *J. Differential Geom.* 62 (2002) 17–38, MR1987375, Zbl 1073.46027.
- [41] E. Lutwak, D. Yang, G. Zhang, On the L_p -Minkowski problem, *Trans. Amer. Math. Soc.* 356 (11) (2004) 4359–4370, MR2067123, Zbl 1069.52010.
- [42] E. Lutwak, D. Yang, G. Zhang, Volume inequalities for subspaces of L_p , *J. Differential Geom.* 68 (2004) 159–184, MR2152912, Zbl 1119.52006.
- [43] E. Lutwak, D. Yang, G. Zhang, L^p John ellipsoids, *Proc. Lond. Math. Soc.* 90 (2005) 497–520, MR2142136, Zbl 1074.52005.
- [44] E. Lutwak, D. Yang, G. Zhang, Optimal Sobolev norms and the L^p Minkowski problem, *Int. Math. Res. Not. IMRN* 62987 (2006) 1–21, MR2211138, Zbl 1110.46023.

- [45] E. Lutwak, D. Yang, G. Zhang, Volume inequalities for isotropic measures, *Amer. J. Math.* 129 (2007) 1711–1723, MR2369894, Zbl 1134.52010.
- [46] E. Lutwak, D. Yang, G. Zhang, Orlicz projection bodies, *Adv. Math.* 223 (2010) 220–242.
- [47] E. Lutwak, D. Yang, G. Zhang, Orlicz centroid bodies, *J. Differential Geom.*, in press.
- [48] Y. Martinez-Maure, Hedgehogs and zonoids, *Adv. Math.* 158 (2001) 1–17, MR1814896, Zbl 0977.52010.
- [49] M. Meyer, E. Werner, On the p -affine surface area, *Adv. Math.* 152 (2000) 288–313, MR1764106, Zbl 0964.52005.
- [50] I. Molchanov, Convex and star-shaped sets associated with multivariate stable distributions. I. Moments and densities, *J. Multivariate Anal.* 100 (2009) 2195–2213.
- [51] V. Oliker, Generalized convex bodies and generalized envelopes, in: *Contemp. Math.*, vol. 140, 1992, pp. 105–113, MR1197592, Zbl 0789.52007.
- [52] G. Paouris, On the ψ_2 -behaviour of linear functionals on isotropic convex bodies, *Studia Math.* 168 (2005) 285–299, MR2146128, Zbl 1078.52501.
- [53] G. Paouris, Concentration of mass on convex bodies, *Geom. Funct. Anal.* 16 (2006) 1021–1049, MR2276533, Zbl 1114.52004.
- [54] G. Paouris, Concentration of mass on isotropic convex bodies, *C. R. Math. Acad. Sci. Paris* 342 (2006) 179–182.
- [55] G. Paouris, Small ball probability estimates for log-concave measures, preprint.
- [56] A.V. Pogorelov, *The Minkowski Multidimensional Problem*, V.H. Winston, Washington, DC, 1978, MR478079, Zbl 0387.53023.
- [57] M.M. Rao, Z.D. Ren, *Theory of Orlicz Spaces*, Monogr. Textbooks Pure Appl. Math., vol. 146, Marcel Dekker, New York, 1991, MR1113700, Zbl 0724.46032.
- [58] D. Ryabogin, A. Zvavitch, The Fourier transform and Firey projections of convex bodies, *Indiana Univ. Math. J.* 53 (2004) 667–682, MR2086696, Zbl 1062.52004.
- [59] R. Schneider, *Convex Bodies: The Brunn–Minkowski Theory*, Encyclopedia Math. Appl., vol. 44, Cambridge University Press, Cambridge, 1993, MR1216521, Zbl 0798.52001.
- [60] C. Schütt, E. Werner, Surface bodies and p -affine surface area, *Adv. Math.* 187 (2004) 98–145, MR2074173, Zbl 1089.52002.
- [61] A. Stancu, The discrete planar L_0 -Minkowski problem, *Adv. Math.* 167 (2002) 160–174, MR1901250, Zbl 1005.52002.
- [62] A. Stancu, On the number of solutions to the discrete two-dimensional L_0 -Minkowski problem, *Adv. Math.* 180 (2003) 290–323, MR2019226, Zbl 1054.52001.
- [63] A. Stancu, The necessary condition for the discrete L_0 -Minkowski problem in \mathbb{R}^2 , *J. Geom.* 88 (2008) 162–168, MR2398486, Zbl 1132.52300.
- [64] A.C. Thompson, *Minkowski Geometry*, Encyclopedia Math. Appl., vol. 63, Cambridge University Press, Cambridge, 1996, MR1406315, Zbl 0868.52001.
- [65] V. Umanskiy, On solvability of two-dimensional L_p -Minkowski problem, *Adv. Math.* 180 (2003) 176–186, MR2019221, Zbl 1048.52001.
- [66] E. Werner, D. Ye, New L_p affine isoperimetric inequalities, *Adv. Math.* 218 (2008) 762–780, MR2414321, Zbl 1155.52002.
- [67] E. Werner, D. Ye, Inequalities for mixed p -affine surface area, *Math. Ann.*, in press.
- [68] V. Yaskin, M. Yaskina, Centroid bodies and comparison of volumes, *Indiana Univ. Math. J.* 55 (2006) 1175–1194, MR2244603, Zbl 1102.52005.
- [69] G. Zhang, The affine Sobolev inequality, *J. Differential Geom.* 53 (1999) 183–202, MR1776095, Zbl 1040.53089.