

# Introduction to the Fourier Transform

## Wave Voodoo

David Nash

William Lyon Mackenzie C.I. Math Club

© David Nash, 2018



# Table of Contents

1 Introduction

2 Fourier Series

3 Fourier Transform



# Background Knowledge

## Introduction

- Even/Odd Functions
- Periodic Functions
- Trigonometry
- Complex Numbers (and  $e^{i\theta}$ )
- Integration and IBP



# What is a Fourier Series?

## Introduction

- Decomposition of a periodic function into sines and cosines
- Expressed as

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \left( \frac{\pi n x}{L} \right) + b_n \sin \left( \frac{\pi n x}{L} \right) \right)$$

- or

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin \left( \frac{\pi n x}{L} + \phi_n \right)$$

- or

$$\sum_{n=-\infty}^{\infty} c_n e^{i \frac{\pi n x}{L}}$$



# What is a Fourier Transform?

## Introduction

- Transformation of a function of time to a function of frequency
- Expressed as

$$\hat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-i2\pi x \xi} dx$$

- Exists both as *continuous* and *discrete* FT
- Today we will be looking at the former only



# Table of Contents

1 Introduction

2 Fourier Series

3 Fourier Transform



# Derivation of the Trigonometric Series

## Fourier Series

- For an even function:



# Derivation of the Trigonometric Series

## Fourier Series

- For an even function:

$$f_e(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right)$$





# Derivation of the Trigonometric Series

## Fourier Series

- For an even function:

$$f_e(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right)$$

$$f_e(x) \cos\left(\frac{\pi m x}{L}\right) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right)$$



# Derivation of the Trigonometric Series

## Fourier Series

- For an even function:

$$f_e(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right)$$

$$f_e(x) \cos\left(\frac{\pi m x}{L}\right) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right)$$

$$\int_{-L}^L f_e(x) \cos\left(\frac{\pi m x}{L}\right) dx = \int_{-L}^L \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right) dx$$



# Derivation of the Trigonometric Series

## Fourier Series

- For an even function:

$$f_e(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right)$$

$$f_e(x) \cos\left(\frac{\pi m x}{L}\right) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right)$$

$$\int_{-L}^L f_e(x) \cos\left(\frac{\pi m x}{L}\right) dx = \int_{-L}^L \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right) dx$$

$$= \sum_{n=0}^{\infty} a_n \int_{-L}^L \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right) dx$$



# Derivation of the Trigonometric Series

## Fourier Series

- For an even function:

$$f_e(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right)$$

$$f_e(x) \cos\left(\frac{\pi m x}{L}\right) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right)$$

$$\int_{-L}^L f_e(x) \cos\left(\frac{\pi m x}{L}\right) dx = \int_{-L}^L \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right) dx$$

$$= \sum_{n=0}^{\infty} a_n \int_{-L}^L \cos\left(\frac{\pi n x}{L}\right) \cos\left(\frac{\pi m x}{L}\right) dx$$

$$= \sum_{n=0}^{\infty} a_n \int_{-L}^L \frac{1}{2} \left( \cos\left((m+n)\frac{\pi x}{L}\right) + \cos\left((m-n)\frac{\pi x}{L}\right) \right) dx$$



- Consider the integral  $\int_{-L}^L \cos\left((m+n)\frac{\pi x}{L}\right) dx$

■

$$\frac{L}{m+n} \sin\left((m+n)\frac{\pi L}{L}\right) - \frac{L}{m+n} \sin\left((m+n)\frac{-\pi L}{L}\right)$$

■

$$= \frac{L}{m+n} (\sin((m+n)\pi) + \sin((m+n)\pi)) = 0$$

- And so we have:

$$\int_{-L}^L f_e(x) \cos\left(\frac{\pi m x}{L}\right) dx = \frac{1}{2} \sum_{n=0}^{\infty} a_n \int_{-L}^L \cos\left((m-n)\frac{\pi x}{L}\right) dx$$



# Another Aside

## Fourier Series

- For the reasons outlined last slide,  $\int_{-L}^L \cos\left(\frac{(m-n)\pi x}{L}\right) dx$  *usually* equals 0
- However, if  $m \equiv n$ :
- $\int_{-L}^L \cos\left(\frac{(n-n)\pi x}{L}\right) dx = \int_{-L}^L 1 dx = 2L$
- EVERY other value of  $m$  will yield a result of 0



# Conclusion

## Fourier Series

$$\begin{aligned}\int_{-L}^L f_e(x) \cos\left(\frac{\pi m x}{L}\right) dx &= \frac{1}{2} \sum_{n=0}^{\infty} \int_{-L}^L \cos\left(\frac{(m-n)\pi x}{L}\right) dx \\&= \frac{1}{2} (0 + 0 + \dots + 0 + 0 + 2La_m + 0 + 0 \dots) \\&= La_m \\a_m &= \frac{1}{L} \int_{-L}^L f_e(x) \cos\left(\frac{\pi m x}{L}\right) dx \\\therefore a_n &= \frac{1}{L} \int_{-L}^L f_e(x) \cos\left(\frac{\pi n x}{L}\right) dx\end{aligned}$$



# Special Case: $n \equiv 0$

## Fourier Series

$$f_e(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{\pi n x}{L}\right)$$

$$\int_{-L}^L f_e(x) dx = \sum_{n=0}^{\infty} a_n \int_{-L}^L \cos\left(\frac{\pi n x}{L}\right) dx$$

$$= 2La_0 + 0 + 0 + 0 \dots$$

$$\therefore a_0 = \frac{1}{2L} \int_{-L}^L f_e(x) dx$$





# Consolidation

## Fourier Series

- $b_n$  can be obtained just as easily, as:

$$b_n = \frac{1}{L} \int_{-L}^L f_o(x) \sin\left(\frac{\pi nx}{L}\right) dx \quad (b_0 = 0)$$

- A function neither even nor odd can be obtained via combination
- We can now express any periodic function as sines and cosines!<sup>1</sup>

■

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos\left(\frac{\pi nx}{L}\right) + b_n \sin\left(\frac{\pi nx}{L}\right) \right)$$

---

<sup>1</sup>Restrictions may apply



# Exponential Form!

## Fourier Series

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i \frac{\pi n x}{L}}$$



# Exponential Form!

## Fourier Series

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i \frac{\pi n x}{L}}$$

$$f(x) e^{-i \frac{\pi m x}{L}} = \sum_{n=-\infty}^{\infty} c_n e^{i \frac{\pi n x}{L}} \times e^{-i \frac{\pi m x}{L}}$$



# Exponential Form!

## Fourier Series

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i \frac{\pi n x}{L}}$$

$$f(x) e^{-i \frac{\pi m x}{L}} = \sum_{n=-\infty}^{\infty} c_n e^{i \frac{\pi n x}{L}} \times e^{-i \frac{\pi m x}{L}}$$

$$\int_{-L}^L f(x) e^{-i \frac{\pi m x}{L}} dx = \sum_{n=-\infty}^{\infty} c_n \int_{-L}^L e^{i(n-m) \frac{\pi x}{L}} dx$$



# An Aside (Look Familiar?)

## Fourier Series

$$\begin{aligned}(n \neq m) \quad \int_{-L}^L e^{i(n-m)\frac{\pi x}{L}} dx &= \int_{-L}^L \cos\left((n-m)\frac{\pi x}{L}\right) dx \\ &\quad + i \int_{-L}^L \sin\left((n-m)\frac{\pi x}{L}\right) dx \\ &= \frac{L}{\pi(n-m)} (\sin((n-m)\pi) - \sin(-(n-m)\pi)) \\ &\quad + i \frac{L}{\pi(n-m)} (\cos((n-m)\pi) - \cos(-(n-m)\pi)) = 0\end{aligned}$$

OR ( $n \equiv m$ )

$$\begin{aligned}&= \int_{-L}^L \cos(0) dx + i \int_{-L}^L \sin(0) dx \\ &= 2L\end{aligned}$$



# Conclusion

## Fourier Series

$$\int_{-L}^L f(x) e^{-i \frac{\pi m x}{L}} dx = 2L c_m$$

$$c_m = \frac{1}{2L} \int_{-L}^L f(x) e^{-i \frac{\pi m x}{L}} dx$$

$$\therefore c_n = \frac{1}{2L} \int_{-L}^L f(x) e^{-i \frac{\pi n x}{L}} dx$$



# Recap

## Fourier Series

$$f(x) = \frac{1}{2L} \int_{-L}^L f(x) dx + \sum_{n=1}^{\infty} \frac{1}{L} \int_{-L}^L \left( f(x) \cos \left( \frac{\pi n x}{L} \right) dx \right) \cos \left( \frac{\pi n x}{L} \right) \\ + \sum_{n=1}^{\infty} \frac{1}{L} \int_{-L}^L \left( f(x) \sin \left( \frac{\pi n x}{L} \right) dx \right) \sin \left( \frac{\pi n x}{L} \right)$$

$$f(x) = \sum_{n=-\infty}^{\infty} \frac{1}{2L} \int_{-L}^L \left( f(x) e^{-i \frac{\pi n x}{L}} dx \right) e^{i \frac{\pi n x}{L}}$$



# Table of Contents

1 Introduction

2 Fourier Series

3 Fourier Transform





# Setup

## Fourier Transform

- Let  $\xi = \frac{n}{2L}$  (The frequency of any term in the sequence), and extend  $L$  to  $\infty$ .
- Now,

$$c_n = \frac{1}{2L} \int_{-\infty}^{\infty} f(x) e^{-i2\pi x \xi} dx$$

- Just let  $2Lc_n = \hat{f}(\xi)$ , and that's our Fourier Transform!

$$\hat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-i2\pi x \xi} dx$$



# Inverse Fourier Transform

## Fourier Transform

- If  $\xi = \frac{n}{2L}$ , then let  $\Delta\xi = \frac{1}{2L}$

$$\begin{aligned}\lim_{L \rightarrow \infty} f(x) &= \lim_{L \rightarrow \infty} \sum_{n=-\infty}^{\infty} \frac{1}{2L} \hat{f}(\xi) e^{i2\pi x \xi} \\ &= \lim_{L \rightarrow \infty} \sum_{n=-\infty}^{\infty} \Delta\xi \hat{f}(\xi) e^{i2\pi x \xi}\end{aligned}$$

- But  $\lim_{L \rightarrow \infty} \frac{1}{2L} = 0$ , and  $\lim_{n \rightarrow \pm\infty} \frac{n}{2L} = \pm\infty$

$$f(x) = \lim_{\Delta\xi \rightarrow 0} \sum_{\xi=-\infty}^{\infty} \Delta\xi \hat{f}(\xi) e^{i2\pi x \xi}$$

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{i2\pi x \xi} d\xi$$



# Disclaimer

## Fourier Transform

- The Fourier Series only applies to *piecewise continuous* functions
- The Fourier Transform is usually impossible to apply properly to functions over an infinite domain
- (Unless you're willing to teach yourself about the Dirac  $\delta$  function)

