


Probability

Vincent Macri



© Vincent Macri, 2017



Table of Contents

1 Introduction

2 Basics

3 Math Elections

4 Generalizations

5 Yahtzee



Probability

Introduction

Probability lets us calculate how likely an event is to occur.



Probability

Introduction

Probability lets us calculate how likely an event is to occur.

Probability has implications in almost every field.



Probability

Introduction

Probability lets us calculate how likely an event is to occur.

Probability has implications in almost every field.

For example, in medicine, probability can tell you the likelihood of an operation being successful.



Probability

Introduction

Probability lets us calculate how likely an event is to occur.

Probability has implications in almost every field.

For example, in medicine, probability can tell you the likelihood of an operation being successful.

In engineering, knowing the probabilities of different components failing could help you troubleshoot problems faster.



Probability

Introduction

Probability lets us calculate how likely an event is to occur.

Probability has implications in almost every field.

For example, in medicine, probability can tell you the likelihood of an operation being successful.

In engineering, knowing the probabilities of different components failing could help you troubleshoot problems faster.

A lot of games have an element of chance. For example, knowing the probabilities of landing on different spaces in Monopoly will tell you which properties will make the most money.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.
- Sets are usually labelled with uppercase letters.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.
- Sets are usually labelled with uppercase letters.
- The elements of a set are separated by commas inside of braces. Example: $A = \{1, 2, 3\}$.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.
- Sets are usually labelled with uppercase letters.
- The elements of a set are separated by commas inside of braces. Example: $A = \{1, 2, 3\}$.
- A subset of a set contains only elements of the set it is a subset of. For example, a subset of A is $\{1, 2\}$.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.
- Sets are usually labelled with uppercase letters.
- The elements of a set are separated by commas inside of braces. Example: $A = \{1, 2, 3\}$.
- A subset of a set contains only elements of the set it is a subset of. For example, a subset of A is $\{1, 2\}$.
- The empty set (also called the null set) is the set containing no elements. It is written as $\{\}$ or \emptyset . It is a subset of all other sets.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.
- Sets are usually labelled with uppercase letters.
- The elements of a set are separated by commas inside of braces. Example: $A = \{1, 2, 3\}$.
- A subset of a set contains only elements of the set it is a subset of. For example, a subset of A is $\{1, 2\}$.
- The empty set (also called the null set) is the set containing no elements. It is written as $\{\}$ or \emptyset . It is a subset of all other sets.
- A set with k elements has 2^k subsets.



A quick overview of sets

Introduction

- Sets are an **unordered** collection of **distinct** elements.
- Sets are usually labelled with uppercase letters.
- The elements of a set are separated by commas inside of braces. Example: $A = \{1, 2, 3\}$.
- A subset of a set contains only elements of the set it is a subset of. For example, a subset of A is $\{1, 2\}$.
- The empty set (also called the null set) is the set containing no elements. It is written as $\{\}$ or \emptyset . It is a subset of all other sets.
- A set with k elements has 2^k subsets.
- The cardinality of a set A , written as $n(A)$ is the number of elements in A . For example, $n(A) = 3$.



A quick overview of set operations

Introduction

There are two main set operations we deal with.



A quick overview of set operations

Introduction

There are two main set operations we deal with.

Set Union $A \cup B$ is the set of all elements in A **or** B .



A quick overview of set operations

Introduction

There are two main set operations we deal with.

Set Union $A \cup B$ is the set of all elements in A **or** B .

Set Intersection $A \cap B$ is the set of all elements in A **and** B .



Probability definitions

Introduction

Sample space The set of all possible outcomes for an activity (experiment) is the sample space of the experiment. We will call this set S .



Probability definitions

Introduction

Sample space The set of all possible outcomes for an activity (experiment) is the sample space of the experiment. We will call this set S .

Event An event is a subset of the sample space. We will call this set E .



Table of Contents

1 Introduction

2 Basics

3 Math Elections

4 Generalizations

5 Yahtzee



Equal probability

Basics

If every possible outcome of an experiment has the same chance of occurring, then we can easily calculate the probability of an event (subset of the sample space) occurring.

The probability function, $p(E)$ gives us the probability of an event occurring.

$$p(E) = \frac{n(E)}{n(S)}$$

Probability is usually written as a number in the range $[0, 1]$.



Example

Basics

We are going to roll a six-sided die.

What is the sample space?



Example

Basics

We are going to roll a six-sided die.

What is the sample space?

$$S = \{1, 2, 3, 4, 5, 6\}$$



Example

Basics

We are going to roll a six-sided die.

What is the sample space?

$$S = \{1, 2, 3, 4, 5, 6\}$$

Let's define two events: we roll a composite number (C), or we roll a non-composite number (N).

$$C = \{4, 6\} \quad N = \{1, 2, 3, 5\}$$



Example

Basics

We are going to roll a six-sided die.

What is the sample space?

$$S = \{1, 2, 3, 4, 5, 6\}$$

Let's define two events: we roll a composite number (C), or we roll a non-composite number (N).

$$C = \{4, 6\} \quad N = \{1, 2, 3, 5\}$$

Then, the chance of rolling a composite number is:

$$p(C) = \frac{n(C)}{n(S)} = \frac{2}{6} = \frac{1}{3}$$



Example

Basics

We are going to roll a six-sided die.

What is the sample space?

$$S = \{1, 2, 3, 4, 5, 6\}$$

Let's define two events: we roll a composite number (C), or we roll a non-composite number (N).

$$C = \{4, 6\} \quad N = \{1, 2, 3, 5\}$$

Then, the chance of rolling a composite number is:

$$p(C) = \frac{n(C)}{n(S)} = \frac{2}{6} = \frac{1}{3}$$

And the chance of rolling a non-composite number is:



Example

Basics

We are going to roll a six-sided die.

What is the sample space?

$$S = \{1, 2, 3, 4, 5, 6\}$$

Let's define two events: we roll a composite number (C), or we roll a non-composite number (N).

$$C = \{4, 6\} \quad N = \{1, 2, 3, 5\}$$

Then, the chance of rolling a composite number is:

$$p(C) = \frac{n(C)}{n(S)} = \frac{2}{6} = \frac{1}{3}$$

And the chance of rolling a non-composite number is:

$$p(N) = \frac{n(N)}{n(S)} = \frac{4}{6} = \frac{2}{3}$$



Factorial

Basics

The factorial operation is the product of all integers from 1 to n .



Factorial

Basics

The factorial operation is the product of all integers from 1 to n .

Definition

$$n! = 1 \times 2 \times \cdots \times (n - 1) \times n$$

$0! = 1$, because otherwise you would break a lot of math.

Factorial is only defined for non-negative integers. However, using very advanced mathematics it is possible to calculate it for other values.



Factorial

Basics

The factorial operation is the product of all integers from 1 to n .

Definition

$$n! = 1 \times 2 \times \cdots \times (n - 1) \times n$$

$0! = 1$, because otherwise you would break a lot of math.

Factorial is only defined for non-negative integers. However, using very advanced mathematics it is possible to calculate it for other values.

Example

$$5! = 1 \times 2 \times 3 \times 4 \times 5 = 120$$



Table of Contents

1 Introduction

2 Basics

3 Math Elections

4 Generalizations

5 Yahtzee



Problem

Math Elections

The student body of Mackenzie is voting on their 3 favourite parts of math at the school.

They have 5 options: calculus, vectors, \LaTeX , Moodle, and math club.

Everyone votes for their favourite part, second favourite part, and third favourite part of math at the school.

These are three separate votes, taken one after another.



Ordered number of options

Math Elections

We reason that:



Ordered number of options

Math Elections

We reason that:

- 1 There are 5 choices for the first spot.



Ordered number of options

Math Elections

We reason that:

- 1 There are 5 choices for the first spot.
- 2 Whatever wins first cannot place second, so we are left with 4 choices for second place.



Ordered number of options

Math Elections

We reason that:

- 1 There are 5 choices for the first spot.
- 2 Whatever wins first cannot place second, so we are left with 4 choices for second place.
- 3 Whatever options placed first or second cannot place third, so we are left with 3 options for third place.



Ordered number of options

Math Elections

We reason that:

- 1 There are 5 choices for the first spot.
- 2 Whatever wins first cannot place second, so we are left with 4 choices for second place.
- 3 Whatever options placed first or second cannot place third, so we are left with 3 options for third place.

This means there are $5 \times 4 \times 3 = 60$ possible combinations.

We can develop the formula for this with factorials:

$$5 \times 4 \times 3 = 5 \times 4 \times 3 \times \frac{2!}{2!} = \frac{5 \times 4 \times 3 \times 2!}{2!} = \frac{5!}{2!} = 60$$



Unordered number of options

Math Elections

If we are only interested in what people's favourite parts of math at the school are, and not the ordering, we can instead calculate how many combinations of 3 elements there are, with no regard to the order.



Unordered number of options

Math Elections

If we are only interested in what people's favourite parts of math at the school are, and not the ordering, we can instead calculate how many combinations of 3 elements there are, with no regard to the order.

Take for example, these 6 outcomes:

First	Second	Third
Math club	Moodle	L ^A T _E X
Math club	L ^A T _E X	Moodle
Moodle	Math club	L ^A T _E X
Moodle	L ^A T _E X	Math club
L ^A T _E X	Math club	Moodle
L ^A T _E X	Moodle	Math club



Unordered number of options

Math Elections

If we are only interested in what people's favourite parts of math at the school are, and not the ordering, we can instead calculate how many combinations of 3 elements there are, with no regard to the order.

Take for example, these 6 outcomes:

First	Second	Third
Math club	Moodle	L ^A T _E X
Math club	L ^A T _E X	Moodle
Moodle	Math club	L ^A T _E X
Moodle	L ^A T _E X	Math club
L ^A T _E X	Math club	Moodle
L ^A T _E X	Moodle	Math club

If we do not care about ordering, these 6 options become 1 option.

This means that there are $\frac{60}{3!} = 10$ options for favourite parts of math at Mackenzie when we ignore the ordering.



Table of Contents

1 Introduction

2 Basics

3 Math Elections

4 Generalizations

5 Yahtzee



Number of ordered options

Generalizations

We can calculate the number of **ordered** subsets of size k from a set of cardinality n with the following formula:

$$P(n, k) = \frac{n!}{(n - k)!}$$



Number of ordered options

Generalizations

We can calculate the number of **ordered** subsets of size k from a set of cardinality n with the following formula:

$$P(n, k) = \frac{n!}{(n - k)!}$$

These ordered sets are called **permutations** of n things taken k at a time.



Number of unordered options

Generalizations

We can calculate the number of **unordered** subsets of size k from a set of cardinality n with the following formula:

$$C(n, k) = \frac{n!}{k!(n-k)!}$$

These unordered selections are called **combinations** of n things taken k at a time.



Number of unordered options

Generalizations

We can calculate the number of **unordered** subsets of size k from a set of cardinality n with the following formula:

$$C(n, k) = \frac{n!}{k!(n-k)!}$$

These unordered selections are called **combinations** of n things taken k at a time.

This formula is also written as:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

And is read as “ n choose k ”.



Table of Contents

1 Introduction

2 Basics

3 Math Elections

4 Generalizations

5 Yahtzee



Probability of rolling a Yahtzee

Yahtzee

Problem

Yahtzee, a game popular with my senior relatives, involves rolling 5 dice. In this game, you get a Yahtzee, which is worth a lot of points, if you roll 5 of a kind.

All of the dice have six sides.

Question

What is the probability of rolling a Yahtzee?



Calculations

Yahtzee

First, we will calculate the number of combinations of dice that can be rolled:



Calculations

Yahtzee

First, we will calculate the number of combinations of dice that can be rolled:

$$6^5 = 7776$$

We will call the set of possible rolls R . The cardinality of R is 7776.



Calculations

Yahtzee

First, we will calculate the number of combinations of dice that can be rolled:

$$6^5 = 7776$$

We will call the set of possible rolls R . The cardinality of R is 7776.

We know that the set of Yahtzee rolls is:

$$Y = \{(1, 1, 1, 1, 1), (2, 2, 2, 2, 2), (3, 3, 3, 3, 3), \\ (4, 4, 4, 4, 4), (5, 5, 5, 5, 5), (6, 6, 6, 6, 6)\}$$



Calculations

Yahtzee

First, we will calculate the number of combinations of dice that can be rolled:

$$6^5 = 7776$$

We will call the set of possible rolls R . The cardinality of R is 7776.

We know that the set of Yahtzee rolls is:

$$Y = \{(1, 1, 1, 1, 1), (2, 2, 2, 2, 2), (3, 3, 3, 3, 3), \\ (4, 4, 4, 4, 4), (5, 5, 5, 5, 5), (6, 6, 6, 6, 6)\}$$

$$p(Y) = \frac{n(Y)}{n(R)} = \frac{6}{7776} = \frac{1}{1296} \doteq 0.00077 = 0.077\%$$

\therefore the chance of rolling a Yahtzee is about 0.077%.



Rolling quadruples

Yahtzee

Question 2

How likely is it to roll four of a kind when rolling 5 dice?

We are looking at the chance of rolling **exactly** four of a kind. 5 of a kind does **not** count.



Quadruples solution

Yahtzee

Let's consider the case of rolling quadruple ones.



Quadruples solution

Yahtzee

Let's consider the case of rolling quadruple ones.

The chance this happening is with four dice is:

$$\left(\frac{1}{6}\right)^4 = \frac{1}{1296}$$



Quadruples solution

Yahtzee

Let's consider the case of rolling quadruple ones.

The chance this happening is with four dice is:

$$\left(\frac{1}{6}\right)^4 = \frac{1}{1296}$$

But we have five dice, so we must multiply this by the chance of rolling something other than one, and we must multiply by the number of different ways to arrange 4 ones and 1 other die:

$$\left(\frac{1}{6}\right)^4 \times \frac{5}{6} \times 5 = \frac{25}{7776}$$



Quadruples solution

Yahtzee

Let's consider the case of rolling quadruple ones.

The chance this happening is with four dice is:

$$\left(\frac{1}{6}\right)^4 = \frac{1}{1296}$$

But we have five dice, so we must multiply this by the chance of rolling something other than one, and we must multiply by the number of different ways to arrange 4 ones and 1 other die:

$$\left(\frac{1}{6}\right)^4 \times \frac{5}{6} \times 5 = \frac{25}{7776}$$

The chance of rolling quadruple ones is the same as rolling quadruple twos, threes, etc.. So:

$$\left(\frac{1}{6}\right)^4 \times \frac{5}{6} \times 5 \times 6 = \frac{25}{1296} \doteq 0.0193$$



Rolling triples

Yahtzee

Question 3

How likely is it to roll three of a kind when rolling 5 dice?

We are looking at the chance of rolling **exactly** three of a kind. 4 or 5 of a kind do **not** count. We **will** count a 2 of a kind and a 3 of a kind as a triple.



Triples solution setup

Yahtzee

This is similar to the last problem, so we will skip some steps for the sake of brevity.



Triples solution setup

Yahtzee

This is similar to the last problem, so we will skip some steps for the sake of brevity.

The main difference is that we don't know how many ways we can order three of a kind with two other numbers. With 4 of a kind, this was trivial. It is not longer trivial with three of a kind.

$$\left(\frac{1}{6}\right)^3 \times \left(\frac{5}{6}\right)^2 \times c \times 6$$

This is essentially the same formula we used last time, but replacing the number of combinations with the variable c . We must use reasoning to figure out c .



Triples solution

Yahtzee

$$\left(\frac{1}{6}\right)^3 \times \left(\frac{5}{6}\right)^2 \times c \times 6$$



Triples solution

Yahtzee

$$\left(\frac{1}{6}\right)^3 \times \left(\frac{5}{6}\right)^2 \times c \times 6$$

We want the number of combinations of 3 elements from a set of 5 elements.



Triples solution

Yahtzee

$$\left(\frac{1}{6}\right)^3 \times \left(\frac{5}{6}\right)^2 \times c \times 6$$

We want the number of combinations of 3 elements from a set of 5 elements.

$$c = \binom{5}{3} = 10$$



Triples solution

Yahtzee

$$\left(\frac{1}{6}\right)^3 \times \left(\frac{5}{6}\right)^2 \times c \times 6$$

We want the number of combinations of 3 elements from a set of 5 elements.

$$c = \binom{5}{3} = 10$$

$$\left(\frac{1}{6}\right)^3 \times \left(\frac{5}{6}\right)^2 \times 10 \times 6 = \frac{125}{648} \doteq 0.193$$

