

- 155.** Use the fact of algebra that a polynomial in  $(x_1, \dots, x_n)$ , which vanishes when  $x_i = x_j$ , is divisible by  $x_i - x_j$ .
- 156.** Divide the  $k$ th column by  $k$  and apply Vandermonde's identity.
- 158.**  $H : \mathbb{R}^7 \rightarrow \mathbb{R}^1$ .
- 160.** The space spanned by columns of  $A$  and  $B$  contains columns of  $A+B$ .
- 166.**  $(A^t \mathbf{a})(\mathbf{x}) = \mathbf{a}(A\mathbf{x})$  for all  $\mathbf{x} \in \mathbb{K}^n$  and  $\mathbf{a} \in (\mathbb{K}^m)^*$ .
- 180.** Modify the Gaussian elimination algorithm of Section 2 by permuting unknowns (instead of equations).
- 181.** Apply the  $LUP$  decomposition to  $M^t$ .
- 182.** Apply  $LPU$ ,  $LUP$ , and  $PLU$  decompositions to  $M^{-1}$ .
- 189.** Consider the map  $\mathbb{R}^n \rightarrow \mathbb{R}^{p+q}$  defined by the linear forms.
- 202.**  $T = A + iB$  where both  $A$  and  $B$  are Hermitian.
- 203.**  $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{z}^\dagger \mathbf{w}$ .
- 205.**  $\langle \mathbf{z}, \mathbf{w} \rangle = \mathbf{z}^\dagger \mathbf{w}$ .
- 206.** Use the previous exercise.
- 208.** The normal forms are:  $i|Z_1|^2 - i|Z_2|^2$ ,  $|Z_1|^2 - |Z_2|^2$ ,  $|Z_1|^2$ .
- 213.** Take the linear form for one of new coordinates.
- 235.** Apply Vieta's formula.
- 239.** Find an eigenvector, and show that its orthogonal complement is invariant.
- 241.** Compute  $\langle \mathbf{x}, A\mathbf{x} \rangle$  for  $\mathbf{x} = \sum t_i \mathbf{v}_i$ , where  $\{\mathbf{v}_i\}$  is an orthonormal basis of eigenvectors, and  $\sum t_i^2 = 1$ .
- 242.** If  $AB = BA$ , then eigenspaces of  $A$  are  $B$ -invariant.
- 247.** Construct  $v_1, \dots, v_n$  by applying the Spectral Theorem to the non-negative Hermitian operator  $A^\dagger A$ , and check that  $A\mathbf{v}_i \in \mathcal{W}$  corresponding to distinct nonzero eigenvalues  $\mu_i$  of  $A^\dagger A$  are pairwise perpendicular.
- 248.** Rephrase the previous exercise in matrix form.
- 251.** The equation  $\langle \mathbf{x} + t\mathbf{y}, \mathbf{x} + t\mathbf{y} \rangle = 0$  quadratic in  $t$  has no real solutions unless  $\mathbf{x}$  is proportional to  $\mathbf{y}$ .
- 253.**  $\langle \mathbf{x}, \mathbf{x} \rangle = (\sum x_i)^2 + \sum x_i^2$ .
- 255.**  $\det(U^t U) = 1$ .
- 264.** See Example 1.
- 274.** Use the previous exercise.
- 276.** Prove first that the radius of the circle must be equal to the middle semiaxis of the ellipsoid.
- 279.** For "exactly onne" claim, use the coordinate-less description of the spectrum provided by the minimax principle.
- 280.** When  $x \approx 0$ ,  $\sin x \approx x$ .

**283.** If  $T$  denotes the cyclic shift of coordinates in  $\mathbb{C}^n$ , then  $C = C_0I + C_1T + \cdots + C_{n-1}T^{n-1}$ .

**297.** Use Newton's binomial formula.

**310.** Pick  $A$  and  $B$  so that  $e^A e^B \neq e^B e^A$ .

**313.** Verify it first for a Jordan cell.

**318.** Make two of the lines the coordinate axes on the plane, and the third one the graph of an isomorphism between them.

**319.** If the first three lines are  $y = 0$ ,  $x = 0$ , and  $y = x$ , then the fourth one is  $y = \lambda x$ .

**320.** If  $d = 0$ ,  $a = \infty$ ,  $b = 1$ , then  $\lambda = c$ .

**322.** Assuming that the spaces  $\mathcal{U}_i$  have the same dimension, and the maps between them are invertible, consider the determinant (or eigenvalues) of the composition of the maps around the cycle.

**325.** Use the theory of Bruhat cells.

**326.** Use Lemma, and apply induction on the length of the shortest of the legs.

**327.** Use Sylvester's rule together with the previous exercise.

**330.** On the unit sphere of area  $4\pi$ , the area of a triangle with the angles  $(\frac{\pi}{p}, \frac{\pi}{q}, \frac{\pi}{r})$  has the area  $\frac{\pi}{p} + \frac{\pi}{q} + \frac{\pi}{r} - \pi$ . Use this to compute the number of required triangles.

**332.** Compute  $R_{\mathbf{v}_1} R_{\mathbf{v}_2}$  to show that it is a Jordan cell of size 2 with the eigenvalue 1.

**333.** They correspond to positive roots  $\mathbf{e}_i - \mathbf{e}_j = \mathbf{v}_{i+1} + \cdots + \mathbf{v}_j$ , where  $0 \leq i < j \leq n$ .

**344.**  $\overrightarrow{AA'} = 3\overrightarrow{AM}/2$ .

**350.**  $\cos^2 \theta \leq 1$ .

**355.** Rotate the regular  $n$ -gon through  $2\pi/n$ .

**358.** Take  $X = A$  first.

**359.**  $\overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA}$ .

**360.** If  $a + b + c + d = 0$ , then  $2(b + c)(c + d) = a^2 - b^2 + c^2 - d^2$ .

**361.**  $XA^2 = |\overrightarrow{OX} - \overrightarrow{OA}|^2 = 2R^2 - 2\langle \overrightarrow{OX}, \overrightarrow{OA} \rangle$  if  $O$  is the center.

**362.** Consider projections of the vertices to any line.

**363.** Project faces to an arbitrary plane and compute signed areas.

**372.** Compute  $(\cos \theta + i \sin \theta)^3$ .

**374.** Divide  $P$  by  $z - z_0$  with a remainder.

**377.**  $(z - z_0)(z - \bar{z}_0)$  is real.

**378.** Apply Vieta's formula.

**382.**  $e^{\pm i\theta} = \cos \theta \pm i \sin \theta$ .

**383.** See Figure spl2-5.

**387.** Show first that  $\binom{n}{k}$  equals the number of  $k$ -element subsets in the set of  $n$  objects.

**393.** Examine the kernel of the map  $\mathbb{R}[x] \rightarrow \mathbb{R}^2$  associating to a polynomial  $P$  the pair  $(P(1), P(-1))$  of its values at  $x = \pm 1$ .

**397.** To a linear form  $f : \mathcal{V}/\mathcal{W} \rightarrow \mathbb{K}$ , associate  $\mathcal{V} \xrightarrow{\pi} \mathcal{V}/\mathcal{W} \xrightarrow{f} \mathbb{K}$ .

**399.** Given  $B : \mathcal{V} \rightarrow \mathcal{W}^*$ , show that evaluation  $(B\mathbf{v})(\mathbf{w})$  of the linear form  $B\mathbf{v} \in \mathcal{W}^*$  on  $\mathbf{w} \in \mathcal{W}$  defines a bilinear form.

**404.** Compute  $q_1 q_2 q_3^* q_1^*$ .

**407.** Introduce multiplication by scalars  $q \in \mathbb{H}$  as  $A \mapsto q^* A$ .

**409.**  $|q|^2 = |-1| = 1$ .

**411.** Prove that  $-1$  is a non-square.

**414.** There are 1 even and 3 odd non-degenerate forms.

**418.**  $\frac{1}{3} = 1 - \frac{2}{3}$ .

**419.** Start with inverting *2-adic units*  $\cdots * * * 1$ . ( $*$  is a wild card).

**421.** Use the previous exercise.



# Answers

7. For normal forms,  $X^2$  and 0 can be taken.
12.  $(n+1)(n+2)/2$ .
15.  $x(t) = x(0)e^{\lambda t}$ .
16.  $x(t) = x(0)e^{3t}, y(t) = y(0)e^{-t}, z(t) = z(0)$ .
19.  $\dot{X}_1 = iX_1, \quad \dot{X}_2 = -iX_2$ .
22. Semiaxes  $2/\sqrt{3}$  and 2.
25. (a)  $\pm(\sqrt{\alpha^2 - \beta^2}, 0)$ , (b)  $\pm(\sqrt{\alpha^2 + \beta^2}, 0)$ .
29. Yes,  $k(x^2 + y^2)$ .
30.  $y = \pm x$ ; level curves of (a) are ellipses, of (c) hyperbolas.
31.  $2X^2 - Y^2 = 1$ .
34. 2nd ellipse, 1st & 4th hyperbolas.
38. A pair of intersecting lines,  $x - 1 = \pm(2y - 1)$ .
45. Yes, some non-equivalent equations represent the empty set.
50.  $\frac{n(n+1)}{2}$ .
65. Points, lines, and planes.
74.  $\dim = 2$ .
76.  $x^k = x_1^k L_1(x) + \cdots + x_n^k L_n(x), k = 0, \dots, n-1$ .
80. yes, no, no, yes, no, yes, yes, no.
82.  $E = \mathbf{va}$ , i.e.  $e_{ij} = v_i a_j, i = 1, \dots, m, j = 1, \dots, n$ .
83.  $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ .
84. Check your answers using associativity, e.g.  $(CB)A = C(BA)$ .

Other answers:  $BAC = \begin{bmatrix} 3 & 1 \\ 3 & 3 \end{bmatrix}, \quad ACB = \begin{bmatrix} 0 & -2 \\ 3 & 6 \end{bmatrix}.$

91.  $\begin{bmatrix} \cos 1^\circ & -\sin 1^\circ \\ \sin 1^\circ & \cos 1^\circ \end{bmatrix}.$

- 92.** If  $B = A^k$ , then  $b_{i,i+k} = 1$ , and all other  $b_{ij} = 0$ .
- 93.** The answer is the same as in the previous problem.
- 94.** (a) If  $A$  is  $m \times n$ ,  $B$  must be  $n \times m$ . (b) Both must be  $n \times n$ .
- 96.** Exactly when  $AB = BA$ .
- 103.** No; yes (of  $x, y \in \mathbb{R}^1$ ); yes (in  $\mathbb{R}^2$ ).
- 104.** I.
- 106.**  $S = 2x_1y_1 + x_1y_2 + x_2y_1$ ,  $A = x_1y_2 - x_2y_1$ .
- 107.**  $(\sum x_i)(\sum y_i)$  and  $\sum_{i \neq j} x_i y_j / 2$ .
- 108.** No, only if  $AB = BA$ .
- 112.**  $T^\dagger = \overline{z_2} w_1$ ;  $S_1 = (\overline{z_1} w_2 + \overline{z_2} w_1) / 2$ ,  $S_2 = (\overline{z_1} w_2 - \overline{z_2} w_1) / 2i$ ;  $H_1 = (\overline{z_1} z_2 + \overline{z_2} z_1) / 2$ ,  $H_2 = (\overline{z_1} z_2 - \overline{z_2} z_1) / 2i$ .
- 113.**  $(\overline{z_1} w_2 + \cdots + \overline{z_{n-1}} w_n + \overline{z_2} w_1 + \cdots + \overline{z_n} z_{n-1}) / 2$ .
- 115.** No, yes,  $n^2$ .
- 118.** 1; 1;  $\cos(\mathbf{x} + y)$ .
- 119.** 2, 1; -2.
- 121.**  $k(k-1)/2$ .
- 123.**  $n(n-1)/2 - l$ .
- 127.** Transpositions  $\tau^{(i)}$  of nearby indices  $(i, i+1)$ ,  $i = 1, \dots, n-1$ .
- 129.** E.g.  $\tau_{14}\tau_{34}\tau_{25}$ .
- 130.** No; yes.
- 131.** + (6 inverted pairs); + (8 inverted pairs).
- 132.** -1 522 200; -29 400 000.
- 134.** 0.
- 135.** Changes sign, if  $n = 4k + 2$  for some  $k$ , and remains unchanged otherwise.
- 136.**  $x = a_1, \dots, a_n$ .
- 138.** Leave it unchanged.
- 141.** (a)  $(-1)^{n(n-1)/2} a_1 a_2 \cdots a_n$ , (b)  $(ad - bc)(eh - fg)$ .
- 142.** (a) 9, (b) 5.
- 143.** (a)  $\begin{bmatrix} -\frac{5}{18} & -\frac{7}{18} & \frac{1}{18} \\ \frac{1}{18} & -\frac{5}{18} & \frac{7}{18} \\ \frac{7}{18} & \frac{1}{18} & -\frac{5}{18} \end{bmatrix}$ , (b)  $\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$ .
- 144.** (a)  $\mathbf{x} = [-\frac{2}{9}, \frac{1}{3}, \frac{1}{9}]^t$ , (b)  $\mathbf{x} = [1, -1, 1]^t$ .
- 146.**  $(\det A)^{n-1}$ , where  $n$  is the size of  $A$ .
- 147.** Those with determinants  $\pm 1$ .
- 148.** (a)  $x_1 = 3, x_2 = x_3 = 1$ , (b)  $x_1 = 3, x_2 = 4, x_3 = 5$ .

**149.** (a)  $-x_1^2 - \cdots - x_n^2$ , (b)  $(ad - bc)^3$ .

**153.**  $\lambda^n + a_1\lambda^{n-1} + \cdots + a_{n-1}\lambda + a_n$ .

**154.** 1.

**156.**  $1!3!5!\cdots(2n-1)!$ .

**158.** 1.

**161.** The system is consistent whenever  $b_1 + b_2 + b_3 = 0$ .

**165.** (a) Yes, (b) yes, (c) yes, (d) no, (e) yes, (f) no, (g) no, (h) yes.

**169.**  $0 \leq \text{codim} \leq k$ .

**171.** Two subspaces are equivalent if and only if they have the same dimension.

**172.** The equivalence class of an ordered pair  $\mathcal{U}, \mathcal{V}$  of subspaces is determined by  $k := \dim U$ ,  $l := \dim V$ , and  $r := \dim(\mathcal{U} + \mathcal{V})$ , where  $k, l \leq r \leq n$  can be arbitrary.

**174.**  $x_1 = 3, x_2 = 1, x_3 = 1$ ; inconsistent;  $x_1 = 1, x_2 = 2, x_3 = -2$ ;

$x_1 = 2t_1 - t_2, x_2 = t_1, x_3 = t_2, x_4 = 1$ ;

$x_1 = -8, x_2 = 3 + t, x_3 = 6 + 2t, x_4 = t$ ;  $x_1 = x_2 = x_3 = x_4 = 0$ ;

$x_1 = \frac{3}{17}t_1 - \frac{13}{17}t_2, x_2 = \frac{19}{17}t_1 - \frac{20}{17}t_2, x_3 = t_1, x_4 = t_2$ ;

$x_1 = -16 + t_1 + t_2 + 5t_3, x_2 = 23 - 2t_1 - 2t_2 - 6t_3, x_3 = t_1, x_4 = t_2, x_5 = t_3$ .

**175.**  $\lambda = 5$ .

**176.** (a)  $\text{rk} = 2$ , (b)  $\text{rk} = 2$ , (c)  $\text{rk} = 3$ , (d)  $\text{rk} = 4$ , (e)  $\text{rk} = 2$ .

**177.** Inverse matrices are:

$$\begin{bmatrix} 1 & -4 & -3 \\ 1 & -5 & -3 \\ -1 & 6 & 4 \end{bmatrix}, \quad \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}, \quad \begin{bmatrix} 2 & -1 & 0 & 0 \\ -3 & 2 & 0 & 0 \\ 31 & -19 & 3 & -4 \\ -23 & 14 & -2 & 3 \end{bmatrix}.$$

**185.**  $\binom{n}{2} - l(\sigma)$ , where  $l(\sigma)$  is the length of the permutation.

**186.** Inertia indices  $(p, q) = (1, 1), (2, 0), (1, 2)$ .

**191.** Empty for  $p = 0$ , has 2 components for  $p = 1$ , and 1 for  $p = 2, 3, 4$ .

**195.**  $z_1^2 + z_2^2 = 1, z_1^2 + z_2^2 = 0, z_2 = z_1^2, z_1^2 = 1, z_1^2 = 0$ .

**196.** Two parallel lines.

**201.** Those all of whose entries are imaginary.

**202.**  $T(\mathbf{z}, \mathbf{w}) =$

$$\frac{1}{2} [T(\mathbf{z} + \mathbf{w}, \mathbf{z} + \mathbf{w}) + iT(i\mathbf{z} + \mathbf{w}, i\mathbf{z} + \mathbf{w}) - (1 + i)(T(\mathbf{z}, \mathbf{z}) + T(\mathbf{w}, \mathbf{w}))].$$

**209.**  $d_{ii} = \Delta_i / \Delta_{i-1}$ .

**210.**  $(p, q) = (2, 2), (3, 1)$ .

**214.** No.

**215.** The sum of the squares of all sides of a parallelogram is equal to the sum of the squares of the diagonals.

**228.** id.

**252.**  $\theta(\mathbf{e}_i, \mathbf{e}_{i+1}) = 2\pi/3$ , while all other pairs are perpendicular.

**254.** One can take  $\mathbf{f}_i = \mathbf{e}_i - \mathbf{e}_0$ , and  $\mathbf{h}_i = \mathbf{e}_i - \mathbf{e}_{i-1}$ , where  $i = 1, \dots, n$ .

**256.** Rotations about the origin, and reflections about a line passing through the origin, which have determinants 1 and  $-1$  respectively.

**262.**  $\pm 1$ .

**269.** The operator is  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ .

**295.** (a)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}$  (b)  $\begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  (c)  $\begin{bmatrix} -3 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$   
 (e)  $\begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$  (f)  $\begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$  (k)  $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$  (l)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & -i \end{bmatrix}$ .

**306.**  $(a_0 - n(a_0 - a_1/3))3^n$ .

**307.** (a)  $(1 + \sqrt{10})/3$ .

**310.**  $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$  will do.

**317.** Pairs of  $m \times n$ -matrices; pairs of matrices of transposed sizes.

**321.**  $\mathbb{K}^1$ .

**323.** The restriction to the eigenline.

**328.** (a)  $(2, 2, r)$ ,  $(2, 3, 3)$ ,  $(2, 3, 4)$ ,  $(2, 3, 5)$ ; (b)  $(3, 3, 3)$ ,  $(2, 4, 4)$ ,  $(2, 3, 6)$ .

**330.** Symmetry planes of regular polyhedron (tetrahedron in case (b), octahedron or cube in case (c), and icosahedron or dodecahedron in case (d)) partition the surface of the sphere circumscribed around the polyhedron into (respectively, 24, 48, and 120) required spherical triangles.

**331.** Since  $\langle \mathbf{v}, \mathbf{v} \rangle = 2$ , we have:

$$\langle \mathbf{x} - \langle \mathbf{x}, \mathbf{v} \rangle \mathbf{v}, \mathbf{y} - \langle \mathbf{y}, \mathbf{v} \rangle \mathbf{v} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle - \langle \mathbf{x}, \mathbf{v} \rangle \langle \mathbf{v}, \mathbf{y} \rangle - \langle \mathbf{y}, \mathbf{v} \rangle \langle \mathbf{x}, \mathbf{v} \rangle + 2 \langle \mathbf{x}, \mathbf{v} \rangle \langle \mathbf{y}, \mathbf{v} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle.$$

**333.**  $\mathcal{V}^{i,j} : \dots \rightarrow 0 \rightarrow \mathbb{K}^1 \xrightarrow{\sim} \dots \xrightarrow{\sim} \mathbb{K}^1 \rightarrow 0 \rightarrow \dots$  ( $i$  zeroes followed by  $j - i$  copies of  $\mathbb{K}^1$ , followed by  $n - j$  zeroes).

**334.**  $\mathcal{V}^{0,n} \oplus \mathcal{V}^{1,n} \oplus \dots \oplus \mathcal{V}^{n-1,n}$ .

**335.**  $X(t) = e^{tA}X(0)e^{-tA}$ .

**336.** The eigenvalues are  $\lambda_i - \lambda_j$ ,  $i, j = 1, \dots, n$ . In the space of linear forms, they form the root system of type  $A_{n-1}$ . The eigenvectors are the elementary matrices  $E_{ij}$  (with the entry 1 at the  $i$ th row and  $j$ th column, and 0 everywhere else). Upper-triangular  $E_{ij}$  correspond to positive roots, and lower-triangular negative.

**337.**  $mg/2$ ,  $mg\sqrt{3}/2$ .



**338.** 18 min (reloading time excluded).

**342.** It rotates with the same angular velocity along a circle centered at the barycenter of the triangle formed by the centers of the given circles.

**346.**  $3/4$ .

**349.**  $7\overrightarrow{OA} = \overrightarrow{OA'} + 2\overrightarrow{OB'} + 4\overrightarrow{OC'}$  for any  $O$ .

**351.**  $3/2$ .

**353.**  $2\langle \mathbf{u}, \mathbf{v} \rangle = |\mathbf{u} + \mathbf{v}|^2 - |\mathbf{u}|^2 - |\mathbf{v}|^2$ .

**354.** (b) No.

**364.** Yes; Yes (0); Yes.

**365.** (a)  $\frac{1+5i}{13}$ ; (b)  $\frac{1-i\sqrt{3}}{2}$ .

**367.**  $|z|^{-2}$ .

**368.**  $\pm i\sqrt{3}$ .

**370.** (a)  $\sqrt{2}, -\pi/4$ ; (b)  $2, -\pi/3$ .

**371.**  $\frac{-1+i\sqrt{3}}{2}$ .

**375.**  $2 \pm i$ ;  $i\frac{1 \pm \sqrt{5}}{2}$ ;  $1 + i$ ,  $1 + i$ ;  $1 \pm \frac{\sqrt{6} - i\sqrt{2}}{2}$ .

**376.**  $-2, 1 \pm i\sqrt{3}$ ;  $i, \frac{\pm\sqrt{3}-i}{2}$ ;  $\pm i\sqrt{2}, \pm i\sqrt{2}$ ;  $\frac{\sqrt{3}\pm i}{2}, \frac{\sqrt{3}\pm i}{2}$ ;  $\pm i, \pm \frac{\sqrt{3}\pm i}{2}$ .

**379.**  $a_k = (-1)^k \sum_{1 \leq i_1 < \dots < i_k \leq n} z_{i_1} \cdots z_{i_k}$ .

**382.**  $\cos \theta = (e^{i\theta} + e^{-i\theta})/2$ ,  $\sin \theta = (e^{i\theta} - e^{-i\theta})/2i$ .

**384.** The upper half-plane  $\operatorname{Im} w > 0$ .

**386.**  $\cos 3\theta = 4\cos^3 \theta - 3\cos \theta$ ,  $\sin 3\theta = 3\sin \theta - 4\sin^3 \theta$ .

**389.**  $p^n$ ;  $p^{mn}$ .

**390.**  $p^{n(n-1)/2}$ .

**392.**  $\operatorname{Ker} D = \mathbb{K}[x^p] \subset \mathbb{K}[x]$ .

**394.** (a)  $n$ ; (b)  $n(n+1)/2$ .

**395.** 2, if  $n > 1$ .

**401.**  $\operatorname{Ker} A^t = \mathcal{V}(\mathcal{V})^\perp$ ,  $A^t(\mathcal{W}^*) = (\mathcal{V}/\operatorname{Ker} A)^* \subset \mathcal{V}^*$ .

**403.**  $-1$ ;  $-q$ .

**408.**  $\begin{bmatrix} z & -\bar{w} \\ w & \bar{z} \end{bmatrix}$ .

**409.**  $\{bi + cj + dk \mid b^2 + c^2 + d^2 = 1\}$ .



# Bibliography

- [1]
- [2] D. K. Faddeev, I. S. Sominsky. *Problems in Linear Algebra*. 8th edition, Nauka, Moscow, 1964 (in Russian).
- [3] A. P. Kiselev. *Kiselev's Geometry / Book I. Planimetry*. Adapted from Russian by Alexander Givental. Sumizdat, El Cerrito, 2006.
- [4] A. P. Kiselev. *Kiselev's Geometry / Book II. Stereometry*. Adapted from Russian by Alexander Givental. Sumizdat, El Cerrito, 2008.
- [5] J.-P. Serre. *A Course in Arithmetic*.
- [6] Ronald Solomon. *Abstract Algebra*. Thomson — Brooks/Cole, Belmont, CA, 2002.
- [7] Berezin
- [8] Van der Waerden
- [9] Hermann Weyl. *Space. Time. Matter*.
- [10] P. Gabriel. *Unzerlegbare Darstellungen, I*. Manuscripta Mathematica 6 (1972), 71–103.
- [11] I. N. Bernstein, I. M. Gelfand, V. A. Ponomarev. *Coxeter functors, and Gabriel's theorem*. Russian mathematical surveys 28, No. 2, (1973), 1732.

# Index

- $n$ -space, 23
- $n$ -vectors, 23
  
- absolute value, 174, 191
- addition of vectors, 21
- additivity, 169
- adjoint form, 42
- adjoint map, 38, 67, 190
- adjoint matrix, 56
- adjoint quaternion, 190
- adjoint systems, 71
- affine subspace, 25, 187
- algebraic number, 182
- algebraically closed field, 133
- annihilator, 189
- anti-Hermitian form, 43, 93
- anti-Hermitian quadratic form, 93
- anti-linear, 42
- anti-symmetric bilinear form, 40
- anti-symmetric matrix, 41
- anti-symmetric operator, 114
- argument of complex number, 175
- associative, 36
- associativity, 166
- augmented matrix, 74
- axiom, 21
- axis of symmetry, 15
  
- Bézout, 180
- Bézout's theorem, 180
- back substitution, 74
- barycenter, 171
- basis, 26, 167
- bijjective, 24, 185
- bilinear form, 39
- bilinearity, 169
- Binet, 60
- Binet–Cauchy formula, 60
  
- block, 51
- block triangular matrix, 52
- Bruhat cell, 84
  
- canonical form, 17
- canonical isomorphism, 30
- canonical projection, 187
- Cartesian coordinates, 170
- category of vector spaces, 23, 184
- Cauchy, 60
- Cauchy – Schwarz inequality, 99, 172
- Cauchy's interlacing theorem, 118
- Cayley transform, 108
- characteristic equation, 103
- characteristic polynomial, 7, 127
- Chebyshev, 30
- Chebyshev polynomials, 30
- classification theorem, 17
- codimension, 69
- cofactor, 55
- cofactor expansion, 55, 58
- column, 183
- column space, 76
- commutative square, 67, 153
- commutativity, 166
- commutator, 97
- complementary multi-index, 58
- complete flag, 82
- completing squares, 12
- complex conjugate, 173
- complex conjugation, 112
- complex multiplication, 116
- complex sphere, 92
- complex vector space, 22
- complexification, 112
- components, 183
- composition, 65
- conic section, 9

- conics, 90
- conjugate quaternion, 190
- connected graph, 151
- continuous time, 139
- coordinate Euclidean space, 111
- coordinate flag, 82
- coordinate quaternionic space, 191
- coordinate system, 167
- coordinate vectors, 23, 183
- coordinates, 23, 27, 167, 183
- Cramer's rule, 57
- cross product, 64
- cross-ratio, 163
- cylinder, 90
  
- Dandelin, 10
- Dandelin's spheres, 9
- Darboux, 197
- Darboux basis, 197
- decomposable representation, 154
- degenerate bilinear form, 125
- Descartes, 170
- determinant, 19, 47
- diagonal matrix, 184
- diagonalizable matrix, 133
- differentiation, 185
- dimension, 27
- dimension of Bruhat cell, 85
- direct sum, 4, 24, 154, 186
- direct sum of representations, 154
- directed segment, 165
- directrix, 16
- discrete dynamical systems, 139
- discrete time, 139
- discriminant, 176, 200
- distance, 99
- distributive law, 34, 174
- distributivity, 166
- dot product, 169
- dot-product, 39
- dual basis, 29, 193
- dual map, 38, 190
- dual space, 25, 185
  
- eccentricity, 16
- edge, 151
- eigenspace, 103, 127
- eigenvalue, 5, 102, 127
- eigenvector, 102, 127
- elementary product, 47
- elementary row operations, 73
- ellipse, 10
- ellipsoid, 119
- entry, 183
- equivalent, 17
- equivalent conics, 91
- equivalent linear maps, 67
- equivalent representations, 152
- Euclidean inner product, 111
- Euclidean space, 111
- Euclidean structure, 111
- Euler's formula, 179
- evaluation, 34
- evaluation map, 185
- even form, 195
- even permutation, 49
- exponential function, 178
  
- Fibonacci, 139
- Fibonacci sequence, 139
- field, 22, 174, 182
- field of  $p$ -adic numbers, 199
- finite dimensional spaces, 27
- flag, 82
- focus of ellipse, 10
- focus of hyperbola, 15
- focus of parabola, 16
- Fourier, 122
- Fourier basis, 122
- Fundamental Formula of Mathematics, 180
  
- Gabriel, 155
- Gaussian elimination, 73
- Gelfand's problem, 125
- golden ratio, 140
- Gram matrix, 107
- Gram-Schmidt process, 96, 100
- graph, 24, 151, 187
- gravitation constant, 126
- group, 161
  
- half-linear, 42
- Hamilton-Cayley equation, 136
- harmonic oscillator, 121
- Hasse, 199

- head, 165
- Hermite, 42
- Hermite polynomials, 30
- Hermitian adjoint, 101
- Hermitian adjoint form, 42
- Hermitian conjugate matrix, 42
- Hermitian form, 43
- Hermitian inner product, 99
- Hermitian isomorphic, 100
- Hermitian operator, 102
- Hermitian quadratic form, 43, 93
- Hermitian space., 99
- Hermitian-anti-symmetric form, 42
- Hermitian-symmetric form, 42
- Hilbert space, 99
- Hirani, 16
- homogeneity, 169
- homogeneous system, 69
- homomorphism theorem, 188
- hyperbola, 10
- hyperplane, 72
- hypersurface, 90
  
- identity matrix, 37
- identity permutation, 49
- imaginary part, 173
- imaginary unit, 173
- inconsistent system, 75
- indecomposable representation, 154
- indices in inversion, 48
- induction hypothesis, 28
- inertia index, 89
- injective, 24, 184
- inner product, 169
- invariant subspace, 103
- inverse matrix, 37
- inverse quaternion, 191
- inverse transformation, 37
- inversion of indices, 48
- involution, 113
- irreducible representation, 155
- isometric Hermitian spaces, 100
- isomorphic spaces, 24, 185
- isomorphism, 24, 185
- iterations of linear maps, 139
  
- Jordan block, 7
- Jordan canonical form, 132
- Jordan cell, 132
- Jordan normal form, 132
- Jordan system, 7
  
- kernel, 24, 184
- kernel of form, 90, 97
- kinetic energy, 120
  
- Lagrange, 58
- Lagrange polynomials, 29
- Lagrange's formula, 58
- law of cosines, 170
- LDU decomposition, 81
- leading coefficient, 74
- leading entry, 74
- leading minors, 95
- left inverse, 71
- left kernel, 189
- left vector space, 191
- length, 99
- length of permutation, 48
- linear combination, 166
- linear dynamical systems, 139
- linear form, 25, 33, 185
- linear function, 25, 33, 185
- linear map, 24, 184
- linear ODE, 139
- linear recursion relation, 139
- linear subspace, 23, 183
- linear transformation, 37
- linearly dependent, 27
- linearly independent, 26
- Linnaeus, 17
- lower triangular, 79
- lower-triangular matrix, 184
- LPU decomposition, 79
- LU decomposition, 81
- LUP decomposition, 85
  
- Möbius band, 30
- mathematical induction, 28
- matrix, 33, 183
- matrix entry, 33
- matrix product, 34, 35
- metric space, 99
- Minkowski, 199
- Minkowski–Hasse theorem, 199
- minor, 55, 58

- multi-index, 58
- multiplication by scalar, 166
- multiplication by scalars, 21
- multiplicative, 174
- multiplicity, 176
  
- nilpotent operator, 129
- non-degenerate bilinear form, 125
- non-degenerate Hermitian form, 95
- nontrivial linear combination, 27
- norm of quaternion, 191
- normal form, 17
- normal operator, 101, 114
- null space, 24, 76, 184
  
- odd form, 195
- odd permutation, 49
- ODE, 139
- operator, 102
- opposite coordinate flag, 83
- opposite vector, 21
- oriented graph, 151
- orthogonal, 100
- orthogonal basis, 87
- orthogonal complement, 104
- orthogonal diagonalization, 107
- Orthogonal Diagonalization Theorem, 117
- orthogonal projection, 107
- orthogonal projector, 108
- orthogonal transformation, 111
- orthogonal vectors, 170
- orthonormal basis, 96, 100, 111
  
- parabola, 14
- partition, 131
- Pascal's triangle, 202
- pendulum, 126
- permutation, 47
- permutation matrix, 79
- phase plane, 121
- pivot, 74
- Plücker, 63
- Plücker coordinates, 72
- Plücker identity, 63
- PLU decomposition, 85
- polar, 175
- polar decomposition, 109
  
- positive definite, 88
- positive operator, 109
- positivity, 169
- potential energy, 120
- power of matrix, 44
- principal axes, 15, 119
- principal minor, 97
- projection, 100
- Pythagorean theorem, 170
  
- quadratic curve, 9
- quadratic form, 4, 11, 41
- quadratic formula, 176
- quaternions, 190
- quiver, 151
- quotient space, 187
  
- range, 24, 184
- rank, 3, 65
- rank of linear system, 69
- rank of matrix, 66
- real normal operators, 112
- real part, 173
- real spectral theorem, 114
- real vector space, 22
- realification, 112
- reduced row echelon form, 74
- reducible representation, 155
- reflection, 161
- regular nilpotent, 129
- representation, 151
- right inverse, 71
- right kernel, 189
- right vector space, 191
- root of unity, 177
- root space, 128
- root system, 162
- row echelon form, 74
- row echelon form of rank  $r$ , 74
- row space, 76
  
- scalar, 21, 22
- scalar product, 169
- semiaxes, 119
- semiaxis of ellipse, 13
- sesquilinear form, 42, 93
- sign of permutation, 47
- similarity, 127

- similarity transformation, 38
- simple problems, 19
- simple quiver, 155
- singular value decomposition, 109
- skew-filed, 190
- span, 26
- spectral theorem, 102
- spectrum, 5, 117
- square matrix, 36, 184
- square root, 109, 177
- standard basis, 26
- standard coordinate flag, 82
- standard coordinate space, 23
- standard Euclidean space, 43
- standard Hermitian space, 43
- subrepresentation, 155
- subspace, 23
- surjective, 24, 185
- Sylvester, 95
- symmetric bilinear form, 40, 87
- symmetric matrix, 41
- symmetric operator, 114
- symmetry, 169
- system of linear equations, 43
  
- tail, 165
- tautology, 189
- time-independent dynamical system, 139
- total anti-symmetry, 50
- trace, 108
- transition matrix, 37
- transposed form, 40
- transposed map, 38
- transposed matrix, 39
- transposed partition, 131
- transposition matrix, 79
- transposition permutation, 49
- triangle inequality, 99, 172
  
- unipotent matrix, 81
- unit coordinate vectors, 26
- unit vector, 169
- unitary rotation, 105
- unitary space, 99
- unitary transformation, 102
- upper triangular, 79
- upper-triangular matrix, 184
  
- Vandermonde, 64
- Vandermonde's identity, 64
- vector, 21, 165
- vector space, 18, 21
- vector subspace, 23
- vector sum, 165
- vertex, 151
- Vieta, 180
- Vieta's theorem, 178
  
- Young tableaux, 130
  
- zero representation, 154
- zero vector, 21, 166