

A Review of Beam Polarization, FCC-ee and Polarizer Ring Development

M. Watson^{1, a)}

Physics Department, University of Surrey UK

(Dated: 29 June 2026)

Your abstract here.

I. INTRODUCTION

Text here.

II. LEP VS. FCC-EE

The Large Electron-Positron Collider (LEP), which operated at CERN between 1989 and 2000, ushered in the era of modern precision electroweak physics and established the benchmark against which the performance goals of the FCC-ee are evaluated¹. LEP initially operated around the Z-resonance before expanding its energy reach past the W-pair production threshold². Through dedicated line-shape scans, compiled in the final combined legacy reports of the LEP collaborations, LEP determined the Z-boson mass and width to $m_Z = 91187.5 \pm 2.1$ MeV and $\Gamma_Z = 2495.2 \pm 2.3$ MeV, respectively^{2,3}. The collider also provided a precise determination of the number of active light neutrino species¹, while measurements of fermion pair-production cross-sections around the Z-pole offered indirect sensitivity to the top-quark mass prior to its discovery². Ultimately, the W-boson mass was measured at LEP to be $m_W = 80.379 \pm 0.012$ GeV.

These precision measurements were made possible by exploiting the natural transverse polarization of the electron and positron beams via a technique known as resonant depolarization (RDP)⁴. RDP allows the spin tune of the beam to be determined by exciting the particles with a transverse oscillating magnetic field until the polarization is destroyed; the frequency of the depolarizing field then directly maps to the spin precession frequency^{4,5}. At LEP, a polarization level of only 5–10% was sufficient for precise energy calibration⁵. However, this technique was fundamentally limited to beam energies below 61 GeV due to the rapid growth of energy spread and spin-orbit resonances observed during high-energy LEP runs^{5–7}. This limitation is mitigated in the FCC-ee layout; its significantly larger bending radius keeps the relative energy spread small enough to preserve beam polarization up to the W-pair threshold, aiming for a W-boson mass measurement precision of 300 keV¹.

The FCC-ee physics programme targets luminosities up to five orders of magnitude greater than those achieved at LEP⁸. Specifically, the Z-pole operational mode is projected to deliver approximately 10^5 times the LEP statistics⁶, enabling profound improvements in the measurement of key

electroweak observables, including the weak mixing angle, W and Z masses, and widths^{8,9}. Current FCC-ee layout studies project uncertainties of approximately 100 keV and 12 keV on the Z mass and width, respectively—representing an order of magnitude improvement over the final LEP results^{4,10}. Fully exploiting this immense statistical data sample requires an equivalent, unprecedented improvement in beam-energy calibration, as well as drastic reductions in theoretical calculation uncertainties to ensure theory errors do not limit the experimental precision¹¹.

Beyond the electroweak arena, the FCC-ee will act as a high-luminosity Higgs and top-quark factory¹². Precise Higgs coupling measurements are vital, as the Higgs boson receives electroweak radiative corrections from heavy Standard Model particles like the top quark^{12,13}. A primary objective of the high-energy operation is a precise threshold scan of the $e^+e^- \rightarrow t\bar{t}$ production cross-section, providing direct sensitivity to the top quark’s mass, width, strong coupling constant, and Yukawa coupling^{9,12}. Collectively, these high-precision measurements form the core strategy of the FCC-ee to probe physics beyond the Standard Model, providing indirect sensitivity to new physics scales approaching 100 TeV^{9,12}.

Achieving these milestones imposes exceptionally demanding requirements on accelerator sub-systems. Because every primary observable is sensitive to systematic shifts in the centre-of-mass energy (E_{CM}), the extreme statistical precision will become entirely dominated by systematic effects unless the beam energies can be determined with a precision significantly surpassing that of the LEP via global Effective Field Theory (EFT) fits^{9,12,14}.

III. FUNDAMENTALS OF SPIN AND POLARIZATION

A. Spin Motion and Tune in Storage Rings

As an electron or positron travels around an accelerator, its spin doesn’t stay aligned with the guiding field, as it precesses at a different rate to the orbit revolution. Its motion is described by the Thomas-BMT equation¹⁵ which originates from the combination of the Larmor and Thomas precession into a single precession vector¹⁶, described by:

$$\boldsymbol{\Omega} = (1 + G\gamma)\mathbf{B}_\perp + (1 + G)\mathbf{B}_\parallel + \left(\frac{1}{\gamma+1} + G\right)\gamma\frac{\mathbf{E} \times \boldsymbol{\beta}}{c}$$

¹⁷ where $\mathbf{B}_\parallel$ and $\mathbf{B}_\perp$ are the magnetic field components parallel and perpendicular to the velocity $\boldsymbol{\beta}$, G is the gyromagnetic anomaly, and γ is the Lorentz factor. the $\mathbf{E} \times \boldsymbol{\beta}$ term is usually negligible in circular accelerators since RF cavity fields run

^{a)} Also at BE-ABP, CERN.

parallel to the velocity, leaving the much simpler

$$\Omega \approx (1 + G\gamma)\mathbf{B}_\perp + (1 + G)\mathbf{B}_\parallel$$

for most practical purposes.^{17,18}

This precession arises because a particle's magnetic moment, itself proportional to its spin, experiences a torque when placed in a magnetic field. Any magnetic moment residing in a magnetic field experiences a torque that causes it to precess around the field instead of aligning with it¹⁷. The bending and focusing magnets in the storage ring are what keep the beam circulating, therefore the spin is continually subject to this torque. Importantly, the rate of this precession is not tied to the rate of orbital revolution: it depends on the gyromagnetic anomaly $a = (g - 2)/2$, a fixed property of the particle species itself^{15,17}. A second, independent contribution comes from Thomas precession, a purely relativistic effect arising because the particle's own rest frame continuously rotates as it is bent around the ring^{15,17}.

Specialising this result to the simplest case of a particle passing through a single bending magnet, where $B_\perp$ dominates, Berglund's derivation for HERA gives the orbit-deflection relation directly¹⁸:

$$\delta\theta_{spin} = (1 + a\gamma)\delta\theta_{orbit}$$

For every degree the momentum vector turns through the bend, the spin turns by $(1 + a\gamma)$ degrees more than the orbit, by a factor that grows with beam energy. Since $a\gamma$ is never an integer at a real beam energy, the spin doesn't return to its original orientation after one turn; it advances to a new phase with each revolution¹⁸.

The excess precession defines the spin tune $\nu = a\gamma$, the number of spin precessions per orbital revolution, evaluated in the frame moving with the particle^{17,19}. The relationship between spin tune and electron energy is linear, making it incredibly useful in the RDP process as the beam energy can be determined directly^{17,20}.

B. Polarization and the Sokolov-Ternov Effect

Polarization is a quantity representative of the particle ensemble rather than a single-particle spin, and describes the degree to which particles align along a specified direction. It is defined as

$$P = \frac{N_\uparrow - N_\downarrow}{N_\uparrow + N_\downarrow}$$

where...¹⁷

The definition is built for 1/2-spin systems so it is implicitly scoped to electrons/positrons²¹.

Radiative spin polarization was first predicted theoretically by Sokolov and Ternov, who derived an asymptotic polarization limit of $P_0 = 8/(5\sqrt{3}) \approx 92.4\%$ for an idealised uniform field²². This prediction was confirmed experimentally within a decade: the first observations of radiative polarization were made at the VEPP-2 storage ring at Novosibirsk and at the

ACO ring in Orsay, with later runs at the improved VEPP-2M and ACO approaching the theoretical limit^{21,23}. Real machines, however, rarely reach this asymptotic value — SPEAR II, for instance, achieved an equilibrium polarization of only about 76% at 3.7 GeV²⁴, and, as mentioned in Section I, LEP itself operated with a transverse polarization around 10%.

The spontaneous build-up of polarization arises directly from synchrotron radiation and spin-flip asymmetry^{22,25}.

FIG. 1. Caption here.

$$P(t) = P_\infty \left(1 - e^{-t/\tau_{ST}}\right) \quad (1)$$

TABLE I. Caption here.

Column 1	Column 2
Value	Value

ACKNOWLEDGMENTS

Acknowledgements here.

¹M. Dam, “Precision electroweak measurements at the fcc-ee,” (2016), <https://doi.org/10.48550/arXiv.1601.03849>.

²the FCC Collaboration, A. Abada, M. Abbrescia, A. S. S. I. Abdyukhanov, A. Fernandez, A. Abramov, M. Aburaia, A. A. O. A. P. R. P. Agrawal, A.-S. J. A. A.-V. J. J. M. Aiba, I. Aichinger, G. Aielli, A. g. Akay, A. Akhundov, H. Aksakal, and A. J. L., “Fcc-ee: The lepton collider: Future circular collider conceptual design report volume 2,” *The European Physical Journal Special Topics* **228**, 261–623 (2019).

³ALEPH Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, SLD Collaboration, SLD Electroweak Heavy Flavour Group, ALEPH Electroweak Working Group, DELPHI Electroweak Working Group, L3 Electroweak Working Group, OPAL Electroweak Working Group, and LEP Electroweak Working Group, “Precision electroweak measurements on the Z resonance,” *Physics Reports* **427**, 257–454 (2006), [arXiv:hep-ex/0509008](https://arxiv.org/abs/hep-ex/0509008).

⁴M. Benedikt, W. Bartmann, J.-P. Burnet, C. Carli, A. Chance, P. Craievich, M. Giovannozzi, C. Grojean, J. Gutleber, K. Hanke, A. Henriques, P. Janot, C. Lourenco, M. Mangano, T. Otto, J. H. Poole, S. Rajagopalan, T. Raubenheimer, E. Todesco, and L. Ulrici, *Future circular collider feasibility study report volume 2: Accelerators, technical infrastructure and safety* (2025).

⁵A. Blondel, P. Janot, J. Wenninger, R. Aßmann, S. Aumon, P. Azzurri, D. P. Barber, M. Benedikt, A. V. Bogomyagkov, E. Gianfelice-Wendt, K. D. El, I. A. Koop, M. Koratzinos, E. Levitchev, T. Lefevre, A. Milanese, N. Muchnoi, S. A. Nikitin, K. Oide, and E. Perez, “Polarization and centre-of-mass energy calibration at fcc-ee,” (2019), <https://doi.org/10.48550/arXiv.1909.12245>.

⁶M. Koratzinos, “The fcc-ee design study: luminosity and beam polarization,” (Sissa Medialab, Vienna, Austria, 2016) p. 518.

⁷R. Assmann, A. Blondel, B. Dehning, P. Grosse-Wiesmann, R. Jacobsen, J. P. Koutchouk, J. Miles, M. Placidi, R. Schmidt, and J. Wenninger, “Spin dynamics in LEP with 40–100 GeV beams,” *The European Physical Journal C* **21**, 1–23 (2001).

⁸M. Dam, *Detector requirements, design, and technologies for the FCC-ee higgs, electroweak, and top factory* (2025).

⁹D. d’Enterria, “Physics case of fcc-ee,” (2016), <https://doi.org/10.48550/arXiv.1601.06640>.

- ¹⁰J. A. Maestre, A. Blondel, M. Dam, and P. Janot, “The z lineshape challenge: ppm and kev measurements,” *The European Physical Journal Plus* **136**, 848 (2021).
- ¹¹A. Blondel, A. Freitas, S. Jadach, P. Janot, and T. Riemann, “Theory requirements and challenges for the FCC-ee,” (2019), arXiv:1901.02648 [hep-ph].
- ¹²T. F. Collaboration, A. Abada, M. Abbrescia, A. S. S. I. Abdyukhanov, F. J. Abelleira, A. Abramov, M. Aburaia, A. A. O. A. P. R. P. Agrawal, A.-S. J. A. A.-V. J. J. M. Aiba, I. Aichinger, G. Aielli, A. Akay, A. Akhundov, H. Aksakal, and A. J. L., “Fcc physics opportunities: Future circular collider conceptual design report volume 1,” *The European Physical Journal C* **79**, 474 (2019).
- ¹³F. Siringo and G. Buccheri, *Electroweak radiative corrections to higgs production via vector boson fusion using SCET: Numerical results* (2025).
- ¹⁴J. Ellis and T. You, “Global analysis of effective field theory constraints on the FCC-ee,” *The European Physical Journal C* **79**, 708 (2019), arXiv:1901.00310 [hep-ph].
- ¹⁵V. Bargmann, L. Michel, and V. L. Telegdi, “Precession of the polarization of particles moving in a homogeneous electromagnetic field,” *American Physical Society* **2**, 435–436 (1959).
- ¹⁶E. Hamwi and G. Hoffstaetter, *Polarization Transmission in the Electron-Ion Collider's Hadron Storage Ring* (2025).
- ¹⁷*Polarized Beam Dynamics and Instrumentation in Particle Accelerators* (Springer International Publishing, Cham, 2023).
- ¹⁸G. Zara and M. Berglund, *Spin-Orbit Maps and Electron Spin Dynamics for the Luminosity Upgrade Project at HERA* (2001).
- ¹⁹H. Huang, *Measurement of the Spin Tune using the coherent spin motion of polarized protons in a storage ring* (2019).
- ²⁰J. Keintzel, Y. Wu, C. Kiel, D. Sagan, G. Wilkinson, K. Oide, J. Bauche, W. Höfle, A. Blondel, H. Jiang, A. Bogomyagkov, G. Roy, A. Martens, M. Hofer, T. Pieloni, D. Sittard, Z. Zhang, K. Hock, I. Nikolaev, and D. Gaskell, “Jacow : Fcc-ee energy calibration and polarization - status and outlook,” *JACoW IPAC 2025*, MOPM007 (2025).
- ²¹J. Buon and J.-P. Koutchouk, “Polarization of electron and proton beams,” (1995), 10.5170/CERN-1995-006.879.
- ²²A. Sokolov and I. Ternov, *ON THE SPIN AND POLARIZATION EFFECTS IN THE THEORY OF SYNCHROTRON RADIATION* (1963).
- ²³V. Baier, *Forty years of acting electron-positron colliders* (2006).
- ²⁴S. R. Mane, “Radiative spin polarization in high energy storage rings,” (2025), arXiv:2512.13958 [physics.acc-ph].
- ²⁵J. D. Kroth and K. Tuchin, “Sokolov-ternov effect in rotating systems,” *Physical Review D* **111** (2025), <https://doi.org/10.1103/physrevd.111.056004>.