

chemical potential of an ideal solvent :

$$\mu_A = \mu_A^* + RT \ln \frac{p_A}{p_A^*} = \mu_A^* + RT \ln x_A = \mu_{A,\text{modified}}^* + RT \ln b_A$$

chemical potential of an ideal solvent :

$$\mu_A = \mu_A^* + RT \ln \frac{p_A}{p_A^*} = \mu_A^* + RT \ln x_A = \mu_{A,\text{modified}}^* + RT \ln b_A$$

general form for real or ideal solvent : $\mu_A = \mu_A^* + RT \ln a_A$; $a_A = \frac{p_A}{p_A^*}$

chemical potential of an ideal solvent :

$$\mu_A = \mu_A^* + RT \ln \frac{p_A}{p_A^*} = \mu_A^* + RT \ln x_A = \mu_{A,\text{modified}}^* + RT \ln b_A$$

general form for real or ideal solvent : $\mu_A = \mu_A^* + RT \ln a_A$; $a_A = \frac{p_A}{p_A^*}$

activity = activity coefficient \times molar concentration relative to standard value

chemical potential of an ideal solvent :

$$\mu_A = \mu_A^* + RT \ln \frac{p_A}{p_A^*} = \mu_A^* + RT \ln x_A = \mu_{A,\text{modified}}^* + RT \ln b_A$$

general form for real or ideal solvent : $\mu_A = \mu_A^* + RT \ln a_A$; $a_A = \frac{p_A}{p_A^*}$

activity = activity coefficient \times molar concentration relative to standard value

$$\mu_A = \mu_A^* + RT \ln a_A; \quad a_A = \frac{p_A}{p_A^*} \quad \left\{ \begin{array}{l} \text{as } x_A \longrightarrow 1 \\ \gamma_A \longrightarrow 1 \\ a_A \longrightarrow x_A \end{array} \right.$$

activity coefficient, γ : $a_A = \gamma_A x_A$

$$\mu_A = \mu_A^* + RT \ln x_A + RT \ln \gamma_A$$

chemical potential of an ideal solvent :

$$\mu_A = \mu_A^* + RT \ln \frac{p_A}{p_A^*} = \mu_A^* + RT \ln x_A = \mu_{A,\text{modified}}^* + RT \ln b_A$$

general form for real or ideal solvent : $\mu_A = \mu_A^* + RT \ln a_A$; $a_A = \frac{p_A}{p_A^*}$

activity = activity coefficient \times molar concentration relative to standard value

$$\mu_A = \mu_A^* + RT \ln a_A; a_A = \frac{p_A}{p_A^*} \quad \left\{ \begin{array}{l} \text{as } x_A \rightarrow 1 \\ \gamma_A \rightarrow 1 \\ a_A \rightarrow x_A \end{array} \right.$$

activity coefficient, γ : $a_A = \gamma_A x_A$

$$\mu_A = \mu_A^* + RT \ln x_A + RT \ln \gamma_A$$

standard state of solvent: $x_A = 1$

In terms of molality, $a_j = \gamma_j \frac{b_j}{b_j^\ominus}$

Table 5.3 Standard states

Component	Basis	Standard state	Activity	Limits
Solid or liquid		Pure	$a = 1$	
Solvent	Raoult	Pure solvent	$a = p/p^*, a = \gamma x$	$\gamma \rightarrow 1$ as $x \rightarrow 1$ (pure solvent)
Solute	Henry	(1) A hypothetical state of the pure solute	$a = p/K, a = \gamma x$	$\gamma \rightarrow 1$ as $x \rightarrow 0$
		(2) A hypothetical state of the solute at molality b^\ominus	$a = \gamma b/b^\ominus$	$\gamma \rightarrow 1$ as $b \rightarrow 0$

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \frac{p_B}{K_B}; \quad a_B = \gamma_B x_B; \quad \begin{cases} \gamma_B \rightarrow 1 \\ \text{as } x_B \rightarrow 0; (\text{not } 1) \end{cases}$$

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \frac{p_B}{K_B}; \quad a_B = \gamma_B x_B; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } x_B \rightarrow 0; \text{ (not 1)} \end{array} \right.$$

in terms of molalities: selection of standard state is arbitrary

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \frac{p_B}{K_B}; \quad a_B = \gamma_B x_B; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } x_B \rightarrow 0; \text{ (not 1)} \end{array} \right.$$

in terms of molalities: selection of standard state is arbitrary

$$\text{ideal-dilute: } \mu_B = \mu_B^\ominus + RT \ln b_B$$

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \frac{p_B}{K_B}; \quad a_B = \gamma_B x_B; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } x_B \rightarrow 0; (\text{not } 1) \end{array} \right.$$

in terms of molalities: selection of standard state is arbitrary

$$\text{ideal-dilute: } \mu_B = \mu_B^\ominus + RT \ln b_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \gamma_B \frac{b_B}{b_B^\ominus}; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } b_B \rightarrow 0; (\text{not } 1) \end{array} \right. ;$$
$$b_B^\ominus = 1 \text{ mol/kg}$$

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \frac{p_B}{K_B}; \quad a_B = \gamma_B x_B; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } x_B \rightarrow 0; (\text{not } 1) \end{array} \right.$$

in terms of molalities: selection of standard state is arbitrary

$$\text{ideal-dilute: } \mu_B = \mu_B^\ominus + RT \ln b_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \gamma_B \frac{b_B}{b_B^\ominus}; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } b_B \rightarrow 0; (\text{not } 1) \end{array} \right. ;$$
$$b_B^\ominus = 1 \text{ mol/kg}$$

$$\left\{ \begin{array}{l} b_B \rightarrow 0 \\ \text{as } \mu_B \rightarrow -\infty \end{array} \right. : \text{difficult to remove last traces of solute from soln.}$$

solute: it approaches ideal dilute solution as $x_B \rightarrow 0$ (not 1).

$$\text{Ideal-dilute: } \mu_B = \mu_B^* + RT \ln \frac{p_B}{p_B^*} = \mu_B^* + RT \ln \frac{K_B}{p_B^*} + RT \ln x_B = \mu_B^\ominus + RT \ln x_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \frac{p_B}{K_B}; \quad a_B = \gamma_B x_B; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } x_B \rightarrow 0; \text{ (not 1)} \end{array} \right.$$

in terms of molalities: selection of standard state is arbitrary

$$\text{ideal-dilute: } \mu_B = \mu_B^\ominus + RT \ln b_B$$

$$\text{Real: } \mu_B = \mu_B^\ominus + RT \ln a_B; \quad a_B = \gamma_B \frac{b_B}{b_B^\ominus}; \quad \left\{ \begin{array}{l} \gamma_B \rightarrow 1 \\ \text{as } b_B \rightarrow 0; \text{ (not 1)} \end{array} \right. ;$$
$$b_B^\ominus = 1 \text{ mol/kg}$$

$$\left\{ \begin{array}{l} b_B \rightarrow 0 \\ \text{as } \mu_B \rightarrow -\infty \end{array} \right. : \text{difficult to remove last traces of solute from soln.}$$

$$\text{General expression: } \boxed{\mu = \mu^\ominus + RT \ln a = \mu^{\text{ideal}} + RT \ln \gamma}$$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

$$\mu(H^+) = \mu^\ominus(H^+) + RT \ln a(H^+) = \mu^\ominus(H^+) - (RT \ln 10) \times pH$$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

$$\mu(H^+) = \mu^\ominus(H^+) + RT \ln a(H^+) = \mu^\ominus(H^+) - (RT \ln 10) \times pH$$

$$\mu^\oplus(H^+) = \mu^\ominus(H^+) - 7RT \ln 10$$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

$$\mu(H^+) = \mu^\ominus(H^+) + RT \ln a(H^+) = \mu^\ominus(H^+) - (RT \ln 10) \times pH$$

$$\mu^\oplus(H^+) = \mu^\ominus(H^+) - 7RT \ln 10$$

$$\text{At } 298 \text{ K, } 7RT \ln 10 = 39.96 \text{ kJ mol}^{-1}$$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

$$\mu(H^+) = \mu^\ominus(H^+) + RT \ln a(H^+) = \mu^\ominus(H^+) - (RT \ln 10) \times pH$$

$$\mu^\oplus(H^+) = \mu^\ominus(H^+) - 7RT \ln 10$$

At 298 K, $7RT \ln 10 = 39.96 \text{ kJ mol}^{-1}$

two standard values differ by about 40 kJ mol^{-1}

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
- ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
 - ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine
- ▶ infinitesimal amount $d\xi$ of A turns into B

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
 - ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine
- ▶ infinitesimal amount $d\xi$ of A turns into B
 - ▶ change in amount of A, $dn_A = -d\xi$
 - ▶ change in amount of B, $dn_B = d\xi$

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
 - ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine
- ▶ infinitesimal amount $d\xi$ of A turns into B
 - ▶ change in amount of A, $dn_A = -d\xi$
 - ▶ change in amount of B, $dn_B = d\xi$
- ▶ If extent of reaction changes by a finite amount $\Delta\xi$

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
 - ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine
- ▶ infinitesimal amount $d\xi$ of A turns into B
 - ▶ change in amount of A, $dn_A = -d\xi$
 - ▶ change in amount of B, $dn_B = d\xi$
- ▶ If extent of reaction changes by a finite amount $\Delta\xi$
 - ▶ change in amount of A, $\Delta n_A = -\Delta\xi$
 - ▶ change in amount of B, $\Delta n_B = \Delta\xi$

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
 - ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine
- ▶ infinitesimal amount $d\xi$ of A turns into B
 - ▶ change in amount of A, $dn_A = -d\xi$
 - ▶ change in amount of B, $dn_B = d\xi$
- ▶ If extent of reaction changes by a finite amount $\Delta\xi$
 - ▶ change in amount of A, $\Delta n_A = -\Delta\xi$
 - ▶ change in amount of B, $\Delta n_B = \Delta\xi$
- ▶ reaction Gibbs energy : $\Delta_r G = \left(\frac{\partial G}{\partial \xi} \right)_{p, T}$; ξ measures extent of reaction

Chemical equilibrium

- ▶ Consider $A \rightleftharpoons B$,
 - ▶ e.g., pentane \rightleftharpoons 2-methylbutane or l-alanine \rightleftharpoons d-alanine
- ▶ infinitesimal amount $d\xi$ of A turns into B
 - ▶ change in amount of A, $dn_A = -d\xi$
 - ▶ change in amount of B, $dn_B = d\xi$
- ▶ If extent of reaction changes by a finite amount $\Delta\xi$
 - ▶ change in amount of A, $\Delta n_A = -\Delta\xi$
 - ▶ change in amount of B, $\Delta n_B = \Delta\xi$
- ▶ reaction Gibbs energy : $\Delta_r G = \left(\frac{\partial G}{\partial \xi} \right)_{p, T}$; ξ measures extent of reaction
 - ▶ $\Delta_r G$ is a derivative

suppose the reaction advances by $d\xi$,

suppose the reaction advances by $d\xi$,

$$dG = \mu_A dn_A + \mu_B dn_B = -\mu_A d\xi + \mu_B d\xi = (\mu_B - \mu_A) d\xi$$

suppose the reaction advances by $d\xi$,

$$dG = \mu_A dn_A + \mu_B dn_B = -\mu_A d\xi + \mu_B d\xi = (\mu_B - \mu_A) d\xi$$


$$\text{or, } \Delta_r G = \left(\frac{\partial G}{\partial \xi} \right)_{p, T} = \mu_B - \mu_A$$


suppose the reaction advances by $d\xi$,

$$dG = \mu_A dn_A + \mu_B dn_B = -\mu_A d\xi + \mu_B d\xi = (\mu_B - \mu_A) d\xi$$


$$\text{or, } \Delta_r G = \left(\frac{\partial G}{\partial \xi} \right)_{p, T} = \mu_B - \mu_A$$

= difference between chemical potentials of reactants and products
at the composition of the reaction mixture







- ▶ As reaction advances :
slope of G changes



- ▶ As reaction advances :
slope of G changes
- ▶ Exergonic:
 $\Delta_r G < 0 \implies$ forward reaction
spontaneous



- ▶ As reaction advances :
slope of G changes
- ▶ Exergonic:
 $\Delta_r G < 0 \Rightarrow$ forward reaction
spontaneous
- ▶ Endergonic:
 $\Delta_r G > 0 \Rightarrow$ reverse reaction
spontaneous



- ▶ As reaction advances :
slope of G changes
- ▶ Exergonic:
 $\Delta_r G < 0 \Rightarrow$ forward reaction
spontaneous
- ▶ Endergonic:
 $\Delta_r G > 0 \Rightarrow$ reverse reaction
spontaneous
- ▶ Equilibrium corresponds to zero slope
 $\Delta_r G = 0 \Rightarrow$ foot of the valley
: reaction at equilibrium

coupling of exergonic and endergonic reactions :

coupling of exergonic and endergonic reactions :



coupling of exergonic and endergonic reactions :



▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature
- ▶ ATP acts as a phosphate donor to a number of acceptors (e.g., glucose), but is recharged by more powerful phosphate donors in a number of biochemical processes

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature
- ▶ ATP acts as a phosphate donor to a number of acceptors (e.g., glucose), but is recharged by more powerful phosphate donors in a number of biochemical processes
 - ▶ Synthesis of ATP is Endergonic $\Delta_r G^\oplus = 31 \text{ kJ mol}^{-1}$

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature
- ▶ ATP acts as a phosphate donor to a number of acceptors (e.g., glucose), but is recharged by more powerful phosphate donors in a number of biochemical processes
 - ▶ Synthesis of ATP is Endergonic $\Delta_r G^\oplus = 31 \text{ kJ mol}^{-1}$
- ▶ To synthesize in cells, we couple the reverse reaction with an exergonic reaction

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature
- ▶ ATP acts as a phosphate donor to a number of acceptors (e.g., glucose), but is recharged by more powerful phosphate donors in a number of biochemical processes
 - ▶ Synthesis of ATP is Endergonic $\Delta_r G^\oplus = 31 \text{ kJ mol}^{-1}$
- ▶ To synthesize in cells, we couple the reverse reaction with an exergonic reaction

At blood temperature, $\Delta_r G^\oplus = -147 \text{ kJ mol}^{-1}$ for oxidation of glucose by NAD^+

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature
- ▶ ATP acts as a phosphate donor to a number of acceptors (e.g., glucose), but is recharged by more powerful phosphate donors in a number of biochemical processes
 - ▶ Synthesis of ATP is Endergonic $\Delta_r G^\oplus = 31 \text{ kJ mol}^{-1}$
- ▶ To synthesize in cells, we couple the reverse reaction with an exergonic reaction

At blood temperature, $\Delta_r G^\oplus = -147 \text{ kJ mol}^{-1}$ for oxidation of glucose by NAD^+


- ▶ coupled reaction :
$$\text{C}_6\text{H}_{12}\text{O}_6(\text{aq}) + 2 \text{NAD}^+(\text{aq}) + 2 \text{ADP(aq)} + 2 \text{P}_i^-(\text{aq}) + 2 \text{H}_2\text{O(l)} \\ \longrightarrow 2 \text{CH}_3\text{COCO}_2^-(\text{aq}) + 2 \text{NADH(aq)} + 2 \text{ATP(aq)} + 2 \text{H}_3\text{O}^+(\text{aq})$$

coupling of exergonic and endergonic reactions :

- ▶ $\text{ATP(aq)} + \text{H}_2\text{O(l)} \longrightarrow \text{ADP(aq)} + \text{P}_i^-(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$
 - ▶ $\Delta_r G^\oplus = -31 \text{ kJ mol}^{-1}$; $\Delta_r H^\oplus = -20 \text{ kJ mol}^{-1}$; $\Delta_r S^\oplus = 34 \text{ J K}^{-1} \text{ mol}^{-1}$
 - ▶ $\therefore \Delta_r S^\oplus$ is large, $\Delta_r G^\oplus$ is sensitive to temperature
- ▶ ATP acts as a phosphate donor to a number of acceptors (e.g., glucose), but is recharged by more powerful phosphate donors in a number of biochemical processes
 - ▶ Synthesis of ATP is Endergonic $\Delta_r G^\oplus = 31 \text{ kJ mol}^{-1}$
- ▶ To synthesize in cells, we couple the reverse reaction with an exergonic reaction

At blood temperature, $\Delta_r G^\oplus = -147 \text{ kJ mol}^{-1}$ for oxidation of glucose by NAD^+

- ▶ coupled reaction :



In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

- ▶ actual cell composition may be far from standard, and ATP reaction may be much more potent

In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

- ▶ actual cell composition may be far from standard, and ATP reaction may be much more potent
- ▶ e.g., biosynthesis of sucrose from glucose and fructose can be driven by plant enzymes though reaction is endergonic $\Delta_r G^\oplus = 23 \text{ kJ mol}^{-1}$

In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

- ▶ actual cell composition may be far from standard, and ATP reaction may be much more potent
- ▶ e.g., biosynthesis of sucrose from glucose and fructose can be driven by plant enzymes though reaction is endergonic $\Delta_r G^\oplus = 23 \text{ kJ mol}^{-1}$
- ▶ biosynthesis of proteins is strongly endergonic, not only due to enthalpy change but also for large decrease in entropy that occurs when many amino acids are assembled into precisely determined sequence

In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

- ▶ actual cell composition may be far from standard, and ATP reaction may be much more potent
- ▶ e.g., biosynthesis of sucrose from glucose and fructose can be driven by plant enzymes though reaction is endergonic $\Delta_r G^\oplus = 23 \text{ kJ mol}^{-1}$
- ▶ biosynthesis of proteins is strongly endergonic, not only due to enthalpy change but also for large decrease in entropy that occurs when many amino acids are assembled into precisely determined sequence
- ▶ e.g., formation of peptide link is endergonic, with $\Delta_r G^\oplus = 17 \text{ kJ mol}^{-1}$

In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

- ▶ actual cell composition may be far from standard, and ATP reaction may be much more potent
- ▶ e.g., biosynthesis of sucrose from glucose and fructose can be driven by plant enzymes though reaction is endergonic $\Delta_r G^\oplus = 23 \text{ kJ mol}^{-1}$
- ▶ biosynthesis of proteins is strongly endergonic, not only due to enthalpy change but also for large decrease in entropy that occurs when many amino acids are assembled into precisely determined sequence
- ▶ e.g., formation of peptide link is endergonic, with $\Delta_r G^\oplus = 17 \text{ kJ mol}^{-1}$
- ▶ but biosynthesis occurs indirectly and is equivalent to consumption of 3 ATP molecules for each link

In the cell, each ATP molecule is used to drive an endergonic reaction for which $\Delta_r G^\oplus \not\approx 31 \text{ kJ mol}^{-1}$

- ▶ actual cell composition may be far from standard, and ATP reaction may be much more potent
- ▶ e.g., biosynthesis of sucrose from glucose and fructose can be driven by plant enzymes though reaction is endergonic $\Delta_r G^\oplus = 23 \text{ kJ mol}^{-1}$
- ▶ biosynthesis of proteins is strongly endergonic, not only due to enthalpy change but also for large decrease in entropy that occurs when many amino acids are assembled into precisely determined sequence
- ▶ e.g., formation of peptide link is endergonic, with $\Delta_r G^\oplus = 17 \text{ kJ mol}^{-1}$
- ▶ but biosynthesis occurs indirectly and is equivalent to consumption of 3 ATP molecules for each link

In moderately small protein like myoglobin, with about 150 peptide links, construction alone requires 450 ATP molecules

Perfect gas equilibria :

Perfect gas equilibria :

When A and B are perfect gases

$$\Delta_r G = \mu_B - \mu_A = (\mu_B^\ominus + RT \ln p_B) - (\mu_A^\ominus + RT \ln p_A)$$

Perfect gas equilibria :

When A and B are perfect gases

$$\begin{aligned}\Delta_r G &= \mu_B - \mu_A = (\mu_B^\ominus + RT \ln p_B) - (\mu_A^\ominus + RT \ln p_A) \\ &= \Delta_r G^\ominus + RT \ln \frac{p_B}{p_A} = \Delta_r G^\ominus + RT \ln Q;\end{aligned}$$

Perfect gas equilibria :

When A and B are perfect gases

$$\begin{aligned}\Delta_r G &= \mu_B - \mu_A = (\mu_B^\ominus + RT \ln p_B) - (\mu_A^\ominus + RT \ln p_A) \\ &= \Delta_r G^\ominus + RT \ln \frac{p_B}{p_A} = \Delta_r G^\ominus + RT \ln Q; \quad Q = \text{reaction quotient}\end{aligned}$$

Perfect gas equilibria :

When A and B are perfect gases

$$\begin{aligned}\Delta_r G &= \mu_B - \mu_A = (\mu_B^\ominus + RT \ln p_B) - (\mu_A^\ominus + RT \ln p_A) \\ &= \Delta_r G^\ominus + RT \ln \frac{p_B}{p_A} = \Delta_r G^\ominus + RT \ln Q; \quad Q = \text{reaction quotient} \\ \text{and } \Delta_r G^\ominus &= \Delta_f G_m^\ominus(B) - \Delta_f G_m^\ominus(A) = \text{standard reaction Gibbs energy}\end{aligned}$$

Perfect gas equilibria :

When A and B are perfect gases

$$\begin{aligned}\Delta_r G &= \mu_B - \mu_A = (\mu_B^\ominus + RT \ln p_B) - (\mu_A^\ominus + RT \ln p_A) \\ &= \Delta_r G^\ominus + RT \ln \frac{p_B}{p_A} = \Delta_r G^\ominus + RT \ln Q; \quad Q = \text{reaction quotient}\end{aligned}$$

and $\Delta_r G^\ominus = \Delta_f G_m^\ominus(B) - \Delta_f G_m^\ominus(A)$ = standard reaction Gibbs energy

At equilibrium, $\Delta_r G = 0$

Perfect gas equilibria :

When A and B are perfect gases

$$\begin{aligned}\Delta_r G &= \mu_B - \mu_A = (\mu_B^\ominus + RT \ln p_B) - (\mu_A^\ominus + RT \ln p_A) \\ &= \Delta_r G^\ominus + RT \ln \frac{p_B}{p_A} = \Delta_r G^\ominus + RT \ln Q; \quad Q = \text{reaction quotient}\end{aligned}$$

and $\Delta_r G^\ominus = \Delta_f G_m^\ominus(B) - \Delta_f G_m^\ominus(A)$ = standard reaction Gibbs energy

At equilibrium, $\Delta_r G = 0 \implies \Delta_r G^\ominus + RT \ln K$ with $K = \left(\frac{p_B}{p_A}\right)_{\text{equilibr}}$

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$,

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$, $\Delta_r G = \Delta_r G^\ominus + RT \ln Q$

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$, $\Delta_r G = \Delta_r G^\ominus + RT \ln Q$

$\Delta_r G^\ominus = \nu_C \mu_C^\ominus + \nu_D \mu_D^\ominus - \nu_A \mu_A^\ominus - \nu_B \mu_B^\ominus$ and $Q = \frac{p_D^{\nu_D} p_C^{\nu_C}}{p_A^{\nu_A} p_B^{\nu_B}}$ if A, ..., D are perfect gases

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$, $\Delta_r G = \Delta_r G^\ominus + RT \ln Q$

$$\Delta_r G^\ominus = \nu_C \mu_C^\ominus + \nu_D \mu_D^\ominus - \nu_A \mu_A^\ominus - \nu_B \mu_B^\ominus \text{ and } Q = \frac{p_D^{\nu_D} p_C^{\nu_C}}{p_A^{\nu_A} p_B^{\nu_B}} \text{ if A, } \dots, \text{ D are perfect gases}$$

At equilibrium: $\Delta_r G = 0 \xrightarrow{Q=K}$

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$, $\Delta_r G = \Delta_r G^\ominus + RT \ln Q$

$\Delta_r G^\ominus = \nu_C \mu_C^\ominus + \nu_D \mu_D^\ominus - \nu_A \mu_A^\ominus - \nu_B \mu_B^\ominus$ and $Q = \frac{p_D^{\nu_D} p_C^{\nu_C}}{p_A^{\nu_A} p_B^{\nu_B}}$ if A, ..., D are perfect gases

At equilibrium: $\Delta_r G = 0 \xrightarrow{Q=K} \boxed{\Delta_r G^\ominus = -RT \ln K}$

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$, $\Delta_r G = \Delta_r G^\ominus + RT \ln Q$

$\Delta_r G^\ominus = \nu_C \mu_C^\ominus + \nu_D \mu_D^\ominus - \nu_A \mu_A^\ominus - \nu_B \mu_B^\ominus$ and $Q = \frac{p_D^{\nu_D} p_C^{\nu_C}}{p_A^{\nu_A} p_B^{\nu_B}}$ if A, ..., D are perfect gases

At equilibrium: $\Delta_r G = 0 \xrightarrow{Q=K} \boxed{\Delta_r G^\ominus = -RT \ln K}$

For real gases, pressures replaced by fugacities and more generally for arbitrary phases, by activities.

In general, for a reaction $\nu_A A + \nu_B B \rightleftharpoons \nu_C C + \nu_D D$, $\Delta_r G = \Delta_r G^\ominus + RT \ln Q$

$$\Delta_r G^\ominus = \nu_C \mu_C^\ominus + \nu_D \mu_D^\ominus - \nu_A \mu_A^\ominus - \nu_B \mu_B^\ominus \text{ and } Q = \frac{p_D^{\nu_D} p_C^{\nu_C}}{p_A^{\nu_A} p_B^{\nu_B}} \text{ if A, } \dots, \text{ D are perfect gases}$$


At equilibrium: $\Delta_r G = 0 \xrightarrow{Q=K} \boxed{\Delta_r G^\ominus = -RT \ln K}$

For real gases, pressures replaced by fugacities and more generally for arbitrary phases, by activities.

$$Q = \frac{\text{activities of products raised to powers of stoichiometric coefficients}}{\text{activities of reactants raised to powers of stoichiometric coefficients}}$$


Writing ν s for reactants as negative integers, $Q = \prod_j a_j^{\nu_j}$

In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$




In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$

- ▶ If mixing of reactants + products ignored
 $\Rightarrow G$ changes linearly from initial (pure reactants) to final value (pure products)




In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$

- ▶ If mixing of reactants + products ignored
 $\Rightarrow G$ changes linearly from initial (pure reactants) to final value (pure products)
- ▶ slope = $\Delta_r G^\ominus$.




In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$




- ▶ If mixing of reactants + products ignored
 $\Rightarrow G$ changes linearly from initial (pure reactants) to final value (pure products)
- ▶ slope = $\Delta_r G^\ominus$.
- ▶ as products are produced :
further contribution to G arising from mixing (lowest curve)

In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$




- ▶ If mixing of reactants + products ignored
 $\implies G$ changes linearly from initial (pure reactants) to final value (pure products)
- ▶ slope = $\Delta_r G^\ominus$.
- ▶ as products are produced :
 further contribution to G arising from mixing (lowest curve)
- ▶ $\Delta G_{\text{mix}} = nRT(x_A \ln x_A + x_B \ln x_B)$

In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$



- ▶ If mixing of reactants + products ignored
 $\Rightarrow G$ changes linearly from initial (pure reactants) to final value (pure products)
- ▶ slope = $\Delta_r G^\ominus$.
- ▶ as products are produced :
further contribution to G arising from mixing (lowest curve)
- ▶ $\Delta G_{\text{mix}} = nRT(x_A \ln x_A + x_B \ln x_B)$
- ▶ sum of two contributions has a minimum

In molecular terms, the minimum at $\Delta_r G^\ominus = 0$ stems from $\Delta_{\text{mix}} G$



- ▶ If mixing of reactants + products ignored
 $\Rightarrow G$ changes linearly from initial (pure reactants) to final value (pure products)
- ▶ slope = $\Delta_r G^\ominus$.
- ▶ as products are produced :
 further contribution to G arising from mixing (lowest curve)
- ▶ $\Delta G_{\text{mix}} = nRT(x_A \ln x_A + x_B \ln x_B)$
- ▶ sum of two contributions has a minimum
 - ▶ corresponds to equilibrium composition

dependence of $\Delta_r G$ on Q : $dG = \sum_j \mu_j dn_j = \sum_j \nu_j \mu_j d\xi$

$$\Delta_r G = \left(\frac{\partial G}{\partial \xi} \right)_{p, T} = \sum_j \nu_j \mu_j = \underbrace{\sum_j \nu_j \mu_j^\ominus}_{\Delta_r G^\ominus} + RT \sum_j \nu_j \ln a_j = \Delta_r G^\ominus + RT \ln \overbrace{\prod_j a_j^{\nu_j}}^Q$$

dependence of $\Delta_r G$ on Q : $dG = \sum_j \mu_j dn_j = \sum_j \nu_j \mu_j d\xi$

$$\begin{aligned}\Delta_r G &= \left(\frac{\partial G}{\partial \xi} \right)_{p, T} = \sum_j \nu_j \mu_j = \underbrace{\sum_j \nu_j \mu_j^\ominus}_{\Delta_r G^\ominus} + RT \sum_j \nu_j \ln a_j = \Delta_r G^\ominus + RT \ln \overbrace{\prod_j a_j^{\nu_j}}^Q \\ &= \Delta_r G^\ominus + RT \ln Q\end{aligned}$$

dependence of $\Delta_r G$ on Q : $dG = \sum_j \mu_j dn_j = \sum_j \nu_j \mu_j d\xi$

$$\Delta_r G = \left(\frac{\partial G}{\partial \xi} \right)_{p, T} = \sum_j \nu_j \mu_j = \underbrace{\sum_j \nu_j \mu_j^\ominus}_{\Delta_r G^\ominus} + RT \sum_j \nu_j \ln a_j = \Delta_r G^\ominus + RT \ln \overbrace{\prod_j a_j^{\nu_j}}^Q$$
$$= \Delta_r G^\ominus + RT \ln Q$$

$$K = \left(\prod_j a_j^{\nu_j} \right)_{\text{equil}} ; \Delta_r G^\ominus = -RT \ln K$$

In elementary applications, activities are often replaced by:

In elementary applications, activities are often replaced by:


- ▶ molalities, by replacing a_j by $\frac{b_j}{b^\ominus}$ where $b^\ominus = 1 \text{ mol kg}^{-1}$


In elementary applications, activities are often replaced by:

- ▶ molalities, by replacing a_j by $\frac{b_j}{b^\ominus}$ where $b^\ominus = 1 \text{ mol kg}^{-1}$
- ▶ molar concentrations, by replacing a_j by $\frac{[j]}{c^\ominus}$, where $c^\ominus = 1 \text{ mol dm}^{-3}$


In elementary applications, activities are often replaced by:

- ▶ molalities, by replacing a_j by $\frac{b_j}{b^\ominus}$ where $b^\ominus = 1 \text{ mol kg}^{-1}$
- ▶ molar concentrations, by replacing a_j by $\frac{[j]}{c^\ominus}$, where $c^\ominus = 1 \text{ mol dm}^{-3}$
- ▶ partial pressures, by replacing a_j by $\frac{p_j}{p^\ominus}$, where $p^\ominus = 1 \text{ bar}$






$$K = a_{\text{CaCO}_3(\text{s})}^{-1} \cdot a_{\text{CaO}(\text{s})} \cdot a_{\text{CO}_2(\text{g})} = \frac{\overbrace{a_{\text{CaO}(\text{s})} \cdot a_{\text{CO}_2(\text{g})}}^{=1}}{\underbrace{a_{\text{CaCO}_3(\text{s})}}_{=1}} = a_{\text{CO}_2(\text{g})}$$




$$K = a_{\text{CaCO}_3(\text{s})}^{-1} \cdot a_{\text{CaO}(\text{s})} \cdot a_{\text{CO}_2(\text{g})} = \frac{\overbrace{a_{\text{CaO}(\text{s})} \cdot a_{\text{CO}_2(\text{g})}}^{=1}}{\underbrace{a_{\text{CaCO}_3(\text{s})}}_{=1}} = a_{\text{CO}_2(\text{g})}$$

Provided CO_2 can be treated as a perfect gas, $K = \frac{p_{\text{CO}_2(\text{g})}}{p^\ominus} = p_{\text{CO}_2(\text{g})}$

=numerical value of decomposition vapour pressure of calcium carbonate



$$\Delta_r G^\ominus = 2\Delta_f G^\ominus(\text{NH}_3, \text{g}) - [\Delta_f G^\ominus(\text{N}_2, \text{g}) + 3\Delta_f G^\ominus(\text{H}_2, \text{g})]$$



$$\begin{aligned}\Delta_r G^\ominus &= 2\Delta_f G^\ominus(\text{NH}_3, \text{g}) - [\Delta_f G^\ominus(\text{N}_2, \text{g}) + 3\Delta_f G^\ominus(\text{H}_2, \text{g})] \\ &= 2\Delta_f G^\ominus(\text{NH}_3, \text{g}) = 2 \times (-16.5) \text{ kJ mol}^{-1}\end{aligned}$$

$$\Delta_r G^\ominus = 2\Delta_f G^\ominus(\text{NH}_3, \text{g}) - [\Delta_f G^\ominus(\text{N}_2, \text{g}) + 3\Delta_f G^\ominus(\text{H}_2, \text{g})]$$

$$\therefore \ln K = -\frac{2 \times (-16.5 \times 10^3)}{8.3145 \times 298} = 6.1 \times 10^5$$

$$K = \frac{a_D^{\nu D} a_C^{\nu C}}{a_A^{\nu A} a_B^{\nu B}}$$

$$K = \frac{a_D^{\nu D} a_C^{\nu C}}{a_A^{\nu A} a_B^{\nu B}} = \frac{\gamma_D^{\nu D} \gamma_C^{\nu C}}{\gamma_A^{\nu A} \gamma_B^{\nu B}} \times \frac{b_D^{\nu D} b_C^{\nu C}}{b_A^{\nu A} b_B^{\nu B}} = K_\gamma K_b, \text{ where } b_j \equiv \frac{b_j}{b_j^\ominus}$$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

For reaction: $R + \nu H^+(aq) \rightarrow P$

$$\Delta_r G = \Delta_r G^\ominus + RT \ln \frac{a_P}{a_R a_{H^+}^\nu} = \Delta_r G^\ominus + RT \ln \frac{a_P}{a_R} - \nu RT \ln a_{H^+}$$

biological standard state: $a_{H^+} = 10^{-7}$; $pH = 7$

For reaction: $R + \nu H^+(aq) \rightarrow P$

$$\Delta_r G = \Delta_r G^\ominus + RT \ln \frac{a_P}{a_R a_{H^+}^\nu} = \Delta_r G^\ominus + RT \ln \frac{a_P}{a_R} - \nu RT \ln a_{H^+}$$

In the biological standard state, both P and R are at unit activity

$$\therefore \Delta_r G = \Delta_r G^\ominus - \nu RT \ln 10 \log a_{H^+} = \Delta_r G^\ominus + \nu RT \ln 10 \cdot pH$$

$$\text{with } pH = 7, \quad \Delta_r G^\oplus = \Delta_r G^\ominus + 7\nu RT \ln 10$$

response of equilibria to the conditions


Le Chatelier's principle :

A system at equilibrium, when subjected to a disturbance, responds in a way that tends to minimize the effect of the disturbance

How equilibria respond to changes of pressure

Consider reaction $A \rightleftharpoons 2B$

$$\alpha = \sqrt{\frac{1}{1 + \frac{4p}{Kp^\ominus}}}$$




label : K

How equilibria respond to changes of pressure

Consider reaction $A \rightleftharpoons 2B$

$$\alpha = \sqrt{\frac{1}{1 + \frac{4p}{Kp^\ominus}}}$$



label : K

even though K is independent of pressure
amounts of A and B do depend on pressure
as p is increased, α decreases
in accord with Le Chatelier's principle

effect of increase in pressure on ammonia synthesis

$$K = \frac{p_{\text{NH}_3}^2 p^{\ominus 2}}{p_{\text{N}_2} p_{\text{H}_2}^3} = \frac{x_{\text{NH}_3}^2 p^2 p^{\ominus 2}}{x_{\text{N}_2} x_{\text{H}_2}^3 p^4} = \frac{K_x p^{\ominus 2}}{p^2}$$


response of equilibria to changes of temperature

$$\frac{d \ln K}{dT} = \frac{\Delta_r H^\ominus}{RT^2}$$


response of equilibria to changes of temperature

$$\frac{d \ln K}{dT} = \frac{\Delta_r H^\ominus}{RT^2}$$


$$\text{or, } \frac{d \ln K}{d\left(\frac{1}{T}\right)} = -\frac{\Delta_r H^\ominus}{R}$$




K vs. T(K)



At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture




At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture



usually dominant species in a mixture at equilibrium is the one with lower set of energy levels


At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture




usually dominant species in a mixture at equilibrium is the one with lower set of energy levels

In a reaction, entropy plays a role as well as energy

an endothermic reaction



At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture




usually dominant species in a mixture at equilibrium is the one with lower set of energy levels


In a reaction, entropy plays a role as well as energy

B energy levels lie higher than the A energy levels, but they are much more closely spaced

an endothermic reaction




At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture



usually dominant species in a mixture at equilibrium is the one with lower set of energy levels


In a reaction, entropy plays a role as well as energy

an endothermic reaction



B energy levels lie higher than the A energy levels, but they are much more closely spaced \Rightarrow their total population may be considerable and B could even dominate in the reaction mixture at equilibrium


At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture



usually dominant species in a mixture at equilibrium is the one with lower set of energy levels

In a reaction, entropy plays a role as well as energy


an endothermic reaction



B energy levels lie higher than the A energy levels, but they are much more closely spaced \Rightarrow their total population may be considerable and B could even dominate in the reaction mixture at equilibrium

Closely spaced energy levels correlate with high entropy, so entropy effects dominate


At a given temperature, there is a specific distribution of populations, and hence specific composition of reaction mixture



usually dominant species in a mixture at equilibrium is the one with lower set of energy levels

In a reaction, entropy plays a role as well as energy

an endothermic reaction



B energy levels lie higher than the A energy levels, but they are much more closely spaced \Rightarrow their total population may be considerable and B could even dominate in the reaction mixture at equilibrium

Closely spaced energy levels correlate with high entropy, so entropy effects dominate

$$K = e^{-\frac{\Delta_r H^\ominus}{RT}} e^{-\frac{\Delta_r S^\ominus}{R}}$$