





Example of a Carnot engine





$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$






$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$

$$\text{adiabats: } V_A T_h^c = V_D T_c^c \text{ (exponent } c = \frac{C_V}{nR})$$

$$\text{and } V_C T_c^c = V_B T_h^c$$



$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$


adiabats: $V_A T_h^c = V_D T_c^c$ (exponent $c = \frac{C_V}{nR}$)

and $V_C T_c^c = V_B T_h^c$

obtain : $V_A V_C T_h^c T_c^c = V_D V_B T_h^c T_c^c$

and $\therefore \frac{V_A}{V_B} = \frac{V_D}{V_C} \implies q_c = -nRT_c \ln \frac{V_B}{V_A}$

$$\therefore \frac{q_h}{q_c} = -\frac{T_h}{T_c}$$



$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$

adiabats: $V_A T_h^c = V_D T_c^c$ (exponent $c = \frac{C_V}{nR}$)

and $V_C T_c^c = V_B T_h^c$


obtain : $V_A V_C T_h^c T_c^c = V_D V_B T_h^c T_c^c$

and $\therefore \frac{V_A}{V_B} = \frac{V_D}{V_C} \implies q_c = -nRT_c \ln \frac{V_B}{V_A}$

$$\therefore \frac{q_h}{q_c} = -\frac{T_h}{T_c}$$

efficiency, $\eta = \frac{\text{work performed}}{\text{heat absorbed}} = \frac{|w|}{q_h}$

$$= \frac{q_h + q_c}{q_h} = 1 + \frac{q_c}{q_h} = 1 - \frac{T_c}{T_h}$$



$$q_h = nRT_h \ln \frac{V_B}{V_A}; \quad q_c = nRT_c \ln \frac{V_D}{V_C}$$

$$\text{adiabats: } V_A T_h^c = V_D T_c^c \quad (\text{exponent } c = \frac{C_V}{nR})$$

$$\text{and} \quad V_C T_c^c = V_B T_h^c$$

$$\text{obtain : } V_A V_C T_h^c T_c^c = V_D V_B T_h^c T_c^c$$

$$\text{and } \therefore \frac{V_A}{V_B} = \frac{V_D}{V_C} \implies q_c = -nRT_c \ln \frac{V_B}{V_A}$$

$$\therefore \frac{q_h}{q_c} = -\frac{T_h}{T_c}$$


$$\begin{aligned} \text{efficiency, } \eta &= \frac{\text{work performed}}{\text{heat absorbed}} = \frac{|w|}{q_h} \\ &= \frac{q_h + q_c}{q_h} = 1 + \frac{q_c}{q_h} = 1 - \frac{T_c}{T_h} \end{aligned}$$

$$\Delta S = \frac{q_h}{T_h} + 0 + \frac{q_c}{T_c} + 0 = 0$$

2nd Law : all reversible engines have same efficiency regardless of their construction

2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction




Let $\eta_A > \eta_B$

engine A takes heat q_h from hot source and dumps q_c in cold reservoir

2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction




Let $\eta_A > \eta_B$

engine A takes heat q_h from hot source and dumps q_c in cold reservoir

Reverse engine B

2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction



Let $\eta_A > \eta_B$


engine A takes heat q_h from hot source and dumps q_c in cold reservoir

Reverse engine B

engine B acquires energy heat q_c from cold reservoir and releases $q'_h < q_h$ to the hot end

2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction




Let $\eta_A > \eta_B$

engine A takes heat q_h from hot source and dumps q_c in cold reservoir

Reverse engine B


engine B acquires energy heat q_c from cold reservoir and releases $q'_h < q_h$ to the hot end

Form composite engine C : A+B



2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction




Let $\eta_A > \eta_B$

engine A takes heat q_h from hot source and dumps q_c in cold reservoir

Reverse engine B

engine B acquires energy heat q_c from cold reservoir and releases $q'_h < q_h$ to the hot end


Form composite engine C : A+B



C takes heat $q_h - q'_h$ from hot source and converts it completely into work

2nd Law : all reversible engines have same efficiency regardless of their construction

► Equivalent to Kelvin-Planck statement : Proof by contradiction




Let $\eta_A > \eta_B$

engine A takes heat q_h from hot source and dumps q_c in cold reservoir

Reverse engine B

engine B acquires energy heat q_c from cold reservoir and releases $q'_h < q_h$ to the hot end

Form composite engine C : A+B




C takes heat $q_h - q'_h$ from hot source and converts it completely into work

⇒ violation of Kelvin-Planck statement of second law


universality of 2nd law :

any reversible cycle \approx
collection of Carnot cycles
 \oint (arbitrary path)
 $= \sum \oint$ (each cycle)



universality of 2nd law :

any reversible cycle \approx
collection of Carnot cycles
 \oint (arbitrary path)
 $= \sum \oint$ (each cycle)




ΔS along an individual path inside is
cancelled by ΔS along the path it shares with
neighbouring cycle

universality of 2nd law :

any reversible cycle \approx
collection of Carnot cycles

\oint (arbitrary path)

$$= \sum \oint \text{(each cycle)}$$



ΔS along an individual path inside is
cancelled by ΔS along the path it shares with
neighbouring cycle

The approxmn. becomes exact as individual
cycles become infinitesimal


$$\oint \text{(arbitrary path)} = \sum \oint \text{(each cycle)} = 0$$

universality of 2nd law :

any reversible cycle \approx
collection of Carnot cycles

\oint (arbitrary path)

$$= \sum \oint \text{(each cycle)}$$



ΔS along an individual path inside is
cancelled by ΔS along the path it shares with
neighbouring cycle

The approxmn. becomes exact as individual
cycles become infinitesimal

$$\oint \text{(arbitrary path)} = \sum \oint \text{(each cycle)} = 0$$


\therefore all entropy changes cancel except for those
along perimeter of the overall cycle

$$\sum_{\text{all}} \frac{q_{\text{rev}}}{T} = \sum_{\text{perimeter}} \frac{q_{\text{rev}}}{T} = 0$$

universality of 2nd law :

any reversible cycle \approx
collection of Carnot cycles
 \oint (arbitrary path)

$$= \sum \oint \text{(each cycle)}$$



ΔS along an individual path inside is cancelled by ΔS along the path it shares with neighbouring cycle

The approxmn. becomes exact as individual cycles become infinitesimal

$$\oint \text{(arbitrary path)} = \sum \oint \text{(each cycle)} = 0$$

\therefore all entropy changes cancel except for those along perimeter of the overall cycle

$$\sum_{\text{all}} \frac{q_{\text{rev}}}{T} = \sum_{\text{perimeter}} \frac{q_{\text{rev}}}{T} = 0$$

In the limit of infinitesimal cycles, the non-cancelling edges of Carnot cycles match the overall cycle exactly, and the sum becomes an integral

$\Rightarrow dS$ is exact differential and S is state function

Clausius inequality

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$ and dw are negative when energy leaves the system as work

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$ and dw are negative when energy leaves the system as work

$$\therefore -dw_{\text{rev}} \geq -dw \text{ or, } dw \geq dw_{\text{rev}}$$

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$ and dw are negative when energy leaves the system as work

$$\therefore -dw_{\text{rev}} \geq -dw \text{ or, } dw \geq dw_{\text{rev}}$$

$$dU = dq + dw = dq_{\text{rev}} + dw_{\text{rev}}$$

Clausius inequality

more work is done when a change is reversible than when it is irreversible,

$$\text{i.e., } |dw_{\text{rev}}| \geq |dw|$$

$\therefore dw_{\text{rev}}$ and dw are negative when energy leaves the system as work

$$\therefore -dw_{\text{rev}} \geq -dw \text{ or, } dw \geq dw_{\text{rev}}$$


$$dU = dq + dw = dq_{\text{rev}} + dw_{\text{rev}}$$

$$dq_{\text{rev}} \geq dq, \therefore \Delta S = \int \frac{dq_{\text{rev}}}{T} \geq \int \frac{dq}{T}$$

Consider the reverse of an engine :
transfer of energy as heat from
one system - hot source-at T_h
to another system - the cold sink- at T_c

Consider the reverse of an engine :
transfer of energy as heat from
one system - hot source-at T_h
to another system - the cold sink- at T_c


thermodynamic refrigerator
?



energy removed from cool source at temp. T_c
 $= \frac{|q_c|}{T_c}$

Consider the reverse of an engine :
transfer of energy as heat from
one system - hot source-at T_h
to another system - the cold sink- at T_c

thermodynamic refrigerator
?




energy removed from cool source at temp. T_c
 $= \frac{|q_c|}{T_c}$

and deposited in warmer sink at temp. T_h

Consider the reverse of an engine :
transfer of energy as heat from
one system - hot source-at T_h
to another system - the cold sink- at T_c

thermodynamic refrigerator
?




energy removed from cool source at temp. T_c
 $= \frac{|q_c|}{T_c}$

and deposited in warmer sink at temp. T_h

$$\Delta S = -\frac{|q_c|}{T_c} + \frac{|q_c|}{T_h} < 0 \implies \text{not spontaneous}$$

Consider the reverse of an engine :
transfer of energy as heat from
one system - hot source-at T_h
to another system - the cold sink- at T_c

thermodynamic refrigerator
?




energy removed from cool source at temp. T_c
 $= \frac{|q_c|}{T_c}$

and deposited in warmer sink at temp. T_h


$$\Delta S = -\frac{|q_c|}{T_c} + \frac{|q_c|}{T_h} < 0 \implies \text{not spontaneous}$$

to generate more entropy, energy must be
added to the stream that enters the warm sink.
task: find minimum energy to be supplied

Thermodynamic refrigerator




Thermodynamic refrigerator



coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

Thermodynamic refrigerator




coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is c

Thermodynamic refrigerator




coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is c

and more efficient the refrigerator.

Thermodynamic refrigerator



coefficient of performance,


$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is c

and more efficient the refrigerator.

$$\frac{1}{c} = \frac{|q_h| - |q_c|}{|q_c|} = \frac{|q_h|}{|q_c|} - 1$$

Thermodynamic refrigerator



coefficient of performance,

$$c = \frac{\text{energy transferred as heat}}{\text{energy transferred as work}} = \frac{|q_c|}{|w|}$$

The less the work required to achieve a given transfer, greater is c

and more efficient the refrigerator.

$$\frac{1}{c} = \frac{|q_h| - |q_c|}{|q_c|} = \frac{|q_h|}{|q_c|} - 1$$

using $\frac{q_h}{q_c} = -\frac{T_h}{T_c}$, we get, $c = \frac{T_c}{T_h - T_c}$

work to maintain a low temperature is relevant to design of refrigerators

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate $\propto \Delta T$

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate $\propto \Delta T$

If rate at which energy leaks = $A(T_h - T_c)$

where A depends on sample size and insulation

then minimum power, P , required to maintain original ΔT by pumping out energy by heating surroundings

$$P \propto \frac{1}{c} \cdot A(T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c}$$

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate $\propto \Delta T$

If rate at which energy leaks = $A(T_h - T_c)$

where A depends on sample size and insulation

then minimum power, P , required to maintain original ΔT by pumping out energy by heating surroundings

$$P \propto \frac{1}{T_c} \cdot A(T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c} \text{ or, } P \propto (\Delta T)^2$$

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate $\propto \Delta T$

If rate at which energy leaks = $A(T_h - T_c)$

where A depends on sample size and insulation

then minimum power, P , required to maintain original ΔT by pumping out energy by heating surroundings

$$P \propto \frac{1}{T_c} \cdot A(T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c} \text{ or, } P \propto (\Delta T)^2$$

power increases as $(\Delta T)^2$

work to maintain a low temperature is relevant to design of refrigerators

No thermal insulation is perfect

there is always a flow of energy as heat into the sample at a rate $\propto \Delta T$

If rate at which energy leaks = $A(T_h - T_c)$

where A depends on sample size and insulation

then minimum power, P , required to maintain original ΔT by pumping out energy by heating surroundings


$$P \propto \frac{1}{T_c} \cdot A(T_h - T_c) = \frac{A(T_h - T_c)^2}{T_c} \text{ or, } P \propto (\Delta T)^2$$

power increases as $(\Delta T)^2$


\therefore air-conditioners more expensive to run on hot days than on mild days

Equivalence of Kelvin-Planck (K) and Clausius statements (C)

Equivalence of Kelvin-Planck (K) and Clausius statements (C)




Equivalence of Kelvin-Planck (K) and Clausius statements (C)




—C —K


Equivalence of Kelvin-Planck (K) and Clausius statements (C)




—C —K



Equivalence of Kelvin-Planck (K) and Clausius statements (C)




—C \subset —K




—K \subset —C

entropy change in isothermal expansion :

entropy change in isothermal expansion :
consider a perfect gas




entropy change in isothermal expansion :
consider a perfect gas




Reversible : $\Delta U = 0$;

entropy change in isothermal expansion :
consider a perfect gas




Reversible : $\Delta U = 0$; $q = -w$;

entropy change in isothermal expansion :
consider a perfect gas



Reversible : $\Delta U = 0$; $q = -w$; $\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$


entropy change in isothermal expansion :
consider a perfect gas



Reversible : $\Delta U = 0$; $q = -w$; $\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

entropy change in isothermal expansion :
consider a perfect gas




Reversible : $\Delta U = 0$; $q = -w$; $\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

Irreversible : $\Delta U = 0$

entropy change in isothermal expansion :
consider a perfect gas




Reversible : $\Delta U = 0$; $q = -w$; $\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

Irreversible : $\Delta U = 0$

\therefore If isothermal expansion occurs freely, and irreversibly, then $q = -w = p_{\text{ex}}(V_f - V_i)$

entropy change in isothermal expansion :
consider a perfect gas



Reversible : $\Delta U = 0$; $q = -w$; $\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$


Irreversible : $\Delta U = 0$

\therefore If isothermal expansion occurs freely, and irreversibly, then $q = -w = p_{\text{ex}}(V_f - V_i)$

$$\therefore \Delta S_{\text{surr (irrev)}} = -\frac{p_{\text{ex}}(V_f - V_i)}{T} = -\frac{nR(V_f - V_i)}{V_f}$$

ΔS_{sys} remains the same as before
(calculated using reversible path)

entropy change in isothermal expansion :
consider a perfect gas



Reversible : $\Delta U = 0$; $q = -w$; $\Delta S_{\text{sys}} = nR \ln \frac{V_f}{V_i}$

$$\Delta S_{\text{surr (rev)}} = -\Delta S_{\text{sys}}$$

Irreversible : $\Delta U = 0$

\therefore If isothermal expansion occurs freely, and irreversibly, then $q = -w = p_{\text{ex}}(V_f - V_i)$

$$\therefore \Delta S_{\text{surr (irrev)}} = -\frac{p_{\text{ex}}(V_f - V_i)}{T} = -\frac{nR(V_f - V_i)}{V_f}$$

ΔS_{sys} remains the same as before
(calculated using reversible path)

$$\begin{aligned} \Delta S_{\text{total (rev)}} &= 0 \text{ rev} \\ &> 0 \text{ irrev} \end{aligned}$$

Phase transition : $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Phase transition : $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same $\Delta_{\text{vap}}S$ ($\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$)

Phase transition : $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same $\Delta_{\text{vap}}S$ ($\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$)

Ex. liq. Br_2 :

- no complexity like hydrogen bonding in liquid Br_2
- there is no unusual behaviour in the gas phase
- safe to use Trouton's rule

Phase transition : $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same $\Delta_{\text{vap}}S$ ($\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$)

Ex. liq. Br_2 :

- no complexity like hydrogen bonding in liquid Br_2
there is no unusual behaviour in the gas phase
- safe to use Trouton's rule

$$\begin{aligned} \text{- boils at } 59.2^\circ\text{C}, \Delta_{\text{vap}}H^\ominus &= T_b \times 85 \text{ J mol}^{-1} \\ &= 332.4 \times 85 = 28 \text{ kJ mol}^{-1} \end{aligned}$$

Phase transition : $\Delta_{\text{trs}}S = \frac{\Delta_{\text{trs}}H}{T_{\text{trs}}}$

Trouton's rule :

wide range of liquids have approx. same $\Delta_{\text{vap}}S$ ($\approx 85 \text{ J K}^{-1} \text{ mol}^{-1}$)

Ex. liq. Br_2 :

- no complexity like hydrogen bonding in liquid Br_2
there is no unusual behaviour in the gas phase
- safe to use Trouton's rule

$$\begin{aligned} \text{- boils at } 59.2^\circ\text{C}, \Delta_{\text{vap}}H^\ominus &= T_b \times 85 \text{ J mol}^{-1} \\ &= 332.4 \times 85 = 28 \text{ kJ mol}^{-1} \end{aligned}$$

experimental value : $29.45 \text{ kJ mol}^{-1}$


entropy change as a function of temperature (varying heat capacity) :

$$S(T_f) = S(T_i) + \int_{T_i}^{T_f} \frac{dq_{\text{rev}}}{T}$$

entropy change as a function of temperature (varying heat capacity) :

$$S(T_f) = S(T_i) + \int_{T_i}^{T_f} \frac{dq_{\text{rev}}}{T}$$

$$\begin{aligned} \text{const. } p(V) : S(T_f) &= S(T_i) + \int_{T_i}^{T_f} \frac{C_{p(V)} dT}{T} \\ &= S(T_i) + C_{p(V)} \ln \frac{T_f}{T_i} \end{aligned}$$



label : heat capacity

solved problem : Calculate ΔS when 0.500 dm^3 of Ar at 25°C and 1.00 bar expands to 1.000 dm^3 and is simultaneously heated to 100°C

solved problem : Calculate ΔS when 0.500 dm^3 of Ar at 25°C and 1.00 bar expands to 1.000 dm^3 and is simultaneously heated to 100°C


S is a state function

\therefore choose a convenient path from initial state

solved problem : Calculate ΔS when 0.500 dm^3 of Ar at 25°C and 1.00 bar expands to 1.000 dm^3 and is simultaneously heated to 100°C

S is a state function


\therefore choose a convenient path from initial state



solved problem : Calculate ΔS when 0.500 dm^3 of Ar at 25°C and 1.00 bar expands to 1.000 dm^3 and is simultaneously heated to 100°C

S is a state function

\therefore choose a convenient path from initial state



$$\Delta S_1 = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i};$$

solved problem : Calculate ΔS when 0.500 dm^3 of Ar at 25°C and 1.00 bar expands to 1.000 dm^3 and is simultaneously heated to 100°C

S is a state function

\therefore choose a convenient path from initial state

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 1 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{heat}} 1 \text{ dm}^3 \text{ Ar}; 100^\circ\text{C}$

$$\Delta S_{\text{I}} = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{\text{II}} = C_V \ln \frac{T_f}{T_i}$$

solved problem : Calculate ΔS when 0.500 dm^3 of Ar at 25°C and 1.00 bar expands to 1.000 dm^3 and is simultaneously heated to 100°C

S is a state function

\therefore choose a convenient path from initial state


$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 1 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{heat}} 1 \text{ dm}^3 \text{ Ar}; 100^\circ\text{C}$

$$\Delta S_I = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{II} = C_V \ln \frac{T_f}{T_i}$$

$$\begin{aligned} \Delta S &= \Delta S_I + \Delta S_{II} = nR \left(\ln \frac{V_f}{V_i} + \frac{3}{2} \ln \frac{T_f}{T_i} \right) = \frac{p_i V_i}{T_i} \ln \left(\frac{V_f}{V_i} \left[\frac{T_f}{T_i} \right]^{\frac{3}{2}} \right) \\ &= \frac{10^5 \text{ Pa} \times 0.5 \times 10^{-3} \text{ m}^3}{298} \ln \left(\frac{1}{0.5} \left[\frac{373}{298} \right]^{\frac{3}{2}} \right) \\ &= \frac{10^5 \text{ Pa} \times 0.5 \times 10^{-3} \text{ m}^3}{298} \ln \left(\frac{1}{0.5} \left[\frac{373}{298} \right]^{\frac{3}{2}} \right) = 0.173 \text{ JK}^{-1} \end{aligned}$$

solved problem : Calculate ΔS when 0.5 dm^3 of Ar at 25°C and 1.00 bar is compressed to 0.05 dm^3 and is simultaneously cooled to -25°C

solved problem : Calculate ΔS when 0.5 dm^3 of Ar at 25°C and 1.00 bar is compressed to 0.05 dm^3 and is simultaneously cooled to -25°C



solved problem : Calculate ΔS when 0.5 dm^3 of Ar at 25°C and 1.00 bar is compressed to 0.05 dm^3 and is simultaneously cooled to -25°C

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 0.05 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{cool}} 0.05 \text{ dm}^3 \text{ Ar}; -25^\circ\text{C}$

$$\Delta S_1 = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i};$$

solved problem : Calculate ΔS when 0.5 dm^3 of Ar at 25°C and 1.00 bar is compressed to 0.05 dm^3 and is simultaneously cooled to -25°C

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 0.05 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{cool}} 0.05 \text{ dm}^3 \text{ Ar}; -25^\circ\text{C}$

$$\Delta S_{\text{I}} = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{\text{II}} = C_V \ln \frac{T_f}{T_i}$$


solved problem : Calculate ΔS when 0.5 dm^3 of Ar at 25°C and 1.00 bar is compressed to 0.05 dm^3 and is simultaneously cooled to -25°C

$0.5 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{isothermal}]{\text{rev. expansion}} 0.05 \text{ dm}^3 \text{ Ar}; 25^\circ\text{C} \xrightarrow[\text{const. V}]{\text{cool}} 0.05 \text{ dm}^3 \text{ Ar}; -25^\circ\text{C}$

$$\Delta S_{\text{I}} = nR \ln \frac{V_f}{V_i} = \frac{p_i V_i}{T_i} \ln \frac{V_f}{V_i}; \quad \Delta S_{\text{II}} = C_V \ln \frac{T_f}{T_i}$$


$$\begin{aligned} \Delta S &= \Delta S_{\text{I}} + \Delta S_{\text{II}} = nR \left(\ln \frac{V_f}{V_i} + \frac{3}{2} \ln \frac{T_f}{T_i} \right) = \frac{p_i V_i}{T_i} \ln \left(\frac{V_f}{V_i} \left[\frac{T_f}{T_i} \right]^{\frac{3}{2}} \right) \\ &= \frac{10^5 \text{ Pa} \times 0.5 \times 10^{-3} \text{ m}^3}{298} \ln \left(\frac{0.05}{0.5} \left[\frac{248}{298} \right]^{\frac{3}{2}} \right) = \\ &= -0.44 = -0.44 \text{ JK}^{-1} \end{aligned}$$

Heat capacity vs. temp.




entropy vs. T


Heat capacity vs. temp.




entropy vs. T



Heat capacity vs. temp.



entropy vs. T



$$S(T) = S(0) + \int_0^{T_f} \frac{C_p(s)dT}{T} + \frac{\Delta_{\text{fus}} H}{T_{\text{fus}}} + \int_{T_1}^{T_b} \frac{C_p(l)dT}{T} + \frac{\Delta_{\text{vap}} H}{T_{\text{vap}}} + \int_{T_b}^T \frac{C_p(g)dT}{T}$$

All terms except $S(0)$ can be measured calorimetrically

All terms except $S(0)$ can be measured calorimetrically

integrals can be evaluated either graphically or by fitting a polynomial to the data and integrating analytically

All terms except $S(0)$ can be measured calorimetrically

integrals can be evaluated either graphically or by fitting a polynomial to the data and integrating analytically

$\therefore \frac{dT}{T} = d \ln T$, one can evaluate area under a plot of C_p against $\ln T$

All terms except $S(0)$ can be measured calorimetrically

integrals can be evaluated either graphically or by fitting a polynomial to the data and integrating analytically

$\therefore \frac{dT}{T} = d \ln T$, one can evaluate area under a plot of C_p against $\ln T$

Debye extrapolation : $\lim_{T \rightarrow 0} C_p \longrightarrow aT^3$

Ex. S_m^\ominus for $\text{N}_2(\text{g})$ at 25°C calculated from following data :

Ex. S_m^\ominus for $\text{N}_2(\text{g})$ at 25°C calculated from following data :

$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$

Ex. S_m^\ominus for $\text{N}_2(\text{g})$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42

Ex. S_m^\ominus for $N_2(g)$ at $25^\circ C$ calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20
Correction for gas imperfection	0.92

Ex. S_m^\ominus for $N_2(g)$ at $25^\circ C$ calculated from following data :

	$S_m^\ominus / J K^{-1} mol^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20
Correction for gas imperfection	0.92
Total	192.06

Ex. S_m^\ominus for $N_2(g)$ at 25°C calculated from following data :

	$S_m^\ominus / \text{J K}^{-1}\text{mol}^{-1}$
Debye extrapolation	1.92
$\int_{10}^{35.61}$	25.25
Phase transition : 35.61K	6.43
$\int_{35.61}^{63.14}$	23.38
Fusion : 63.14K	11.42
$\int_{63.14}^{77.32}$	11.41
Vaporization : 77.32K	72.13
$\int_{77.32}^{298.15}$	39.20
Correction for gas imperfection	0.92
Total	192.06

$$S_m(298.15\text{K}) = S_m(0) + 192.06 \text{ J K}^{-1}\text{mol}^{-1}$$

Nernst heat theorem : $\lim_{T \rightarrow 0} \Delta S \longrightarrow 0$ for any transformation provided all substances involved are perfectly crystalline

Nernst heat theorem : $\lim_{T \rightarrow 0} \Delta S \longrightarrow 0$ for any transformation provided all substances involved are perfectly crystalline

3rd law (?) :

The entropy of all perfect crystalline substances is zero at $T = 0$

Boltzmann formula : $S = k_B \ln \Omega$

$\Omega =$ #ways the molecules can be arranged in a crystalline form

Boltzmann formula : $S = k_B \ln \Omega$

$\Omega = \# \text{ways the molecules can be arranged in a crystalline form}$

In most cases, $\lim_{T \rightarrow 0} \Omega = 1$, $\therefore \lim_{T \rightarrow 0} S = 0$

Boltzmann formula : $S = k_B \ln \Omega$

$\Omega = \# \text{ways the molecules can be arranged in a crystalline form}$

In most cases, $\lim_{T \rightarrow 0} \Omega = 1$, $\therefore \lim_{T \rightarrow 0} S = 0$

when $\Omega \neq 1$, then $S = S(0)$, residual entropy

Ice-I_h :

each H atom can lie either close to or far
from its 'parent' O atom

total # of arrangements in sample of N

H₂O molecules with $2N$ H atoms = 2^{2N}


Ice-I_h :

each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of N H₂O molecules with $2N$ H atoms = 2^{2N}

consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H₂O molecule is $2^4 = 16$



Ice-I_h :


each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of N H₂O molecules with $2N$ H atoms = 2^{2N}

consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H₂O molecule is $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds



Ice-I_h :

each H atom can lie either close to or far from its 'parent' O atom


total # of arrangements in sample of N H₂O molecules with $2N$ H atoms = 2^{2N}

consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H₂O molecule is $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

only $\frac{6}{16} = \frac{3}{8}$ of all arrangements are possible, and for N molecules only $(3/8)^N$ of all arrangements are possible



Ice-I_h :

each H atom can lie either close to or far from its 'parent' O atom


total # of arrangements in sample of N H₂O molecules with $2N$ H atoms = 2^{2N}

consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H₂O molecule is $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

only $\frac{6}{16} = \frac{3}{8}$ of all arrangements are possible, and for N molecules only $(3/8)^N$ of all arrangements are possible



\therefore for N water molecules
of possible configurations
 $= 2^{2N} (3/8)^N = \left(\frac{3}{2}\right)^N$

Ice-I_h :


each H atom can lie either close to or far from its 'parent' O atom

total # of arrangements in sample of N H₂O molecules with $2N$ H atoms = 2^{2N}
consider a single central O atom.

total number of arrangements of locations of H atoms around central O atom of one H₂O molecule is $2^4 = 16$

Of these , only 6 correspond to two short and two long bonds

only $\frac{6}{16} = \frac{3}{8}$ of all arrangements are possible, and for N molecules only $(3/8)^N$ of all arrangements are possible



\therefore for N water molecules
of possible configurations
 $= 2^{2N} (3/8)^N = \left(\frac{3}{2}\right)^N$

$$S(0) = Nk_B \ln \left(\frac{3}{2}\right) = 3.37 \text{ J mol}^{-1}\text{K}^{-1}$$

molecular interpretation of increase in entropy with volume


molecular interpretation of increase in entropy with volume

As the system size increases
energy levels move closer together and more
become accessible to the molecules

molecular interpretation of increase in entropy with volume

As the system size increases
energy levels move closer together and more
become accessible to the molecules

ways of achieving same energy (Ω) increases



Maxwell's demon : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers

Maxwell's demon : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers

Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

Maxwell's demon : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers

Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

expansion is irreversible as there is no way to get the gas back into the initial chamber without doing work.

Maxwell's demon : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum


Open the tap and gas in first chamber expands to fill both chambers

Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

expansion is irreversible as there is no way to get the gas back into the initial chamber without doing work.

Maxwell imagined that the tap was operated by a microscopic intelligent creature. If it sees a molecule heading from the second chamber back into the first, it opens the tap and then shuts it, just letting the molecule through. If it sees a molecule heading from the first chamber back into the second chamber, it keeps the tap closed



Maxwell's demon : a thought experiment

A gas initially in one chamber, connected via a closed tap to a second chamber containing only vacuum

Open the tap and gas in first chamber expands to fill both chambers


Equilibrium established and pressure in each

chamber is now half of what it was in the first chamber at the start

expansion is irreversible as there is no way to get the gas back into the initial chamber without doing work.

Maxwell imagined that the tap was operated by a microscopic intelligent creature. If it sees a molecule heading from the second chamber back into the first, it opens the tap and then shuts it, just letting the molecule through. If it sees a molecule heading from the first chamber back into the second chamber, it keeps the tap closed

Demon does no work and yet it makes molecules in second chamber all go back into the first chamber



Thus it creates a pressure difference between the two chambers where none existed before the demon started its mischief

Thus it creates a pressure difference between the two chambers where none existed before the demon started its mischief

as if the demon could therefore cause entropy to decrease in a system with no consequent increase in entropy anywhere else

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it
- ▶ However, erasing information is irreversible

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it
- ▶ However, erasing information is irreversible
- ▶ Erasing information has an associated increase in entropy

- ▶ Conventional explanation : All observations and measurements require energy consumption and the demon is no exception
- ▶ Modern explanation : demon needs to have a memory to operate, this act of storing information (actually it is erasing information) is associated with an increase of entropy, and this increase cancels out any decrease in entropy that the demon might be able to effect in the system
- ▶ The demon is a computational device that processes and stores information
- ▶ It is possible to design a computational process that proceeds entirely reversibly, and therefore has no increase in entropy associated with it
- ▶ However, erasing information is irreversible
- ▶ Erasing information has an associated increase in entropy
- ▶ Maxwell's demon can operate reversibly therefore, but only if it has a large enough hard disc that it doesn't ever need to clear space to continue operation

- ▶ Connection between Shannon's information entropy and the thermodynamic entropy

- ▶ Connection between Shannon's information entropy and the thermodynamic entropy
- ▶ If a bit is fixed in a certain state, then its entropy = 0

- ▶ Connection between Shannon's information entropy and the thermodynamic entropy
- ▶ If a bit is fixed in a certain state, then its entropy = 0
- ▶ if it is equally likely to be in either of the two states, then entropy = $\ln 2$

- ▶ Connection between Shannon's information entropy and the thermodynamic entropy
- ▶ If a bit is fixed in a certain state, then its entropy= 0
- ▶ if it is equally likely to be in either of the two states, then entropy= $\ln 2$
- ▶ generally, $S = k_B H$, where H =information entropy

The Landauer principle : “information is physical” and it has an energy equivalent

- quantifies the thermodynamic cost of the recording/erasure of one bit of information

The Landauer principle : “information is physical” and it has an energy equivalent

- quantifies the thermodynamic cost of the recording/erasure of one bit of information

- ▶ Erasure of one bit of information requires a minimum energy cost equal to $k_B T \ln 2 \approx 0.018$ eV, where T is the temperature of a thermal reservoir used in the process.

- ▶ Information theory is usually supplied in a form that is independent of any physical embodiment

- ▶ Information theory is usually supplied in a form that is independent of any physical embodiment
- ▶ Landauer argued that information is physical and it has an energy equivalent. It may be stored in physical systems such as books and memory chips and it is transmitted by physical devices exploiting electrical or optical signals.

- ▶ Information theory is usually supplied in a form that is independent of any physical embodiment
- ▶ Landauer argued that information is physical and it has an energy equivalent. It may be stored in physical systems such as books and memory chips and it is transmitted by physical devices exploiting electrical or optical signals.
- ▶ \therefore it must obey the laws of physics and, first and foremost, the laws of thermodynamics.

(1) any logically irreversible process must result in an entropy increase in the non-information-bearing degrees of freedom of the information-processing system or its environment

- (1) any logically irreversible process must result in an entropy increase in the non-information-bearing degrees of freedom of the information-processing system or its environment
- (2) any logically reversible process can be implemented thermodynamically reversibly.

- (1) any logically irreversible process must result in an entropy increase in the non-information-bearing degrees of freedom of the information-processing system or its environment
 - (2) any logically reversible process can be implemented thermodynamically reversibly.
- an amount of energy equal to $k_B T \ln 2$ ($k_B T$ = thermal noise per unit bandwidth) is needed to transmit a bit of information, and more if quantized channels are used with photon energies $h\nu > kT$