

Laplace Approximation

Zeel B Patel, Nipun Batra

August 27, 2023

IIT Gandhinagar

Outline

Taylor Series Expansion
ND Taylor Series
Laplace Approximation



Brook Taylor



Pierre-Simon Laplace

Overall idea

- Posterior distribution $p(\boldsymbol{\theta}|\mathcal{D})$ might be intractable but we can compute the MAP estimate.
- We know that posterior would be in form: $p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z}p(\mathcal{D}, \boldsymbol{\theta})$, where Z is the normalizing constant.
- We can approximate this posterior using Taylor series expansion around the MAP estimate and it turns out that, after making a few assumptions, the resulting distribution is a Gaussian:
 $p(\boldsymbol{\theta}|\mathcal{D}) \approx \mathcal{N}(\boldsymbol{\theta}|\boldsymbol{\theta}_{MAP}, H^{-1})$, where H is the Hessian matrix of the log joint evaluated at $\boldsymbol{\theta}_{MAP}$.

Taylor Series Expansion


1D Taylor Series

$$\tilde{f}(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \frac{f'''(x_0)}{3!}(x - x_0)^3 + \dots$$

Taylor Approximation of a 1D Function


Consider the following function:

$$f(x) = \sin(1 + x)$$




Taylor Approximation of a 1D Function

Taylor approximation at $x_0 = 0$:




Taylor Approximation of a 1D Function

Taylor approximation at $x_0 = 0$:



Taylor Approximation of a 1D Function


Taylor approximation at $x_0 = 0$:



Taylor Approximation of a 1D Function

Taylor approximation at $x_0 = 0$:


$$\tilde{f}(x) = 0.84 + 0.54 \frac{(x-0.00)^1}{1!} - 0.84 \frac{(x-0.00)^2}{2!} - 0.54 \frac{(x-0.00)^3}{3!}$$



Taylor Approximation of a 1D Function

Taylor approximation at $x_0 = 0$:


$$\tilde{f}(x) = 0.84 + 0.54 \frac{(x - 0.00)^1}{1!} - 0.84 \frac{(x - 0.00)^2}{2!} - 0.54 \frac{(x - 0.00)^3}{3!} + 0.84 \frac{(x - 0.00)^4}{4!}$$



Taylor Approximation of a 1D Function

Taylor approximation at $x_0 = 0$:


$$\tilde{f}(x) = 0.84 + 0.54 \frac{(x-0.00)^1}{1!} - 0.84 \frac{(x-0.00)^2}{2!} - 0.54 \frac{(x-0.00)^3}{3!} + 0.84 \frac{(x-0.00)^4}{4!} + 0.54 \frac{(x-0.00)^5}{5!}$$



Taylor Approximation of a 1D Function

Taylor approximation at $x_0 = 0$:


$$\tilde{f}(x) = 0.84 + 0.54 \frac{(x-0.00)^1}{1!} - 0.84 \frac{(x-0.00)^2}{2!} - 0.54 \frac{(x-0.00)^3}{3!} + 0.84 \frac{(x-0.00)^4}{4!} + 0.54 \frac{(x-0.00)^5}{5!} + \dots$$



Taylor Approximation of a 1D Gaussian Function

Consider the standard normal distribution: $p(x) \sim \mathcal{N}(x|0,1)$

$$p(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$



Taylor Approximation of a 1D Gaussian Function

We will try to approximate log of the above function using Taylor series expansion around $x_0 = 0$.

Taylor Approximation of a 1D Gaussian Function

We will try to approximate log of the above function using Taylor series expansion around $x_0 = 0$.

$$f(x) = \log p(x) = \log \left(\frac{1}{\sqrt{2\pi}} \right) - \frac{x^2}{2}$$

Taylor Approximation of a 1D Gaussian Function

We will try to approximate log of the above function using Taylor series expansion around $x_0 = 0$.

$$f(x) = \log p(x) = \log \left(\frac{1}{\sqrt{2\pi}} \right) - \frac{x^2}{2}$$

$$f(0) = \log \left(\frac{1}{\sqrt{2\pi}} \right)$$

Taylor Approximation of a 1D Gaussian Function

We will try to approximate log of the above function using Taylor series expansion around $x_0 = 0$.

$$f(x) = \log p(x) = \log \left(\frac{1}{\sqrt{2\pi}} \right) - \frac{x^2}{2}$$

$$f(0) = \log \left(\frac{1}{\sqrt{2\pi}} \right)$$

$$f'(0) \cdot x = -x_0 \cdot x = 0$$

Taylor Approximation of a 1D Gaussian Function

We will try to approximate log of the above function using Taylor series expansion around $x_0 = 0$.

$$f(x) = \log p(x) = \log \left(\frac{1}{\sqrt{2\pi}} \right) - \frac{x^2}{2}$$

$$f(0) = \log \left(\frac{1}{\sqrt{2\pi}} \right)$$

$$f'(0) \cdot x = -x_0 \cdot x = 0$$

$$f''(0) \cdot \frac{x^2}{2} = -1 \cdot \frac{x^2}{2}$$

Taylor Approximation of a 1D Gaussian Function

We will try to approximate log of the above function using Taylor series expansion around $x_0 = 0$.

$$f(x) = \log p(x) = \log \left(\frac{1}{\sqrt{2\pi}} \right) - \frac{x^2}{2}$$

$$f(0) = \log \left(\frac{1}{\sqrt{2\pi}} \right)$$

$$f'(0) \cdot x = -x_0 \cdot x = 0$$

$$f''(0) \cdot \frac{x^2}{2} = -1 \cdot \frac{x^2}{2}$$

$$\text{Taylor approximated } \tilde{f}(x) = \log \left(\frac{1}{\sqrt{2\pi}} \right) - \frac{x^2}{2}$$


ND Taylor Series

$$\tilde{f}(\mathbf{x}) = f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0)^T (\mathbf{x} - \mathbf{x}_0) + \frac{1}{2} (\mathbf{x} - \mathbf{x}_0)^T \nabla^2 f(\mathbf{x}_0) (\mathbf{x} - \mathbf{x}_0) + \dots$$

Approximate a 2d function

We take the following function:



$$f(x_1, x_2) = \sin(1 + x_1 + x_2)$$



Approximate a 2d function

Taylor approximation at $x_0 = (0, 0)$:



Taylor approximation
Polynomial degree: 1



Approximate a 2d function

Taylor approximation at $x_0 = (0, 0)$:

Taylor approximation
Polynomial degree: 2



Laplace Approximation

$$p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z}p(\mathcal{D}, \boldsymbol{\theta})$$

$$p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z} p(\mathcal{D}, \boldsymbol{\theta})$$

We can rewrite this as:

$$p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z} e^{-f(\boldsymbol{\theta})}$$
$$f(\boldsymbol{\theta}) = -\log p(\mathcal{D}, \boldsymbol{\theta})$$

$$p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z}p(\mathcal{D}, \boldsymbol{\theta})$$

We can rewrite this as:

$$p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z}e^{-f(\boldsymbol{\theta})}$$
$$f(\boldsymbol{\theta}) = -\log p(\mathcal{D}, \boldsymbol{\theta})$$

Note that $f(\boldsymbol{\theta})$ is the negative log joint which is used as a loss function to estimate $\boldsymbol{\theta}_{MAP}$.

- Highest mass is concentrated around θ_{MAP} and hence it makes sense to get Taylor approximation around that point.

Laplace Approximation

- Highest mass is concentrated around θ_{MAP} and hence it makes sense to get Taylor approximation around that point.
- In other words, if our approximation is bad where we have low probability mass, it doesn't matter much.

Laplace Approximation

- Highest mass is concentrated around θ_{MAP} and hence it makes sense to get Taylor approximation around that point.
- In other words, if our approximation is bad where we have low probability mass, it doesn't matter much.
- Thus, we approximate $f(\theta)$ as $\tilde{f}(\theta)$ around θ_{MAP} using Taylor series expansion up to second derivative:

$$\begin{aligned}\tilde{f}(\theta) = & f(\theta_{MAP}) + \nabla f(\theta_{MAP})^T (\theta - \theta_{MAP}) \\ & + \frac{1}{2} (\theta - \theta_{MAP})^T \nabla^2 f(\theta_{MAP}) (\theta - \theta_{MAP})\end{aligned}$$

$$\begin{aligned}\tilde{f}(\boldsymbol{\theta}) &= f(\boldsymbol{\theta}_{MAP}) + \nabla f(\boldsymbol{\theta}_{MAP})^T (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP}) \\ &\quad + \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP}) (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})\end{aligned}$$

Laplace Approximation

$$\begin{aligned}\tilde{f}(\boldsymbol{\theta}) &= f(\boldsymbol{\theta}_{MAP}) + \nabla f(\boldsymbol{\theta}_{MAP})^T (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP}) \\ &\quad + \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP}) (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})\end{aligned}$$

Since, $\boldsymbol{\theta}_{MAP}$ is minima of $f(\boldsymbol{\theta})$, $\nabla f(\boldsymbol{\theta}_{MAP}) = 0$.

Laplace Approximation

$$\begin{aligned}\tilde{f}(\boldsymbol{\theta}) &= f(\boldsymbol{\theta}_{MAP}) + \nabla f(\boldsymbol{\theta}_{MAP})^T (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP}) \\ &\quad + \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP}) (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})\end{aligned}$$

Since, $\boldsymbol{\theta}_{MAP}$ is minima of $f(\boldsymbol{\theta})$, $\nabla f(\boldsymbol{\theta}_{MAP}) = 0$.

$$\tilde{f}(\boldsymbol{\theta}) = f(\boldsymbol{\theta}_{MAP}) + \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP}) (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})$$

Laplace Approximation

$$\begin{aligned}\tilde{f}(\boldsymbol{\theta}) &= f(\boldsymbol{\theta}_{MAP}) + \nabla f(\boldsymbol{\theta}_{MAP})^T (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP}) \\ &\quad + \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP}) (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})\end{aligned}$$

Since, $\boldsymbol{\theta}_{MAP}$ is minima of $f(\boldsymbol{\theta})$, $\nabla f(\boldsymbol{\theta}_{MAP}) = 0$.

$$\tilde{f}(\boldsymbol{\theta}) = f(\boldsymbol{\theta}_{MAP}) + \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP}) (\boldsymbol{\theta} - \boldsymbol{\theta}_{MAP})$$

where $\nabla^2 f(\boldsymbol{\theta}_{MAP})$ is the Hessian matrix of $f(\boldsymbol{\theta})$ evaluated at $\boldsymbol{\theta}_{MAP}$.

Laplace Approximation

Plugging this back to the posterior equation:

$$p(\boldsymbol{\theta}|\mathcal{D}) = \frac{1}{Z} e^{-f(\boldsymbol{\theta})} \quad \text{where } f(\boldsymbol{\theta}) = -\log p(\mathcal{D}, \boldsymbol{\theta})$$

Laplace Approximation

Plugging this back to the posterior equation:

$$\begin{aligned} p(\boldsymbol{\theta}|\mathcal{D}) &= \frac{1}{Z} e^{-f(\boldsymbol{\theta})} \quad \text{where } f(\boldsymbol{\theta}) = -\log p(\mathcal{D}, \boldsymbol{\theta}) \\ &\approx \frac{1}{Z} e^{-f(\boldsymbol{\theta}_{MAP})} e^{-\frac{1}{2}(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP})(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})} \end{aligned}$$

Laplace Approximation

Plugging this back to the posterior equation:

$$\begin{aligned} p(\boldsymbol{\theta}|\mathcal{D}) &= \frac{1}{Z} e^{-f(\boldsymbol{\theta})} \quad \text{where } f(\boldsymbol{\theta}) = -\log p(\mathcal{D}, \boldsymbol{\theta}) \\ &\approx \frac{1}{Z} e^{-f(\boldsymbol{\theta}_{MAP})} e^{-\frac{1}{2}(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP})(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})} \\ &= \frac{1}{Z} p(\mathcal{D}, \boldsymbol{\theta}_{MAP}) e^{-\frac{1}{2}(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP})(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})} \end{aligned}$$

Laplace Approximation

Plugging this back to the posterior equation:

$$\begin{aligned} p(\boldsymbol{\theta}|\mathcal{D}) &= \frac{1}{Z} e^{-f(\boldsymbol{\theta})} \quad \text{where } f(\boldsymbol{\theta}) = -\log p(\mathcal{D}, \boldsymbol{\theta}) \\ &\approx \frac{1}{Z} e^{-f(\boldsymbol{\theta}_{MAP})} e^{-\frac{1}{2}(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP})(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})} \\ &= \frac{1}{Z} p(\mathcal{D}, \boldsymbol{\theta}_{MAP}) e^{-\frac{1}{2}(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})^T \nabla^2 f(\boldsymbol{\theta}_{MAP})(\boldsymbol{\theta}-\boldsymbol{\theta}_{MAP})} \end{aligned}$$

$$\begin{aligned} p(\boldsymbol{\theta}|\mathcal{D}) &\approx \mathcal{N}\left(\boldsymbol{\theta}|\boldsymbol{\theta}_{MAP}, (\nabla^2 f(\boldsymbol{\theta}_{MAP}))^{-1}\right) \\ Z &= p(\mathcal{D}, \boldsymbol{\theta}_{MAP}) \cdot (2\pi)^{D/2} \cdot |\nabla^2 f(\boldsymbol{\theta}_{MAP})|^{-\frac{1}{2}} \end{aligned}$$

Pros and Cons of Laplace Approximation

- Pros:
 - Simple to implement
 - Computationally efficient
 - Can be used to approximate any intractable function

Pros and Cons of Laplace Approximation

- Pros:
 - Simple to implement
 - Computationally efficient
 - Can be used to approximate any intractable function
- Cons:
 - It can give bad approximation when posterior is not unimodal
 - Gaussian assumption can be too restrictive at times
 - Hessian matrix inversion can be numerically unstable and expensive. A diagonal or block-wise approximation can be applied to resolve this. Checkout [Laplace-Redux](#) for more details.

Beta-Bernoulli Coin Toss

Let's take Beta-Bernoulli Coin Toss example since we know the closed form posterior for it. Consider the following scenario:

Beta-Bernoulli Coin Toss


Let's take Beta-Bernoulli Coin Toss example since we know the closed form posterior for it. Consider the following scenario:

- $\mathcal{D} = \{1, 1, 1, 1, 1, 1, 1, 1, 0\}$
- $p(\theta) = \text{Beta}(\alpha = 2, \beta = 2)$
- $h = \theta$
- $p(y|h) = h^y(1 - h)^{1-y}$

Beta-Bernoulli Coin Toss


Let's take Beta-Bernoulli Coin Toss example since we know the closed form posterior for it. Consider the following scenario:

- $\mathcal{D} = \{1, 1, 1, 1, 1, 1, 1, 1, 0\}$
- $p(\theta) = \text{Beta}(\alpha = 2, \beta = 2)$
- $h = \theta$
- $p(y|h) = h^y(1 - h)^{1-y}$




Beta-Bernoulli Coin Toss

MAP estimate:



Beta-Bernoulli Coin Toss


Laplace Approximation:



Normal Prior for Coin Toss


Consider the following coin toss experiment scenario:

- $\mathcal{D} = \{1, 1, 1, 1, 1, 1, 1, 1, 0\}$
- $p(\theta) = \mathcal{N}(\theta | \mu = -2, \sigma = 1)$
- $h = \sigma(\theta)$
- $p(y|\theta) = h^y(1 - h)^{1-y}$




Normal Prior for Coin Toss

First, we find the MAP estimate.



Normal Prior for Coin Toss

Now, according to the Laplace Approximation, the posterior is:



We got True (MC) Posterior by a Monte Carlo estimation of the evidence.

Multi-Mode example

Consider a Gaussian Mixture distribution with two modes. We assume that, it is an unnormalized density and we want to get normalized Laplace approximation of it.

Multi-Mode example


Consider a Gaussian Mixture distribution with two modes. We assume that, it is an unnormalized density and we want to get normalized Laplace approximation of it.

$$p(\theta) = \frac{7}{10}\mathcal{N}(\theta|-2, 1) + \frac{3}{10}\mathcal{N}(\theta|2, 1)$$

Multi-Mode example

Consider a Gaussian Mixture distribution with two modes. We assume that, it is an unnormalized density and we want to get normalized Laplace approximation of it.

$$p(\theta) = \frac{7}{10}\mathcal{N}(\theta|-2, 1) + \frac{3}{10}\mathcal{N}(\theta|2, 1)$$



Multi-Mode example

Laplace Approximation:

