

# TTK4250 Sensor Fusion

## Assignment 2

**Hand in:** *Friday 13. September* on Blackboard.

---

**Read Python setup guide on BB**, it can be found under *Course work*.

This assignment should be handed in on Blackboard, as a PDF pluss **the zip created by running `create_handin.py`**, before the deadline.

You are supposed to show how you got to each answer unless told otherwise.

If you struggle, we encourage you to ask for help from a classmate or come to the exercise class on Friday.

---

### Task 1: *Transformation of Gaussian random variables*

Let  $x \in \mathbb{R}^n$  be  $\mathcal{N}(\mu, \Sigma)$ . Find the distribution and see if you recognize it:

Hint: they are all given in the book.

(a)  $z = \Sigma^{-\frac{1}{2}}(x - \mu)$ , where  $\Sigma^{\frac{1}{2}}(\Sigma^{\frac{1}{2}})^T = \Sigma$

*Hint:* If you are using theorem 2.4.1, you might need  $\det(A^{\frac{1}{2}}) = \det(A)^{\frac{1}{2}}$ ,  $(A^{-1})^T = (A^T)^{-1}$ , and  $\det(A^T) = \det(A)$  whenever  $A$  has full rank.

(b) Use transformation of random variables to find  $y_i = z_i^2$ , where  $z_i$  is the  $i$ 'th variable in the vector  $z$ .

(c)  $y = (x - \mu)^T \Sigma^{-1} (x - \mu) = z^T z = \sum z_i^2 = \sum y_i$ .

*Hint:* The MGF of  $y_i$  is given in the book through example 2.8 and 2.10. Example 2.6 might also be handy.

**Task 2: Sensor fusion**

In this task we want to find out if a boat is above the line  $x_2 = x_1 + 5$ . In order to do this we will fuse measurements from two sensors with our prior belief: A drone-mounted camera, and a maritime surveillance radar. You have some prior knowledge of the state of the boat. You get 1 measurement from each sensor that are processed so that you know them to be (approximately) Gaussian conditioned on the position.

To be more specific, let us denote the state by  $x$  and our prior Gaussian by  $\mathcal{N}(x; \bar{x}, P)$ . The measurement from the camera is given by  $z^c = H^c x + v^c$  and the measurement from the radar by  $z^r = H^r x + v^r$ , where  $v^c, v^r$  denotes the measurement noise and is distributed according to  $\mathcal{N}(0, R^c)$  and  $\mathcal{N}(0, R^r)$ , respectively.

Only insert the numbers when asked to. The needed values are given by

$$\begin{aligned} \bar{x} &= \begin{bmatrix} 0 & 0 \end{bmatrix}^T, & P &= 25I_2, & H^c &= H^r = I_2, \\ R^c &= \begin{bmatrix} 79 & 36 \\ 36 & 36 \end{bmatrix}, & R^r &= \begin{bmatrix} 28 & 4 \\ 4 & 22 \end{bmatrix}, & z^c &= \begin{bmatrix} 2 & 14 \end{bmatrix}^T, & z^r &= \begin{bmatrix} -4 & 6 \end{bmatrix}^T \end{aligned}$$

- (a) What is  $p(z^c|x)$ ?
- (b) Show that the joint  $p(x, z^c)$  can be written as a Gaussian distribution.  

*Hint:* Use conditional probability and the proof of theorem 3.3.1.
- (c) Find the marginal  $p(z^c)$  and the conditional  $p(x|z^c)$ , using the above and either theorems from the book or calculations.
- (d) Given what you found above, what is the marginal  $p(z^r)$  and the conditional  $p(x|z^r)$ ?
- (e) What is the MMSE and MAP estimate of  $x$  given  $z^c$ ? You do not need to do calculations to find the answer, but briefly state what you would do if you had to.
- (f) Finish the `sensor_model.LinearSensorModel2d.get_pred_meas` method that can be used to calculate marginal probabilities  $p(z)$ .
- (g) Finish the `conditioning.get_cond_state` function that can be used to calculate conditional probabilities  $p(x|z)$ .
- (h) Finish the `task2.get_conds` function that is used to calculate the conditional probabilities  $p(x|z^c)$  and  $p(x|z^r)$ .
- (i) Finish the `task2.get_double_conds` function that is used to calculate the conditional probabilities  $p(x|z^c, z^r)$ , i.e. the posterior of  $x$  conditioned on  $z^c$  then  $z^r$ , and  $p(x|z^r, z^c)$ , i.e. the posterior of  $x$  conditioned on  $z^r$  then  $z^c$ . Does it matter which order we condition?
- (j) Finish the `gaussian.MultiVarGauss2d.get_transformed` method that is used to calculate the probability  $p(Tx)$ , where  $T$  is a linear transformation.
- (k) You now want to know the probability that the boat is above the line,  $x_2 = x_1 + 5$ .  
 Finish `task2.get_prob_over_line` using the appropriate linear transform and the CDF.

*Hint:* This is the same as finding  $\Pr(x_2 - x_1 > 5) = \Pr(\begin{bmatrix} -1 & 1 \end{bmatrix} x > 5)$

To get the cdf you can use `from scipy.stats import norm`, and then use

`norm.cdf(value, mean, std)`. Note that it takes the standard deviation (std) and not the variance as input.

**Task 3:** *Working with the canonical form*

In Section 3.3 the fundamental product identity was studied using a moment-based parametrization. Clearly, it must also be possible to establish an equivalent result using the canonical representation. In this exercise we shall therefore consider the product

$$\mathcal{N}^{-1}(\mathbf{x}; \mathbf{a}, \mathbf{B}) \mathcal{N}^{-1}(\mathbf{y}; \mathbf{C}\mathbf{x}, \mathbf{D}). \quad (6)$$

- (a) Show that (6) is identical to the Gaussian

$$\mathcal{N}^{-1}\left(\begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix}; \begin{bmatrix} \mathbf{a} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \mathbf{B} + \mathbf{C}^\top \mathbf{D}^{-1} \mathbf{C} & -\mathbf{C}^\top \\ -\mathbf{C} & \mathbf{D} \end{bmatrix}\right) \quad (7)$$

*Hint:*

Taking the logarithm of the form (3.17) in the book with (3.20) inserted give a relatively simple way to the goal, after the terms constant in  $\mathbf{x}$  and  $\mathbf{y}$  are subtracted. Also

$$\mathbf{a}^\top \mathbf{A} \mathbf{a} + \mathbf{b}^\top \mathbf{B} \mathbf{b} + 2\mathbf{a}^\top \mathbf{C} \mathbf{b} = \mathbf{a}^\top \mathbf{A} \mathbf{a} + \mathbf{b}^\top \mathbf{B} \mathbf{b} + \mathbf{a}^\top \mathbf{C} \mathbf{b} + \mathbf{b}^\top \mathbf{C}^\top \mathbf{a} = \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix}^\top \begin{bmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{C}^\top & \mathbf{B} \end{bmatrix} \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix},$$

is handy, and valid for any vectors  $\mathbf{a}$  and  $\mathbf{b}$  and matrices  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$  of interest (variable names are not related to the task).

- (b) Show that the marginal distribution of  $\mathbf{y}$ , from the joint density (7), is

$$\mathcal{N}^{-1}(\mathbf{y}; \mathbf{C}^\top (\mathbf{B} + \mathbf{C}^\top \mathbf{D}^{-1} \mathbf{C})^{-1} \mathbf{a}, \mathbf{D} - \mathbf{C}(\mathbf{B} + \mathbf{C}^\top \mathbf{D}^{-1} \mathbf{C})^{-1} \mathbf{C}^\top). \quad (8)$$

*Hint:* Theorem 3.4.1

- (c) Show that the conditional distribution of  $\mathbf{x}$  given  $\mathbf{y}$  is

$$\mathcal{N}^{-1}(\mathbf{x}; \mathbf{a} + \mathbf{C}^\top \mathbf{y}, \mathbf{B} + \mathbf{C}^\top \mathbf{D}^{-1} \mathbf{C}). \quad (9)$$

*Hint:* Theorem 3.4.1

- (d) Let us now return to the original formulation of the product identity in Theorem 3.3.1. Use the result from c) to show that

$$\hat{\mathbf{P}}^{-1} = \mathbf{P}^{-1} + \mathbf{H}^\top \mathbf{R}^{-1} \mathbf{H}. \quad (10)$$

*Hint:* Match variables in (3.21) with (3.10) in the book.