

REFERENCES

- [1] J. Schoffelen and J. Gross, "Source connectivity analysis with MEG and EEG," *Human Brain Mapping*, vol. 30, no. 6, pp. 1857–1865, Jun. 2009. [Online]. Available: <https://onlinelibrary.wiley.com/doi/10.1002/hbm.20745>
- [2] A. Coito, M. Genetti, F. Pittau, G. R. Iannotti, A. Thomschewski, Y. H"oller, E. Trinka, R. Wiest, M. Seeck, C. M. Michel, G. Plomp, and S. Vulliemoz, "Altered directed functional connectivity in temporal lobe epilepsy in the absence of interictal spikes: A high density EEG study," *Epilepsia*, vol. 57, no. 3, pp. 402–411, Mar. 2016. [Online]. Available: <https://onlinelibrary.wiley.com/doi/10.1111/epi.13308>
- [3] S. Pelle, A. Scarabello, L. Ferri, G. Ricci, F. Bisulli, and M. Ursino, "Enhancing non-invasive pre-surgical evaluation through functional connectivity and graph theory in drugresistant focal epilepsy," *Journal of Neuroscience Methods*, vol. 413, p. 110300, Jan. 2025. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S0165027024002450>
- [4] H. Becker, L. Albera, P. Comon, R. Gribonval, F. Wendling, and I. Merlet, "Brain-source imaging: From sparse to tensor models," *IEEE Signal Processing Magazine*, vol. 32, no. 6, pp. 100–112, 2015.
- [5] W. Xie, R. T. Toll, and C. A. Nelson, "EEG functional connectivity analysis in the source space," *Developmental Cognitive Neuroscience*, vol. 56, p. 101119, Aug. 2022. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S1878929322000627>
- [6] F. Tadel, S. Baillet, J. C. Mosher, D. Pantazis, and R. M. Leahy, "Brainstorm: A User-Friendly Application for MEG/EEG Analysis," *Computational Intelligence and Neuroscience*, vol. 2011, pp. 1–13, 2011. [Online]. Available: <http://www.hindawi.com/journals/cin/2011/879716/>
- [7] E. K"oksai-Ers"oz, M. Yochum, P. Benquet, and F. Wendling, "eCOALIA: Neocortical neural mass model for simulating electroencephalographic signals," *SoftwareX*, vol. 28, p. 101924, Dec. 2024.
- [8] R. S. Desikan, F. S'egonne, B. Fischl, B. T. Quinn, B. C. Dickerson, D. Blacker, R. L. Buckner, A. M. Dale, R. P. Maguire, B. T. Hyman, M. S. Albert, and R. J. Killiany, "An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest," *NeuroImage*, vol. 31, no. 3, pp. 968–980, Jul. 2006. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S1053811906000437>
- [9] N. A. Anstey, "CORRELATION TECHNIQUES –A REVIEW*," *Geophysical Prospecting*, vol. 12, no. 4, pp. 355–382, Dec. 1964. [Online]. Available: <https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2478.1964.tb01911.x>
- [10] R. Srinivasan, W. R. Winter, J. Ding, and P. L. Nunez, "EEG and MEG coherence: Measures of functional connectivity at distinct spatial scales of neocortical dynamics," *Journal of Neuroscience Methods*, vol. 166, no. 1, pp. 41–52, Oct. 2007. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S016502700700307X>
- [11] J. F. Hipp, D. J. Hawellek, M. Corbetta, M. Siegel, and A. K. Engel, "Large-scale cortical correlation structure of spontaneous oscillatory activity," *Nature Neuroscience*, vol. 15, no. 6, pp. 884–890, Jun. 2012. [Online]. Available: <https://www.nature.com/articles/nn.3101>
- [12] R. Bru"na, F. Maest'u, and E. Pereda, "Phase locking value revisited: teaching new tricks to an old dog," *Journal of Neural Engineering*, vol. 15, no. 5, p. 056011, Oct. 2018. [Online]. Available: <https://iopscience.iop.org/article/10.1088/1741-2552/aacfe4>
- [13] M. S. H"am"al"ainen and R. J. Ilmoniemi, "Interpreting magnetic fields of the brain: minimum norm estimates," *Medical & Biological Engineering & Computing*, vol. 32, no. 1, pp. 35–42, Jan. 1994. [Online]. Available: <http://link.springer.com/10.1007/BF02512476>