

Limitations of Basing Screening Policies on Screening Trials

The US Preventive Services Task Force and Prostate Cancer Screening

Ruth Etzioni, PhD,* Roman Gulati, MS,* Matt R. Cooperberg, MD,† David M. Penson, MD,‡
Noel S. Weiss, PhD,§ and Ian M. Thompson, MD||

Background: The US Preventive Services Task Force recently recommended against prostate-specific antigen screening for prostate cancer based primarily on evidence from the European Randomized Study of Screening for Prostate Cancer (ERSPC) and the US Prostate, Lung, Colorectal, and Ovarian (PLCO) cancer screening trial.

Objective: To examine limitations of basing screening policy on evidence from screening trials.

Methods: We reviewed published modeling studies that examined population and trial data. The studies (1) project the roles of screening and changes in primary treatment in the US mortality decline; (2) extrapolate the ERSPC mortality reduction to the long-term US setting; (3) estimate overdiagnosis based on US incidence trends; and (4) quantify the impact of control arm screening on PLCO mortality results.

Results: Screening plausibly explains 45% and changes in primary treatment can explain 33% of the US prostate cancer mortality decline. Extrapolating the ERSPC results to the long-term US setting implies an absolute mortality reduction at least 5 times greater than that observed in the trial. Approximately 28% of screen-detected cases are overdiagnosed in the United States versus 58% of screen-detected cases suggested by the ERSPC results. Control arm screening can explain the null result in the PLCO trial.

Conclusions: Modeling studies indicate that population trends and trial results extended to the long-term population setting are consistent with greater benefit of prostate-specific antigen screening—and more favorable harm-benefit tradeoffs—than has been suggested by empirical trial evidence.

Key Words: mass screening, policy development, prostatic neoplasms, simulation modeling

(*Med Care* 2013;51: 295–300)

Recently, the US Preventive Services Task Force recommended against prostate-specific antigen (PSA) screening for prostate cancer based on moderate certainty that the benefits of screening do not outweigh the harms.¹ The evidence of screening benefit was based primarily on mortality results from the European Randomized Study of Screening for Prostate Cancer (ERSPC)^{2,3} and the US-based Prostate, Lung, Colorectal, and Ovarian (PLCO) cancer screening trial.^{4,5} Restating the trial mortality results after medians of 13 and 11 years of follow-up, respectively, the Task Force concluded that “there is adequate evidence that the benefit of PSA screening and early treatment ranges from 0 to 1 prostate cancer deaths avoided per 1000 men screened.”¹

Although randomized trials are the gold standard for evidence-based decision making in medicine, they can have important limitations as a basis for screening policies. First, screening policy development demands information about long-term benefits and harms because these policies generally pertain to interventions conducted over an individual’s healthy lifetime. Unfortunately, most screening trials provide short-term results rather than the long-term outcomes generated by a typical population-based screening program. Second, screening trial results can be highly influenced by the trial population and by patterns of compliance with the trial protocol. Third, any inferences about screening benefit are limited to the screening strategy or strategies tested in the trial. This does not permit policy makers to identify and compare alternative policies that might be more acceptable.

In this commentary, we argue that the ERSPC and PLCO are subject to these limitations and that taking their results at face value misrepresents the likely long-term population impact of PSA screening (relative to no screening) in the United States. To examine these limitations, we review modeling studies that analyze population trends and extrapolate trial evidence. A recent publication defines the models used in these studies as “mathematical frameworks that facilitate estimation of the consequences of health care decisions.”⁶ These models enable in-depth analysis of observed data that can reveal information about the disease

From the *Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA; †Department of Urology, UCSF Helen Diller Family Comprehensive Cancer Center, San Francisco, CA; ‡Vanderbilt University Medical Center, Nashville, TN; §Department of Epidemiology, University of Washington, Seattle, WA; and ||Department of Urology, University of Texas, San Antonio, TX.

Supported by Award Numbers R01 CA131874 and U01 CA88160 from the National Cancer Institute and Award Number U01 CA157224 from the National Cancer Institute and the Centers for Disease Control.

The authors declare no conflict of interest.

Reprints: Ruth Etzioni, PhD, Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N, M2-B230, Seattle, WA 98109-1024. E-mail: retzioni@fhcrc.org.

Copyright © 2013 by Lippincott Williams & Wilkins
ISSN: 0025-7079/13/5104-0295

process and facilitate extrapolation beyond the trial setting and horizon.

We summarize 3 main points from these studies. First, changes in primary treatments explain only a minority of the observed decline in prostate cancer mortality; assuming screening benefits would-be metastatic cases detected while still in a localized stage, PSA screening plausibly explains nearly half of the decline. Second, estimates of overdiagnoses and lives saved based on ERSPC results overstate the likely long-term harm-benefit tradeoff in the United States. Third, results from the PLCO trial, which found no significant mortality difference between the intervention and control arms, do not rule out a clinically significant benefit of PSA screening; instead, the key lesson of the PLCO trial is that more intensive screening does not necessarily yield a greater benefit than less intensive screening. This observation opens the door to investigations of more efficient screening strategies that preserve benefit while reducing cost and harms, and we can use modeling to identify such strategies.

Our discussion draws on several models developed as part of the Cancer Intervention and Surveillance Modeling Network. To model population mortality trends, we review a microsimulation model of prostate cancer progression, detection, and survival that projects mortality with and without screening or changes in treatment. We examine the same microsimulation model and an analytic model to estimate overdiagnosis associated with PSA screening from population incidence trends. To extrapolate long-term harm-benefit tradeoffs, we combine the estimated number of overdiagnosed lives with the estimated lives saved because of screening, based on applying the relative reduction in prostate cancer deaths from the ERSPC to long-term US prostate cancer mortality without screening. Finally, to interrogate PLCO results, we review a microsimulation model to replicate the PLCO trial with and without control arm contamination under a clinically significant screening benefit.

We conclude that model-based analysis of evidence is consistent with an important long-term benefit of PSA screening and that policies should be developed that improve the targeting of screening and treatment rather than eliminating the opportunity for early detection and intervention. Although we focus on the case of prostate cancer screening, our arguments have bearing on the methods by which screening policies are developed, particularly when the only available trial evidence has clear limitations and extrapolation beyond the specific trial settings, horizons, and protocols is needed.

POPULATION OUTCOMES

From the time of rapid adoption of PSA screening in the United States in the early 1990s, prostate cancer death rates have fallen each year. The latest figures released by the Surveillance, Epidemiology, and End Results program for 2009 indicate that age-adjusted prostate cancer death rates have declined by 44% since their peak in 1991.⁷ However, other aspects of prostate cancer management have also

changed both before and during the PSA era. Primary treatment patterns have evolved considerably, beginning with a marked increase in the frequency of radical prostatectomy in the 1980s,⁸ followed by the dissemination of hormonal therapies as adjuvant to radiation therapy in the 1990s.⁹ Both of these treatments have been shown to be efficacious in randomized treatment trials,^{10,11} although recent results from a US trial of radical prostatectomy have suggested that, in a population undergoing screening, benefit may be limited to men with intermediate or high-risk disease.¹² In addition, radiation therapy technologies have evolved and are now able to deliver more intense, targeted doses. Advances in primary treatment have been cited¹³ as the most likely explanation for the decline in prostate cancer deaths in the United States. Nonetheless, 3 independently developed simulation models of prostate cancer progression, diagnosis, and survival,¹⁴ making favorable assumptions regarding treatment efficacy based on the results of randomized trials and recent comparative effectiveness studies,^{15,16} estimated that primary treatment changes explained only up to one third of the drop in disease-specific deaths¹⁴ by 2005, leaving a large fraction of the decline to be explained by other factors.

One of the 3 models¹⁷ was used to study the role of screening in the absence of treatment changes using data through 2000 and found that screening explained 45% of the mortality decline.¹⁸ The model was rigorously calibrated to US prostate cancer incidence trends before and after the introduction of screening¹⁷ and assumed that cases detected in an earlier (localized) stage by screening received a corresponding post-lead-time survival improvement. This screening benefit assumption was supported by a later analysis in which the same model was used to simulate the ERSPC and this assumption was found to be consistent with the 27% drop in prostate cancer deaths observed after a median follow-up of 9 years (after adjustment for non-attendance).¹⁹

Although these results do not prove that screening explains the observed drop in prostate cancer deaths, a decline of 70% in the incidence of advanced stage disease since 1990 supports a role for PSA screening in explaining the part of the mortality decline that is not accounted for by changes in primary treatment.²⁰ A direct connection between PSA screening and the incidence of metastatic disease is evident from the ERSPC, in which the incidence of metastatic disease in the screened group (278 cases) was about half that in the control group (567 cases).³ Although there may be other explanations for the drop in US prostate cancer deaths, these are not clearly supported by available data. For example, there have been improvements in treatment for metastatic disease, but these have been relatively recent, and a study of trends in survival among newly diagnosed metastatic cases shows little change over time through 2005.²¹ Earlier detection of progressive disease using PSA monitoring could also be a factor, but to date there have been no conclusive studies of the efficacy of earlier versus later treatment of metastatic disease following diagnosis. No etiologic factors that might have induced a change in the underlying risk of developing or dying of the disease in the population have been identified. It has been suggested that misattribution of

other-cause deaths as due to prostate cancer could have caused a spurious rise in deaths due to the disease just before the peak in 1991, but studies of misattribution have indicated that it is a relatively rare event.^{22,23} We conclude that, although population prostate cancer trends are likely a complex result of many factors, modeling can help to disentangle the contributions of specific factors like treatment and screening. The observed decline in disease-specific deaths is not easily explained by other changes in disease management and is indicative of screening benefit in the population.

TRIAL RESULTS I: THE ERSPC

The ERSPC observed a short-term relative reduction in deaths (fraction of disease-specific deaths prevented by screening) of 21% after a median follow-up of 11 years³ and a corresponding absolute reduction (lives saved per 1000 men enrolled in screening) of 1.07 deaths per 1000 men invited. It is this measure of absolute benefit that formed the basis for the Task Force's conclusion that the reduction in prostate cancer mortality 10–14 years after PSA screening is “at most, very small.”¹

How can such a large relative effect of screening translate into such a modest absolute benefit and, ultimately, into an impression of an ineffective screening test?²⁴ This is where the issues of follow-up duration (time from start of enrollment to analysis) and trial population (population from which trial participants are sampled) become essential to understand.

The absolute benefit of screening depends on the baseline disease-specific mortality rate in the unscreened population. This rate was extremely low in the ERSPC control group, amounting to only 0.5 deaths per 1000 person years after a median of 11 years of follow-up.³ Even though the 21% relative reduction was substantial, the absolute difference in rates amounted to only 0.1 deaths per 1000 person years and this translated into 1.07 lives saved per 1000 men screened.³

However, in a population screening program, where men begin testing at a specified starting age and continue screening until they either die of other causes or reach an age beyond which screening is no longer recommended, the relevant baseline disease-specific mortality approaches the lifetime probability of prostate cancer death. In addition, this lifetime probability should apply to the population of interest. The ERSPC results not only reflect short follow-up; they also pertain to a European population. In the United States, in the absence of screening, the lifetime risk of a prostate cancer death based on 1990 death rates, just before the spread of PSA screening, was 32 per 1000 men. A reduction of 21% yields 6.7 ($=32 \times 0.21$) lives saved per 1000 men screened and an number needed to screen (NNS) of 149 ($=1000/6.7$). Recognizing that advances in treatment may have mitigated prostate cancer mortality, and using more recent mortality rates from 2006, we obtain a lifetime risk of prostate cancer death of 28 per 1000 men, 5.9 lives saved per 1000 men screened, and an NNS of 170. Using endpoints of the 95% confidence interval for ERSPC mortality reduction implies a corresponding range of 2.5 ($=28 \times 0.09$) to 9.0

($=28 \times 0.32$) lives saved per 1000 men screened and a range of NNS of 112–397. These NNS estimates are considerably lower than the short-term estimates from the trial (1410 and 1055 at 9 and 11 y, respectively).

The short-term perspective also distorts estimates of screening harm, particularly overdiagnosis. This is the detection by screening of cancers that would never have become clinically apparent without the screening test. Because prostate cancer is known to be a disease that is highly prevalent, particularly in older men, the potential for overdiagnosis is great. Indeed, after a median follow-up of 9 years, the frequency of overdiagnosis among men invited to screening was estimated to be 34 per 1000 men screened, which was the observed excess incidence in the screened group relative to the control group.² Combining this estimated frequency of overdiagnosis with the 9-year estimate of 0.7 lives saved per 1000 men screened yielded a projected ratio of 48 overdiagnoses per life saved, or 48 additional cancers needed to detect (NND) to save 1 life.² This figure has been cited in the media as an illustration of the high harm-to-benefit tradeoff associated with PSA screening.^{25,26} However, the excess incidence over the short term produces an inflated assessment of overdiagnosis amounting in this case to 58% of screen-detected cases (>1 case overdiagnosed for every 2 screen-detected cancers). After 11 years, the NND was revised down to 37, but both overdiagnosis and the corresponding NND are highly dependent on the setting and length of follow-up.³

Two different models were used to estimate the long-term fraction of screen-detected cases overdiagnosed in the US population.^{27,28} In contrast to the trial-based estimates, which use excess incidence in the screened group as a proxy for overdiagnosis, both models estimated the lead time associated with PSA screening then derived the fraction overdiagnosed as the fraction of screen-detected cases dying of other causes within their lead time. The lead-time-based estimates, namely, 23% and 28% overdiagnosed among screen-detected cases, are lower than the ERSPC estimate because they represent true overdiagnosis rather than excess incidence and because they were estimated in the US setting. Given that 16% of men will be diagnosed under current screening practices,²⁹ then even assuming that all new cases are screen detected and using the higher estimate of overdiagnosis, we project that 44.8 ($=0.16 \times 0.28 \times 1000$) per 1000 will be overdiagnosed during their lifetimes. With 5.9 lives saved per 1000 over the long-term based on US population mortality, the long-term NND is more accurately estimated as 7.6 ($=44.8/5.9$) rather than 37, giving a very different picture of the harm-benefit profile of PSA screening.

TRIAL RESULTS II: THE PLCO

The PLCO trial was originally designed to compare annual screening with no screening for prostate cancer. However, the trial was initiated in 1993, at the tail of the first wave of PSA screening in the US population.²⁹ As a consequence, screening was common among control arm subjects; an investigation into control arm contamination

revealed “the intensity of PSA screening in the control arm was estimated to be approximately half of that in the intervention arm.”⁵ In fact, trial investigators estimated that 74% of men in the control arm received at least 1 routine PSA test during the 6 years of the trial compared with 95% of the intervention arm, with over half of the men in the control tested every 1–2 years. Moreover, the incidence of prostate cancer in the control group was 20% higher than expected based on comparable population rates, suggesting that control arm subjects were screened even more intensively than the general population.³⁰

The PLCO trial has been interpreted as providing evidence that screening is not beneficial.¹³ Indeed, the Task Force’s estimate of the mortality reduction associated with screening includes the possibility of 0 lives saved over a 10- to 14-year period based on the trial result of no reduction in prostate cancer deaths in the intervention group relative to the control group. In fact, both published reports from the trial^{4,5} have shown a slight excess of prostate cancer deaths in the intervention group. When modeling was used to replicate the trial, it was found that the trial has extremely low power (range, 9%–25% across three models at 13 y) to infer a difference between the control and intervention groups even under a clinically significant benefit of screening.¹⁹ This is due not only to the control arm contamination but also to the lower-than-expected frequency of prostate cancer deaths in the trial population.³¹ Given the low numbers of deaths, the mortality results are highly variable, and the chance of excess mortality in the intervention arm, even in the presence of a significant screening benefit, is 15%–29% after 13 years.¹⁹

We conclude that the PLCO does not provide actionable information regarding screening benefit or lack thereof and that inclusion by the Task Force of 0 as a potential lower bound for the benefit of screening on the basis of the reported PLCO trial results is not warranted. Instead, the study provides important evidence of the equivalence of more intensive (annual) versus less intensive (approximately biennial) screening.

CONCLUSIONS

Clinical trials are generally perceived to reflect the highest level of evidence for policy development as they are designed to eliminate systematic differences between the treatment and control groups that may lead to biased results. There have been many nontrial-based studies of PSA screening efficacy,^{32–38} but these have either been inconclusive or subject to extensive criticism because of their observational nature.

We are in agreement that published results from clinical trials should be used to inform policy so long as they are directly applicable to the policy-relevant setting. In the case of prostate screening and the Task Force recommendations, this setting is the US population over the long term. We have made the case that the published screening trial results do not accurately reflect the outcomes of most import in this setting. Thus, our primary concern relates to relevance of the published trial results, and our goal is in essence to make these

results more relevant for policy. In doing so, we consider the published results to be reliable; we do not address issues of data quality although we are aware that some have questioned the quality and reporting of the ERSPC findings.^{13,39}

There is a further handicap associated with limiting policy development to published trials: inferences about screening harm and benefit will pertain only to those strategies actually tested. If there are strategies with more favorable harm-benefit tradeoffs than those tested, these cannot be considered or evaluated. This is critically important in the case of prostate cancer screening with its high price tag in terms of unnecessary biopsies, which can cause serious infections including sepsis in 0.5%–1.0% of biopsy cases, and overtreatment, which carries the burden of impotence and incontinence in 20%–30% of cases.⁴⁰

There have been many suggestions for targeted screening strategies, including increasing the interval between screens,⁴¹ using more conservative criteria for biopsy referral in older men who are at higher risk of false-positive tests and overdiagnosis,⁴² and screening adaptively, such as lengthening the interval to the next screen if the current PSA level is below a specified threshold.⁴³ Testing all competing policies in randomized trials is infeasible; consequently, this is a setting in which well-calibrated and validated models are proving to be of great value.^{44,45} We therefore advocate the use of models alongside clinical trial results, not only to interpret and extrapolate trial findings to the relevant population setting, but also to ensure that a recommendation against screening takes into account feasible alternative strategies.

The universe of models is massive and heterogeneous in terms of both model quality and methodology. Models have many limitations. They invariably represent a simplified version of disease progression, intervention, and impact. The accuracy of model results is dependent on the underlying assumptions of the model. Results themselves are subject to uncertainty which can be difficult to quantify. A recently published series of reports from a joint Task Force of the International Society for Pharmaceutical Outcomes Research and the Society for Medical Decision Making^{6,46–52} codifies good modeling practices and should help to ensure that these limitations are acknowledged and addressed to the extent possible. However, the question of how to weigh the evidence from different data sources and study designs, including models, when formulating screening recommendations remains. Task Force recommendations for breast and colorectal screening relied on both trials and modeling studies, yet a systematic framework for synthesizing evidence from multiple study designs has yet to be established.

In conclusion, we submit these comments out of a sense that a critically important policy decision may have been made on the basis of an incomplete picture of the benefits and harms of PSA screening. We consider this to reflect a weakness not of the specific decision—a more complete picture may have led the panel to the same recommendation. Rather our concern is about the process—a process in which the most reliable evidence is defined as the observed outcome of clinical trials and there is no formal mandate to go beyond published trial estimates of benefit

and harm. With a disease whose hallmark is a lengthy natural history, the harms of developing cancer screening policies based primarily on limited-duration screening trials may well outweigh the benefits.

REFERENCES

- Moyer VA. On behalf of the USPSTF. Screening for Prostate Cancer: US Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2012;157:120–134.
- Schröder FH, Hugosson J, Roobol MJ, et al. Screening and prostate-cancer mortality in a randomized European study. *N Engl J Med.* 2009;360:1320–1328.
- Schröder FH, Hugosson J, Roobol MJ, et al. Prostate-cancer mortality at 11 years of follow-up. *N Engl J Med.* 2012;366:981–990.
- Andriole GL, Crawford ED, Grubb RL 3rd, et al. Mortality results from a randomized prostate-cancer screening trial. *N Engl J Med.* 2009;360:1310–1319.
- Andriole GL, Crawford ED, Grubb RL III, et al. Prostate cancer screening in the randomized prostate, lung, colorectal, and ovarian cancer screening trial: mortality results after 13 years of follow-up. *J Natl Cancer Inst.* 2012;104:1–8.
- Caro JJ, Briggs AH, Siebert U, et al. Modeling Good Research Practices—Overview: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-1. *Med Decis Making.* 2012;32:667–677.
- Surveillance, Epidemiology, and End Results (SEER) Program. Available at: <http://www.seer.cancer.gov>. Mortality-All COD, Public-Use With State, Total US (1969-2005), National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released April 2008. Underlying mortality data provided by NCHS. Available at: <http://www.cdc.gov/nchs>. Accessed October 15, 2012.
- Lu-Yao GL, Greenberg ER. Changes in prostate cancer incidence and treatment in USA. *Lancet.* 1994;343:251–254.
- Cooperberg MR, Grossfeld GD, Lubeck DP, et al. National practice patterns and time trends in androgen ablation for localized prostate cancer. *J Natl Cancer Inst.* 2003;95:981–989.
- Bolla M, Gonzalez D, Warde P, et al. Improved survival in patients with locally advanced prostate cancer treated with radiotherapy and goserelin. *N Engl J Med.* 1997;337:295–300.
- Bill-Axelsson A, Holmberg L, Ruutu M, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. *N Engl J Med.* 2011;364:1708–1717.
- Wilt T, Brawer MK, Jones K, et al. Radical prostatectomy versus observation for localized prostate cancer. *N Engl J Med.* 2012;367:203–213.
- Chou R, LeFevre ML. Prostate cancer screening—the evidence, the recommendations, and the clinical implications. *JAMA.* 2011;306:2721–2722.
- Etzioni R, Gulati R, Tsodikov A, et al. The prostate cancer conundrum revisited: treatment changes and prostate cancer mortality declines. *Cancer.* 2012;118:5955–5963.
- Cooperberg MR, Vickers AJ, Broering JM, et al. Comparative risk-adjusted mortality outcomes after primary surgery, radiotherapy, or androgen-deprivation therapy for localized prostate cancer. *Cancer.* 2010;116:5226–5234.
- Boorjian SA, Kames RJ, Viterbo R, et al. Long-term survival after radical prostatectomy versus external-beam radiotherapy for patients with high-risk prostate cancer. *Cancer.* 2011;117:2883–2891.
- Gulati R, Inoue L, Katcher J, et al. Calibrating disease progression models using population data: a critical precursor to policy development in cancer control. *Biostatistics.* 2010;11:707–719.
- Etzioni R, Tsodikov A, Mariotto A, et al. Quantifying the role of PSA screening in the US prostate cancer mortality decline. *Cancer Causes Control.* 2008;19:175–181.
- Gulati R, Tsodikov A, Wever EM, et al. The impact of PLCO control arm contamination on perceived PSA screening efficacy. *Cancer Causes Control.* 2012;23:827–835.
- Surveillance, Epidemiology, and End Results (SEER) Program. Available at: <http://www.seer.cancer.gov>. Incidence—SEER 9 Regs Research Data, Nov 2011 Sub, Vintage 2009 Pops (1973-2009) <Katrina/Rita Population Adjustment> - Linked To County Attributes—Total US, 1969-2010 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2012, based on the November 2011 submission. Accessed May 1, 2012.
- Cetin K, Beebe-Dimmer JL, Fryzek JP, et al. Recent time trends in the epidemiology of stage IV prostate cancer in the United States: analysis of data from the Surveillance, Epidemiology, and End Results Program. *Urology.* 2010;75:1396–1404.
- Penson DF, Albertsen PC, Nelson PS, et al. Determining cause of death in prostate cancer: are death certificates valid? *J Natl Cancer Inst.* 2001;93:1822–1823.
- Hoffman RM, Stone SN, Hunt WC, et al. Effects of misattribution in assigning cause of death on prostate cancer mortality rates. *Ann Epidemiol.* 2003;13:450–454.
- Hanley JA. Mortality reductions produced by sustained prostate cancer screening have been underestimated. *J Med Screen.* 2010;17:147–151.
- Kolata G. *Studies Show Prostate Test Saves Few Lives.* New York, NY: New York Times; 2009.
- Ablin R. *The Great Prostate Mistake.* Tuscon, AZ: New York Times; 2010.
- Telesca D, Etzioni R, Gulati R. Estimating lead time and overdiagnosis associated with PSA screening from prostate cancer incidence trends. *Biometrics.* 2008;64:10–19.
- Draisma G, Etzioni R, Tsodikov A, et al. Lead time and overdiagnosis in prostate-specific antigen screening: importance of methods and context. *J Natl Cancer Inst.* 2009;101:374–383.
- Mariotto A, Etzioni R, Krapcho M, et al. Reconstructing prostate-specific antigen (PSA) testing patterns among black and white men in the US from Medicare claims and the National Health Interview Survey. *Cancer.* 2007;109:1877–1886.
- Pinsky PF, Black A, Kramer BS, et al. Assessing contamination and compliance in the prostate component of the Prostate, Lung, Colorectal, and Ovarian (PLCO) Cancer Screening Trial. *Clin Trials.* 2010;7:303–311.
- Prorok PC, Andriole GL, Bresalier RS, et al. Design of the Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial. *Control Clin Trials.* 2000;21:273S–309S.
- Farkas A, Schneider D, Perrotti M, et al. National trends in the epidemiology of prostate cancer, 1973 to 1994: evidence for the effectiveness of prostate-specific antigen screening. *Urology.* 1998;52:444–448; discussion 448–449.
- Shibata A, Ma J, Whittemore AS. Prostate cancer incidence and mortality in the United States and the United Kingdom. *J Natl Cancer Inst.* 1998;90:1230–1231.
- Lu-Yao G, Albertsen PC, Stanford JL, et al. Natural experiment examining impact of aggressive screening and treatment on prostate cancer mortality in two fixed cohorts from Seattle area and Connecticut. *BMJ.* 2002;325:740.
- Chu KC, Tarone RE, Freeman HP. Trends in prostate cancer mortality among black men and white men in the United States. *Cancer.* 2003;97:1507–1516.
- Coldman AJ, Phillips N, Pickles TA. Trends in prostate cancer incidence and mortality: an analysis of mortality change by screening intensity. *CMAJ.* 2003;168:31–35.
- Baade PD, Coory MD, Aitken JF. International trends in prostate-cancer mortality: the decrease is continuing and spreading. *Cancer Causes Control.* 2004;15:237–241.
- Concato J, Wells CK, Horwitz RI, et al. The effectiveness of screening for prostate cancer: A nested case-control study. *Arch Intern Med.* 2006;166:38–43.
- Miller AB. New data on prostate-cancer mortality after PSA screening. *N Engl J Med.* 2012;366:1047–1048.
- Chou R, Crosswell JM, Dana T, et al. Screening for prostate cancer: a review of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med.* 2011;155:762–771.
- Wolf AM, Wender RC, Etzioni RB, et al. American Cancer Society guideline for the early detection of prostate cancer: update 2010. *CA Cancer J Clin.* 2010;60:70–98.
- Polascik TJ, Oesterling JE, Partin AW. Prostate specific antigen: a decade of discovery—what we have learned and where we are going. *J Urol.* 1999;162:293–306.

43. Vickers AJ, Lilja H. Urological cancer: time for another rethink on prostate cancer screening. *Nat Rev Clin Oncol*. 2012;9:7–8.
44. Zauber AG, Lansdorp-Vogelaar I, Knudsen AB, et al. Evaluating test strategies for colorectal cancer screening: a decision analysis for the US Preventive Services Task Force. *Ann Intern Med*. 2008;149:659–669.
45. Mandelblatt JS, Cronin KA, Bailey S, et al. Effects of mammography screening under different screening schedules: model estimates of potential benefits and harms. *Ann Intern Med*. 2009;151:738–747.
46. ISPOR-SMDM Joint Modeling Good Research Practices Task Force. Available at: <http://www.ispor.org/TaskForces/GRPModelingTf.asp>. Accessed May 6, 2012.
47. Roberts M, Russell LB, Paltiel AD, et al. Conceptualizing a Model: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-2. *Value Health*. 2012;15:804–811.
48. Siebert U, Alagoz O, Bayoumi AM, et al. State-Transition Modeling: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-3. *Value Health*. 2012;15:812–820.
49. Karnon J, Stahl J, Brennan A, et al. Modeling Using Discrete Event Simulation: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-4. *Med Decis Making*. 2012;32:701–711.
50. Pitman R, Fisman D, Zaric GS, et al. Dynamic Transmission Modeling: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-5. *Value Health*. 2012;15:828–834.
51. Briggs AH, Weinstein MC, Fenwick EA, et al. Model Parameter Estimation and Uncertainty Analysis: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force Working Group-6. *Med Decis Making*. 2012;32:722–732.
52. Eddy DM, Hollingworth W, Caro JJ, et al. Model Transparency and Validation: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-7. *Med Decis Making*. 2012;32:733–743.