

# Toward a Unified Framework for Quantum-Relativistic Systems: A Higher-Categorical and Noncommutative Geometric Formulation

## Abstract

We present a conceptual framework that integrates quantum mechanics and gravitational phenomena using advanced mathematical tools. This approach combines higher category theory, Topological Quantum Field Theory (TQFT), derived functors, homotopy-theoretic methods, noncommutative geometry, spectral theory, category-theoretic logic, and topological data analysis. Rather than a single equation or conventional model, we outline a meta-structure in which quantum states, spacetime configurations, and their mutual interactions are represented in an  $(\infty, 1)$ -categorical setting. The noncommutative geometric input encodes curvature into a quantum Hamiltonian through commutators of covariant derivatives. TQFT connects topological changes in spacetime to transformations of quantum states, while derived functors and homotopy theory ensure stable invariants under deformation. Category-theoretic logic provides a foundational language for observables and measurements, and topological data analysis supplies robust topological invariants of evolving state spaces. We conclude with a prototype equation that synthesizes these elements, serving as a starting point for a more complete unified theory of quantum gravity.

## 1 Introduction

A consistent unification of quantum mechanics and general relativity remains one of the central open problems in theoretical physics. Quantum field theories describe matter and non-gravitational forces with great precision, but when it comes to spacetime itself, classical concepts of geometry and the dynamical curvature described by general relativity pose formidable conceptual and mathematical challenges.

To move toward a unified framework, we must consider new mathematical tools. Higher category theory and TQFT shed light on how global topological features of spacetime might influence quantum behavior. Derived functors and homotopy theory reveal stable invariants under deformations, addressing anomalies and singular structures in a controlled manner. Noncommutative geometry replaces classical point-set notions with operator-algebraic structures, making curvature emerge naturally as a noncommutative spectral invariant. Spectral theory then links geometric data to the eigenvalues of certain operators, while category-theoretic logic provides a logical and conceptual blueprint for understanding observables and measurements. Finally, topological data analysis (TDA) offers a data-driven approach to extracting robust topological features from evolving quantum states.

In this paper, we outline a conceptual scaffolding that weaves together these advanced mathematical domains. Rather than proposing a final theory, we present a flexible and extensible framework that shows how each component can fit into a broader unifying picture. We conclude with a prototype equation that incorporates these ideas into a single symbolic expression.

## 2 Higher-Categorical and TQFT Structures

Consider an  $(\infty, 1)$ -category of spacetime configurations, denoted

$$(\infty, \mathbf{1})\text{-Spacetime}.$$

Its objects can represent geometric configurations or possibly noncommutative analogs of manifolds, while its morphisms represent transformations of these configurations (e.g., changes in topology or boundary conditions). Higher morphisms (2-, 3-, and so forth) encode gauge transformations, dualities, and higher symmetries.

We also consider an  $(\infty, 1)$ -category of quantum states,

$$(\infty, \mathbf{1})\text{-QStates},$$

whose objects are not just Hilbert spaces but possibly derived objects such as chain complexes of vector spaces or Hilbert spaces equipped with additional structure.

TQFT provides a functorial bridge:

$$Z : (\infty, \mathbf{1})\text{-Cobordisms} \rightarrow \mathbf{Vect},$$

assigning topological data (cobordisms) to vector spaces and linear maps. Integrating  $Z$  with

our geometric and quantum categories allows topological changes in spacetime to induce well-defined transformations in the quantum state categories, capturing global, topological aspects of quantum-gravitational behavior.

### 3 Derived Functors and Homotopy Theory

The complexities of quantum gravity often involve singularities, anomalies, and nontrivial topological features. Derived functors allow us to handle these subtleties systematically. We introduce a derived functor that connects geometry to states:

$$\mathcal{H}^{(\infty)} : (\infty, \mathbf{1})\text{-Spacetime} \xrightarrow{\text{derived}} (\infty, \mathbf{1})\text{-QStates}.$$

This functor ensures that as we move through different spacetime configurations, the corresponding quantum states incorporate homotopy-invariant and cohomological corrections. Homotopy theory stabilizes the physical content against local deformations, focusing attention on robust invariants rather than fragile, coordinate-dependent details.

### 4 Noncommutative Geometry and Spectral Theory

A key step toward unification is the recognition that spacetime at the Planck scale may not be described by classical geometry. Instead, we adopt a noncommutative geometric perspective, where points of spacetime are replaced by elements of a noncommutative algebra. The curvature of spacetime emerges from commutators of covariant derivatives  $D_\mu$ :

$$[D_\mu, D_\nu].$$

This leads to a noncommutative generalization of the Hamiltonian:

$$\mathcal{H}(t) = \frac{\hbar c}{l_p^2} [D_\mu, D_\nu].$$

Its spectral decomposition provides eigenvalues linked to geometric invariants, connecting the energy spectrum of quantum states to underlying geometric and topological data.

## 5 Category-Theoretic Logic and Observables

Quantum observables can be represented as objects in a suitable topos, and their logical relationships (e.g., implications, consistency conditions) become morphisms in a category of propositions. This categorical logic transforms questions about measurement and uncertainty into statements about morphisms and subobject classifiers, integrating quantum observables into the higher-categorical fabric of the theory.

## 6 Topological Data Analysis of Evolving States

As parameters such as time or curvature change, quantum states trace out trajectories in high-dimensional spaces. TDA techniques like persistent homology identify stable topological features in these evolving data sets. By integrating TDA, we discover topological invariants that persist across parameter changes, offering new insights into phase structures and stable features of quantum-gravitational states.

## 7 A Prototype Unifying Equation

While a fully unified theory is more complex than any single expression, we propose a guiding equation as a starting point:

$$\frac{d}{dt}\Psi(t) = \frac{\hbar c}{l_p^2}[D_\mu, D_\nu]\Psi(t) \oplus Z(\text{Cobordisms}) \oplus \delta_{\text{derived}}(\Psi(t)).$$

Here:

- $\Psi(t)$  is an object in  $(\infty, \mathbf{1})\text{-QStates}$ , capturing a quantum state enriched with topological and homotopy-theoretic structure.
- $\frac{\hbar c}{l_p^2}[D_\mu, D_\nu]$  encodes noncommutative geometric curvature contributions, tying space-time structure to quantum evolution.
- $Z(\text{Cobordisms})$  represents topological transitions in spacetime as functorial assignments from TQFT, altering the state space in line with topological changes.
- $\delta_{\text{derived}}(\Psi(t))$  indicates derived and homotopy-theoretic corrections, ensuring stable invariants and refining the evolution beyond what a simple operator could capture.

The use of  $\oplus$  is schematic, indicating that the effective “Hamiltonian” is a composite of geometric, topological, and derived corrections. This equation is a conceptual placeholder, illustrating how contributions from various advanced frameworks might be combined.

## 8 Conclusion

This paper outlines a conceptual meta-structure for unifying quantum mechanics with gravitational phenomena. By leveraging higher category theory, TQFT, derived functors, homotopy theory, noncommutative geometry, spectral theory, categorical logic, and TDA, we gain a flexible language and toolkit for tackling quantum gravity. While much work remains to refine and concretize this framework, the approach presented here suggests a path forward, where complexity is not shunned but embraced as a signpost to deeper, more foundational understanding.