Subgradient Descent

Xintian Han & David S. Rosenberg

NYU CDS

Feb 13, 2019

Contents

1. Motivation and Review: Lasso
2. Convexity and Sublevel Sets
3. Convex and Differentiable Functions
4. Subgradients
5. Subgradient Descent

Motivation and Review: Lasso

The Lasso Problem

- Lasso problem can be parametrized as

$$\min_{w \in \mathbf{R}^d} J(w) = \frac{1}{n} \sum_{i=1}^{n} \left\{ w^T x_i - y_i \right\}^2 + \lambda \|w\|_1$$

- We could solve Lasso by Shooting Method and Projected SGD.
- How about using SGD?
- $\|w\|_1 = | w_1 | + | w_2 |$ is not differentiable!

Gradient Descent on Lasso Objective?

- The partial gradient of the Lasso objective is

$$\nabla_w J(w) = \frac{1}{n}\sum_{j=1}^{n} 2\{w^T x_j - y_j\}x_j + \lambda \cdot \text{sign}(w)$$

  when $w_i \neq 0$ for all i, and **otherwise is undefined**.

Potential arguments for why we shouldn't care about the points of nondifferentiability:

- If we start with a random w, will we ever hit exactly $w_i = 0$?
- If we did, could we perturb the step size by ε to miss such a point?
- Does it even make sense to check $w_i = 0$ with floating point numbers?

Gradient Descent on Lasso Objective?

- If we blindly apply gradient descent from a random starting point
  - seems unlikely that we'll hit a point where the gradient is undefined.
  - So it's clear how we could apply gradient descent or SGD to a function that's not differentiable everywhere
- Still, doesn't mean that gradient descent will work if objective not differentiable!
- Theory of subgradients and subgradient descent will clear up any uncertainty.

Convexity and Sublevel Sets

Convex Sets

Definition

A set C is **convex** if the line segment between any two points in C lies in C.

KPM Fig. 7.4

Convex and Concave Functions

Definition

A function $f : \mathbf{R}^d \to \mathbf{R}$ is **convex** if the line segment connecting any two points on the graph of f lies above the graph. f is **concave** if $-f$ is convex.

KPM Fig. 7.5

Examples of Convex Functions on $\mathbf{R}$

Examples

- $x \mapsto ax + b$ is both convex and concave on $\mathbf{R}$ for all $a, b \in \mathbf{R}$.
- $x \mapsto |x|^p$ for $p \geqslant 1$ is convex on $\mathbf{R}$
- $x \mapsto e^{ax}$ is convex on $\mathbf{R}$ for all $a \in \mathbf{R}$
- Every norm on $\mathbf{R}^n$ is convex (e.g. $\|x\|_1$ and $\|x\|_2$)
- Max: $(x_1, \ldots, x_n) \mapsto \max\{x_1 \ldots, x_n\}$ is convex on $\mathbf{R}^n$

Simple Composition Rules

Examples

- If g is convex, and $Ax + b$ is an affine mapping, then $g(Ax + b)$ is convex.
- If g is convex then $\exp g(x)$ is convex.
- If g is convex and nonnegative and $p \geqslant 1$ then $g(x)^p$ is convex.
- If g is concave and positive then $\log g(x)$ is concave
- If g is concave and positive then $1/g(x)$ is convex.

Main Reference for Convex Optimization

- Boyd and Vandenberghe (2004)
  - Very clearly written, but has a ton of detail for a first pass.
  - See the Extreme Abridgement of Boyd and Vandenberghe.

Level Sets and Sublevel Sets

Let $f : \mathbf{R}^d \to \mathbf{R}$ be a function. Then we have the following definitions:

Definition

A **level set** or **contour line** for the value c is the set of points $x \in \mathbf{R}^d$ for which $f(x) = c$.

Definition

A **sublevel** set for the value c is the set of points $x \in \mathbf{R}^d$ for which $f(x) \leqslant c$.

Theorem

*If $f : \mathbf{R}^d \to \mathbf{R}$ is **convex**, then the **sublevel sets are convex**.*

(Proof straight from definitions.)

Convex Function

Plot courtesy of Brett Bernstein.

Contour Plot Convex Function: Sublevel Set

Is the sublevel set $\{x \mid f(x) \leqslant 1\}$ convex?

Plot courtesy of Brett Bernstein.

Nonconvex Function

Plot courtesy of Brett Bernstein.

Contour Plot Nonconvex Function: Sublevel Set

Is the sublevel set $\{x \mid f(x) \leqslant 1\}$ convex?

Plot courtesy of Brett Bernstein.

Fact: Intersection of Convex Sets is Convex

Plot courtesy of Brett Bernstein.

Level and Superlevel Sets

Level sets and superlevel sets of convex functions are **not** generally convex.

Plot courtesy of Brett Bernstein.

Convex Optimization Problem: Standard Form

Convex Optimization Problem: Standard Form

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leqslant 0, \quad i = 1, \ldots, m \end{array}$$

where $f_0, \ldots, f_m$ are convex functions.

- What can we say about each constraint set $\{x \mid f_i(x) \leqslant 0\}$? (convex)
- What can we say about the feasible set $\{x \mid f_i(x) \leqslant 0, i = 1, \ldots, m\}$? (convex)

Convex Optimization Problem: Implicit Form

Convex Optimization Problem: Implicit Form

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & x \in C \end{array}$$

where f is a convex function and C is a convex set.
An alternative "generic" convex optimization problem.

Convex and Differentiable Functions

First-Order Approximation

- Suppose $f : \mathbf{R}^d \to \mathbf{R}$ is **differentiable.**
- Predict $f(y)$ given $f(x)$ and $\nabla f(x)$?
- Linear (i.e. "**first order**") approximation:

$$f(y) \approx f(x) + \nabla f(x)^T (y - x)$$

Boyd & Vandenberghe Fig. 3.2

First-Order Condition for Convex, Differentiable Function

- Suppose $f : \mathbf{R}^d \to \mathbf{R}$ is **convex** and **differentiable.**
- Then for any $x, y \in \mathbf{R}^d$

$$f(y) \geqslant f(x) + \nabla f(x)^T (y - x)$$

- The linear approximation to f at x is a **global underestimator** of f:

Figure from Boyd & Vandenberghe Fig. 3.2; Proof in Section 3.1.3

First-Order Condition for Convex, Differentiable Function

- Suppose $f : \mathbf{R}^d \to \mathbf{R}$ is **convex** and **differentiable**
- Then for any $x, y \in \mathbf{R}^d$

$$f(y) \geqslant f(x) + \nabla f(x)^T (y - x)$$

Corollary

If $\nabla f(x) = 0$ then x is a global minimizer of f.

For convex functions, **local information gives global information.**

Subgradients

Subgradients

Definition

A vector $g \in \mathbf{R}^d$ is a **subgradient** of $f : \mathbf{R}^d \to \mathbf{R}$ at x if for all z,

$$f(z) \geqslant f(x) + g^T(z - x).$$

Blue is a graph of $f(x)$.
Each red line $x \mapsto f(x_0) + g^T(x - x_0)$ is a global lower bound on $f(x)$.

Subdifferential

Definitions

- f is **subdifferentiable** at x if $\exists$ at least one subgradient at x.
- The set of all subgradients at x is called the **subdifferential:** $\partial f(x)$

Basic Facts

- f is convex and differentiable $\implies \partial f(x) = \{\nabla f(x)\}$.
- Any point x, there can be 0, 1, or infinitely many subgradients.
- $\partial f(x) = \emptyset \implies f$ is not convex.

Globla Optimality Condition

Definition

A vector $g \in \mathbf{R}^d$ is a **subgradient** of $f : \mathbf{R}^d \to \mathbf{R}$ at x if for all z,

$$f(z) \geqslant f(x) + g^T(z - x).$$

Corollary

If $0 \in \partial f(x)$, *then* x *is a* ***global minimizer*** *of* f.

Subdifferential of Absolute Value

- Consider $f(x) = |x|$

- Plot on right shows $\{(x, g) \mid x \in \mathbf{R},\ g \in \partial f(x)\}$

Boyd EE364b: Subgradients Slides

$f(x_1, x_2) = |x_1| + 2|x_2|$

Plot courtesy of Brett Bernstein.

Subgradients of $f(x_1, x_2) = |x_1| + 2|x_2|$

- Let's find the subdifferential of $f(x_1, x_2) = |x_1| + 2|x_2|$ at $(3, 0)$.
- First coordinate of subgradient must be 1, from $|x_1|$ part (at $x_1 = 3$).
- Second coordinate of subgradient can be anything in $[-2, 2]$.
- So graph of $h(x_1, x_2) = f(3, 0) + g^T(x_1 - 3, x_2 - 0)$ is a global underestimate of $f(x_1, x_2)$, for any $g = (g_1, g_2)$, where $g_1 = 1$ and $g_2 \in [-2, 2]$.

Underestimating Hyperplane to $f(x_1, x_2) = |x_1| + 2|x_2|$

Plot courtesy of Brett Bernstein.

Important Properties of Subdifferential

- If $f_1, \ldots, f_m : \mathbf{R}^d \to \mathbf{R}$ are convex functions and $f = f_1 + \cdots + f_m$, then $\partial f(x) = \partial f_1(x) + \cdots + \partial f_m(x)$.
- For $\alpha \geqslant 0$, $\partial(\alpha f)(x) = \alpha \partial f(x)$.

Subgradients of $f(x) = \|x\|_1$

- Let's find the subdifferential of $f(x) = \|x\|_1 = \sum_{i=1}^{d} |x_i|$ at any given point $x^0 = (x_1^0, x_2^0, \ldots, x_d^0)$.
- By an important property of subdifferential: If $f = f_1 + \cdots + f_m$, then $\partial f(x) = \partial f_1(x) + \cdots \partial f_m(x)$.
- We could calculate the subgradient of $f^i(x) = |x_i|$ and sum them up.
- The subgradient $g^i = (g_1^i, \ldots, g_d^i)$ of $f^i(x) = |x_i|$ at $x^0 = (x_1^0, x_2^0, \ldots, x_d^0)$ is:

$$g_j^i = 0, \quad j \neq i; \quad g_j^i = s(x_j^0), \quad j = i,$$

  where $s(x) = \text{sign}(x)$ if $x \neq 0$ and $s(x) \in [-1, 1]$ if $x = 0$
- We sum all the g^i up to get the subgradient $g = (g_1, \ldots, g_d)$ of $f(x)$ at x^0:

$$g_i = s(x_i^0) \quad \text{for all } i$$

Subgradient Descent

Subgradient Descent

- Suppose f is convex, and we start optimizing at x_0.
- Repeat
  - Step in a negative subgradient direction:

$$x = x_0 - tg,$$

    where $t > 0$ is the step size and $g \in \partial f(x_0)$.

- $-g$ not a descent direction – can this work?

Convergence Theorem for Fixed Step Size

Assume $f : \mathbf{R}^n \to \mathbf{R}$ is convex and

- f is Lipschitz continuous with constant $G > 0$:

$$|f(x) - f(y)| \leqslant G\|x - y\| \text{ for all } x, y$$

Theorem

For fixed step size t, subgradient method satisfies:

$$\lim_{k \to \infty} f(x_{best}^{(k)}) \leqslant f(x^*) + G^2 t/2$$

Based on https://www.cs.cmu.edu/~ggordon/10725-F12/slides/06-sg-method.pdf

Convergence Theorems for Decreasing Step Sizes

Assume $f : \mathbf{R}^n \to \mathbf{R}$ is convex and

- f is Lipschitz continuous with constant $G > 0$:

$$|f(x) - f(y)| \leqslant G\|x - y\| \text{ for all } x, y$$

Theorem

For step size respecting Robbins-Monro conditions,

$$\lim_{k \to \infty} f(x_{best}^{(k)}) = f(x^*)$$

Based on https://www.cs.cmu.edu/~ggordon/10725-F12/slides/06-sg-method.pdf

Subgradient Descent for Lasso Problem

- Lasso problem can be parametrized as

$$\min_{w \in \mathbf{R}^d} J(w) = \frac{1}{n} \sum_{i=1}^{n} \left\{ w^T x_i - y_i \right\}^2 + \lambda \|w\|_1$$

- Subgradients of $J(w)$ are

$$\frac{1}{n} \sum_{i=1}^{n} 2\{w^T x_i - y_i\} x_i + \lambda s,$$

where $s_i = \text{sign}(w_i)$ if $w_i \neq 0$ and $s_i \in [-1, 1]$ if $w_i = 0$.

Subgradient Descent for Lasso Problem: Potential Issues

- Subgradient descent will work for all convex and Lipschitz continuous objective functions.
- BUT, convergence can be very **slow** for non-differentiable functions
- One can often find better approaches by closer examination of the objective function. For example, shooting method or projected SGD.
- Taking small steps in the direction of the (sub)gradient usually may **not** lead to zero coordinates.
- BUT, in practice, we can threshold small values.

Appendix: Subgradient Gets Us Closer To Minimizer

Theorem

Suppose f is convex.

- *Let $x = x_0 - tg$, for $g \in \partial f(x_0)$.*
- *Let z be any point for which $f(z) < f(x_0)$.*
- *Then for small enough $t > 0$,*

$$\|x - z\|_2 < \|x_0 - z\|_2.$$

- Apply this with $z = x^* \in \arg\min_x f(x)$.

$\Longrightarrow$ **Appendix: Negative subgradient step gets us closer to minimizer**.

Subgradient Gets Us Closer To Minimizer (Proof)

- Let $x = x_0 - tg$, for $g \in \partial f(x_0)$ and $t > 0$.
- Let z be any point for which $f(z) < f(x_0)$.
- Then

$$\begin{aligned} \|x - z\|_2^2 &= \|x_0 - tg - z\|_2^2 \\ &= \|x_0 - z\|_2^2 - 2tg^T(x_0 - z) + t^2\|g\|_2^2 \\ &\leqslant \|x_0 - z\|_2^2 - 2t[f(x_0) - f(z)] + t^2\|g\|_2^2 \end{aligned}$$

- Consider $-2t[f(x_0) - f(z)] + t^2\|g\|_2^2$.
  - It's a convex quadratic (facing upwards).
  - Has zeros at $t = 0$ and $t = 2(f(x_0) - f(z))/\|g\|_2^2 > 0$.
  - Therefore, it's negative for any

$$t \in \left(0, \frac{2(f(x_0) - f(z))}{\|g\|_2^2}\right).$$

Based on Boyd EE364b: Subgradients Slides