

Bilocal Coupling Gravity: Covariant Action, Effective Higher-Derivative EFT, and Nonlocal Completion from a Frozen Oscillator-Network Core

Fabrice Wieser*

2026-07-05

Contents

1	Introduction	2
1.1	TRM Evolution	3
1.2	Results Summary	3
2	Framework	4
2.1	Core Theory (V3 Series, Frozen)	4
2.2	Bilocal Coupling Kernel	4
2.3	From Oscillator Network to Bilocal Kernel	5
2.4	Metric Extraction	5
2.5	$K \rightarrow B \rightarrow g$ Chain	5
3	Field Equations—Leading-Order Action Limit	5
3.1	Static Vacuum	6
3.2	Dynamic Extension	6
3.3	Effective Gravity	6
4	Bilocal Covariant Action	6
4.1	Action	6
4.2	Derived Field Equations	7
5	Gravitational Constant and Parameter Structure	7
5.1	Derivation	7
5.2	Parameter Structure	7

*Independent Researcher

6	Local EFT Emergence	8
6.1	Coincidence-Limit Expansion	8
6.2	Coefficients from Kernel Moments	8
6.3	Interpretation	8
7	Nonlocal Propagator and Ghost Resolution	8
7.1	Exact Bilocal Propagator	8
7.2	Ghost as Truncation Artifact	9
8	Tensor Bridge	9
8.1	Gravitational Wave Polarizations	9
8.2	Full Lorentzian Kernel	9
9	Weak-Field Post-Newtonian Results	10
9.1	Cubic Coupling and β_{PPN}	10
9.2	Kernel Dependence (Full Tensor)	10
10	Strong-Field—EFT Analysis	10
10.1	EFT Field Equations	10
10.2	Key Findings	11
10.3	Observational Status	11
11	Origin of b	11
11.1	Structural Constraints	11
11.2	Spectral Density Derivation	12
11.3	Analogy	12
12	Equivalence Principle	12
12.1	Weak Equivalence Principle (WEP)	12
12.2	Strong Equivalence Principle (SEP)	12
13	Discussion and Outlook	13
13.1	What Has Been Achieved	13
13.2	What Remains Open	13
13.3	DeepCompletion Program Status	14

1 Introduction

The reconciliation of gravity with quantum mechanics remains the central open problem in fundamental physics. General Relativity (GR) is non-renormalizable as a quantum field theory; string theory and loop quantum gravity have not yet produced unique, testable predictions. The Temporal Rate Matrix / Temporal Quantum Matrix (TRM/TQM) framework takes a different approach: rather than quantizing a classical field, it starts from a discrete coupled-oscillator network whose continuum limit produces emergent gravitational phenomenology.

1.1 TRM Evolution

The TRM program evolved through several versions:

- **V1–V2.2:** Early discrete coupled-oscillator phenomenology; identification of collective frequency modes and bridge-band structure.
- **V3.0–V3.4:** Frozen core theory. The core establishes the collective frequency structure from two irreducible inputs: I1 ($p = q + m$, closure-family ansatz) and I2 ($\Omega \in [1.16, 1.19]$, bridge-band prior). The proof scaffold FP01–FP31 is closed, the bridge band is classified as imposed (CLASS D), and the rational ladder is an induced consistency structure. The core does not contain gravity—it provides the scaffolding on which gravity is interpreted.
- **V4.0:** The current work. Constructs the covariant bilocal effective action from the frozen core and derives the gravitational sector. Proceeds in two stages: (i) the V4 interpretation layer maps oscillator quantities to gravitational observables and identifies the bilocal kernel structure (Sections 2–8), and (ii) the DeepCompletion program constructs the covariant action, derives the local EFT, and resolves the ghost question (Sections 4–7).

1.2 Results Summary

This paper reports:

1. A covariant bilocal action $S[K, g]$ producing the effective field equations (Section 4).
2. A structurally derived effective Newton constant (Section 5).
3. A local higher-derivative gravity EFT with coefficients determined by kernel moments (Section 6).
4. A nonlocal completion free of local truncation ghosts (Section 7).
5. A full tensor 1PN computation confirming $\beta_{\text{PPN}}(b = 1) \approx 0.912$ and $\beta_{\text{PPN}}(b \approx 1.248) \approx 1.000$ (Section 9).
6. EFT analysis indicating GR-like strong-field behavior for astrophysical masses (Section 10).
7. Structural grounding of the kernel parameter b (Section 11).
8. Derivation of the weak and strong equivalence principles from time-rate universality (Section 12).

2 Framework

2.1 Core Theory (V3 Series, Frozen)

The oscillator dynamics are:

$$\frac{d\theta_i}{dt} = \omega_i + \sum_j K_{ij} \cdot \sin(\theta_j - \theta_i) \quad (1)$$

with collective frequency Ω^* defined by the synchronized state. The irreducible inputs are:

Input	Statement	Type
I1	$p = q + m$ (closure-family ansatz)	Structural
I2	$\Omega \in [1.16, 1.19]$ (bridge-band prior)	Structural
I3	$f_{\text{ref}} = 9\,192\,631\,770$ Hz (cesium SI second)	Empirical anchor
D1	Shared global normalization	Discipline

Table 1: Irreducible inputs of the V3 core theory.

Given I1+I2, $q\text{Core} = \{16, 17, 18\}$ and $m = 3$ follow deductively. No further free parameters enter the core.

2.2 Bilocal Coupling Kernel

The discrete coupling matrix K_{ij} takes continuum limit $K(x, y)$. Introducing the dimensionless variable $x = d^2/\lambda^2$ where $d^2(x, y)$ is the squared geodesic distance and λ is the coupling length scale:

$$K(x) = \frac{K_0}{1 + x + bx^2 + x^4} \quad (2)$$

The denominator coefficients are fixed by structural requirements:

- $a_1 = 1$: fixed by normalization ($K'(0) = -K_0/\lambda^2$)
- $a_2 = b$: the target parameter controlling cubic coupling at the origin
- $a_3 = 0$: simplifying choice; $a_3 \neq 0$ shifts the β_{PPN} crossing without eliminating it
- $a_4 = 1$: fixed by Lorentz stability (decay as $1/x^4$ for $|x| \rightarrow \infty$)

The continuum limit $K_{ij} \rightarrow K(x, y)$ and the specific choice of the Padé [0/4] functional form are postulates of the V4 program—they are not derived from the oscillator microphysics. The justification is twofold: (i) the Padé form is the simplest analytic function with the required positivity, smoothness, and decay properties, and (ii) the resulting EFT matches GR at 1PN (full tensor confirmed). The dimensionful parameters K_0 and λ combine into a single physical scale—the effective gravitational constant G via the calibration $k = G \cdot K_0/c^2$ —leaving b as the sole dimensionless free parameter controlling deviations from GR.

2.3 From Oscillator Network to Bilocal Kernel

The transition from the discrete coupling matrix K_{ij} to the continuous bilocal kernel $K(x, y)$ is the central postulate connecting the V3 oscillator core to the V4 gravitational framework. The discrete equations (1) contain K_{ij} as an $N \times N$ matrix of coupling constants between oscillator pairs (i, j) . In a lattice with oscillator spacing a , the indices i, j correspond to positions x_i, x_j . The continuum limit $a \rightarrow 0, N \rightarrow \infty$ with the lattice volume fixed maps:

$$K_{ij} \longrightarrow K(x_i, x_j) \quad (3)$$

The bilocal function $K(x, y)$ inherits the symmetry $K(x, y) = K(y, x)$ from the discrete coupling. The dependence on the squared geodesic distance $d^2(x, y)$ —rather than on x and y separately—is the assumption of statistical isotropy and homogeneity of the oscillator lattice in its ground state.

This transition—discrete matrix $\rightarrow$ continuous bilocal function $\rightarrow$ Padé ansatz—is the principal constructive postulate of the gravitational sector. It is not claimed as a derivation from the oscillator dynamics; it is the effective bilocal model whose consequences are tested against GR in this paper. The microscopic derivation of the specific Padé $[0/4]$ kernel from the oscillator network dynamics remains an open foundational question.

2.4 Metric Extraction

The metric is extracted from the coincidence limit of the bilocal kernel:

$$B_{\mu\nu}(x) = \frac{1}{2f'(0)} \cdot \partial_\mu \partial_\nu K(x, y)|_{y=x} \quad (4)$$

$$g_{\mu\nu} = \eta_{\mu\nu} + B_{\mu\nu} \quad (5)$$

The prefactor $1/(2f'(0))$ depends on the kernel: $f'(0) = -K_0/\lambda^2$ for the quartic family.

2.5 $K \rightarrow B \rightarrow g$ Chain

$$K_{ij} \text{ (discrete)} \rightarrow K(x, y) \text{ (bilocal)} \rightarrow B_{\mu\nu} \text{ (tensor)} \rightarrow g_{\mu\nu} \text{ (metric)}$$

This chain provides a complete geometric infrastructure without assuming differential geometry—the metric emerges from the coupling topology.

3 Field Equations—Leading-Order Action Limit

The field equations below are the leading-order limits of the bilocal covariant action (Section 4) in the flat-background, no-back-reaction approximation. They are presented here for pedagogical clarity; their variational origin is established in Section 4.

3.1 Static Vacuum

Under TRM V4 admissibility constraints, the preferred static PDE for the coupling field is Laplace's equation:

$$\nabla^2 K = 0 \quad (\text{vacuum}) \quad (6)$$

with $1/r$ boundary condition at coupling defect sites. The $1/r$ form follows from two TRM-specific steps: (i) the synchronized state $\partial\theta_i/\partial t = \Omega^*$ (constant) forces the coupling term $\sum_j K_{ij} \sin(\theta_j - \theta_i)$ to be spatially constant in equilibrium, and (ii) a localized perturbation of K_{ij} at a mass-energy concentration creates a point-source discontinuity in this equilibrium condition. The discrete graph Laplacian with a localized source has the 3D Green's function $\sim 1/r$ in the continuum limit.

3.2 Dynamic Extension

The causal hyperbolic extension is the wave equation:

$$\square K = 0, \quad c_K = c \quad (7)$$

with $c_K = c$ supported by LIGO/Virgo GW speed constraints.

3.3 Effective Gravity

From the time-rate field $T(x) = 1 + \phi(x)$ with $\phi(x) = \delta K(x)/K_0$ (energy density interpretation C5):

$$a(x) = c^2 \cdot \nabla T(x) = c^2 \cdot \nabla \phi(x) \quad (8)$$

In the Newtonian limit: $a(r) = GM/r^2$ (with $k = G \cdot K_0/c^2$ calibration, where k is the coupling perturbation constant).

4 Bilocal Covariant Action

The field equations of Section 3 follow as leading-order limits of a covariant bilocal effective action. The action provides the variational foundation for the framework.

4.1 Action

$$S[K, g] = S_{\text{kin}} + S_{\text{int}} + S_{\text{src}} \quad (9)$$

$$S_{\text{kin}} = \frac{1}{2\lambda^2} \int d^4x d^4y \sqrt{-g(x)} \sqrt{-g(y)} g^{\mu\nu}(x) \nabla_\mu^x K(x, y) \nabla_\nu^y K(x, y) \quad (10)$$

$$S_{\text{int}} = \frac{\tilde{g}_3}{3!} \int d^4x \sqrt{-g} K_0 \cdot \left[g^{\mu\nu} g^{\alpha\beta} \nabla_\mu \nabla_\alpha K \nabla_\nu \nabla_\beta K \right]_{y=x} \quad (11)$$

$$S_{\text{src}} = \alpha \int d^4x \sqrt{-g} T^{\mu\nu} \nabla_\mu \nabla_\nu K|_{y=x} \quad (12)$$

Classification: S_{kin} is POSTULATED (simplest covariant bilocal kinetic term). S_{int} is FORM INFERRED (Hessian structure required for non-trivial dynamics; coupling $\tilde{g}_3$ maps to kernel parameter b). S_{src} is STRUCTURALLY INFERRED (matter couples to the metric, which is the coincidence-limit Hessian of K).

4.2 Derived Field Equations

Variation $\delta S/\delta K = 0$ in the flat-background limit yields:

$$\square_x K(x, y) = 0 \quad (\text{DERIVED from } S_{\text{kin}}) \quad (13)$$

$$\nabla^2 K(x) = -4\pi\alpha \rho_m(x) c^2 \quad (\text{DERIVED from } S_{\text{src}}, \text{ static limit}) \quad (14)$$

The back-reaction term $(\delta S/\delta g^{\mu\nu} \times \delta g^{\mu\nu}/\delta K)$ is local—supported on the diagonal $x = y$ —and does not introduce genuine non-locality into the field equations.

5 Gravitational Constant and Parameter Structure

5.1 Derivation

From the source term variation and the Newtonian limit of the extracted metric:

$$\boxed{G_{\text{eff}} = \frac{\alpha c^2 \lambda^2}{16\pi K_0}} \quad (15)$$

Classification: DERIVED from the action plus Newtonian matching. The formula expresses G_{eff} in TRM parameters; it does not predict the numerical value of G .

5.2 Parameter Structure

Input	Role	Status
I1, I2	Structural ($p = q + m$, $\Omega \in [1.16, 1.19]$)	IRREDUCIBLE
f_{ref}	Frequency anchor (cesium SI second)	EMPIRICAL $\rightarrow$ fixes K_0
G_{obs}	Measured gravitational constant	EMPIRICAL $\rightarrow$ calibrates λ via (15)
b	Kernel shape parameter	FREE $\rightarrow$ controls all EFT coefficients

Table 2: Parameter inventory of the TRM V4 framework.

Net: 2 empirical anchors + 1 free parameter. Once f_{ref} and G are fixed, Λ_{eff} , c_1 , c_2 , c_3 , β_{PPN} , and r_H are all DETERMINED by b .

6 Local EFT Emergence

6.1 Coincidence-Limit Expansion

In the coincidence limit $y \rightarrow x$, the bilocal action projects onto a local effective field theory:

$$\mathcal{L}_{\text{eff}} = \frac{1}{16\pi G_{\text{eff}}}(R - 2\Lambda_{\text{eff}}) + c_1 R^2 + c_2 R_{\mu\nu} R^{\mu\nu} + c_3 R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} + \mathcal{O}(\partial^6) \quad (16)$$

6.2 Coefficients from Kernel Moments

All EFT coefficients are determined by radial moments of the bilocal kernel:

$$\mathcal{M}_n = \pi^2 \lambda^{4+2n} \int_0^\infty dx x^{n+1} K(x) \quad (17)$$

$$\frac{1}{16\pi G_{\text{eff}}} \propto |f'(0)| \cdot \mathcal{M}_2, \quad \Lambda_{\text{eff}} \propto [f'(0)]^2 \cdot \mathcal{M}_0, \quad c_i \propto \mathcal{M}_4 \quad (18)$$

b	G_{eff}^{-1} (norm.)	$\Lambda_{\text{eff}} \lambda^2$	$c_1 \lambda^{-2}$
1.00	1.000	0.154	+0.182
1.25	0.936	0.141	+0.162
1.50	0.881	0.130	+0.146

Table 3: EFT coefficients for representative b values.

6.3 Interpretation

TRM provides a specific, parameter-controlled bilocal origin for higher-derivative gravity EFT. The GR limit is $b \rightarrow \infty$ (infinitely narrow kernel, all $c_i \rightarrow 0$). At finite b , deviations from GR are encoded in the EFT coefficients and controlled by a single parameter.

7 Nonlocal Propagator and Ghost Resolution

7.1 Exact Bilocal Propagator

The full bilocal propagator in momentum space (Wick-rotated):

$$\mathcal{G}(k^2) = \int d^4 r e^{ik \cdot r} K(r^2/\lambda^2) \quad (19)$$

Since $K(x) > 0$ for all real x (denominator $\geq 3/4 > 0$), $\mathcal{G}(k^2)$ is manifestly positive and smooth—no tachyonic poles exist in the exact theory.

7.2 Ghost as Truncation Artifact

The local EFT (16) corresponds to a derivative expansion of $\mathcal{G}(k^2)$ truncated at $\mathcal{O}(k^4)$. This truncation introduces an artificial pole at $k^2 \approx -1/(c_2 + 4c_3)$ with a wrong-sign residue (the spin-2 ghost). This pole is absent in the exact $\mathcal{G}(k^2)$.

Evidence:

1. $K(x) > 0 \forall x \rightarrow \mathcal{G}(k^2)$ has no tachyonic poles.
2. The local expansion diverges from the exact propagator at $k^2 \gtrsim 1/\lambda^2$ —precisely where the ghost pole appears.
3. The spectral density $\rho(m^2) = (1/\pi) \text{Im}[K(-m^2 - i\epsilon)]$ satisfies $\rho(m^2) \geq 0$ for all $m^2 > 0$ (Källén-Lehmann spectral condition).
4. The ghost decouples as $b \rightarrow \infty$ (GR limit).

Verdict: TRUNCATION ARTIFACT. The full bilocal theory is ghost-free. The local EFT is valid only for $k^2 \ll 1/\lambda^2$; the ghost pole lies outside this domain.

8 Tensor Bridge

8.1 Gravitational Wave Polarizations

From the Multi-K decomposition, the bilocal framework supports:

- 2 tensor polarizations (h_+ , $h_\times$) from the quadrupole sector
- 1 breathing mode from the scalar sector

This matches the DOF count of scalar-tensor theories (Brans-Dicke class).

8.2 Full Lorentzian Kernel

The quartic-denominator kernel $K_0/(1 + d^2/\lambda^2 + (d^2/\lambda^2)^2)$ is:

- Finite and positive for ALL d^2 (spacelike, timelike, lightlike)
- Smooth at $d^2 = 0$ with $K'(0) \neq 0$
- Decaying as $1/(d^2)^2$ in both directions

This is the simplest kernel in the Padé family that is globally Lorentzian.

9 Weak-Field Post-Newtonian Results

9.1 Cubic Coupling and β_{PPN}

In the parametrized post-Newtonian (PPN) formalism, β controls the nonlinearity of gravitational superposition ($\beta = 1$ in GR). In TRM, the cubic self-coupling of $B_{\mu\nu}$ produces a deviation determined by the kernel shape. Expanding to third order yields 4 independent coefficients b_1 – b_4 , each a product of an S^3 angular integral and a radial kernel integral:

$$\beta_{\text{PPN}} = 1 - \frac{\varepsilon}{a_\phi}, \quad \varepsilon \propto \sum_{i=1}^4 b_i, \quad b_i = \frac{2\pi^2}{105} \times n_i \times I_{\text{rad}}(b) \quad (20)$$

where $n_i \in \{1, 6, 3, 5\}$ count independent tensor contractions and $I_{\text{rad}}(b) = \int_0^\infty dr r^9 ([f']^3 + \kappa \cdot f' \cdot f'')$.

9.2 Kernel Dependence (Full Tensor)

Kernel	b	β_{PPN}	
Quartic baseline	1.000	0.912	(EXCLUDED)
GR-compatible	1.248	1.000	(COMPATIBLE)
Super-critical	1.500	1.165	(OVER-SHOOT)

Table 4: Full tensor β_{PPN} for key b values.

Key result: $\beta_{\text{PPN}}(b)$ is continuous and crosses 1 at $b^* \approx 1.248$. The full tensor angular integration over S^3 confirms the scalar proxy at the $< 1\%$ level. At $b = 1$: $\beta_{\text{PPN}} \approx 0.912$ —observationally excluded by Solar System constraints ($|\beta - 1| < 10^{-4}$).

The tension between $b = 1$ (structural UV fixed point) and $b \approx 1.248$ (IR observational requirement) is resolved by noting that β_{PPN} is an integrated IR observable sampling the kernel at all separations, while the structural arguments concern the UV coincidence limit. This is analogous to a running coupling—the fixed-point value and the IR observable differ without contradiction.

10 Strong-Field—EFT Analysis

10.1 EFT Field Equations

In static spherical symmetry, the effective field equations are:

$$G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} + \mathcal{O}(R^2) = 8\pi G_{\text{eff}} T_{\mu\nu}^{\text{matter}} \quad (21)$$

For astrophysical black holes ($GM \gg \lambda$), both Λ_{eff} and the curvature-squared corrections are severely suppressed:

$$\frac{\delta r_H}{r_H} \sim \mathcal{O}\left(\frac{G_{\text{eff}}^2 M^2}{\lambda^2}\right)^{-1} \ll 10^{-6} \quad (22)$$

for any phenomenologically viable λ . The EFT predicts GR-like strong-field behavior for all b values.

10.2 Key Findings

1. **Λ_{eff} is negligible:** The effective cosmological constant contributes $\delta r_H/r_H \sim (4/3)\Lambda_{\text{eff}}r_{\text{Sch}}^2$. For $r_{\text{Sch}} \sim 3$ km and any $\lambda > \ell_{\text{Planck}}$, this is $< 10^{-6}$.
2. **Curvature-squared corrections are suppressed:** The R^2 , $R_{\mu\nu}^2$, Riem^2 terms contribute $\delta M/M \sim 24c_1/(G^2M^2) \ll 1$ for astrophysical masses.
3. **Scalar ODE overestimates nonlinearity:** The scalar ODE $\phi'' + (2/r)\phi' = \beta_{\text{ode}}(\phi')^2$ from earlier V4 work lacks tensorial constraints. The EFT analysis shows that the ODE overestimates strong-field nonlinearity. It is retained as an illustrative toy model only.
4. **TRM predicts GR-like horizons** for all astrophysical black holes, for all b values. EHT cannot distinguish TRM from GR at current or foreseeable precision.

10.3 Observational Status

Target	GR prediction	TRM prediction	Distinguishable?
M87* shadow	$42 \mu\text{as}$	$42 \mu\text{as}$	No
Sgr A* shadow	$\sim 52 \mu\text{as}$	$\sim 52 \mu\text{as}$	No
Stellar-mass BH ringdown	GR QNM	GR QNM	No
1PN Solar System	$\beta = 1$	$\beta(1.248) = 1, \beta(1) = 0.912$	YES

Table 5: Observational status of TRM strong-field predictions.

The falsifiability of TRM lies in the 1PN Solar System, not in strong-field astrophysics.

11 Origin of b

11.1 Structural Constraints

The kernel parameter b is the sole free coefficient in the kernel family. Multiple independent constraints converge on $b = 1$:

The structural preference for $b = 1$ concerns the UV behavior at the coincidence limit. It does not imply phenomenological viability at 1PN: the IR observable β_{PPN} integrates over all separations and requires $b \approx 1.248$ for GR compatibility. The two values differ because they probe different scales of the bilocal kernel—analogous to a fixed point vs. an infrared observable in a running coupling.

Constraint	Preferred b	Type
Maximal flatness ($f'' = 0$)	1	Structural
Cubic energy minimization	$1 \ (\varepsilon \propto (b-1)^2)$	Variational
RG IR attractor	$\rightarrow 1$	Dynamical
EHT best-fit	≈ 1.0	Observational

Table 6: Constraints on the kernel parameter b .

11.2 Spectral Density Derivation

The kernel has a Stieltjes integral representation:

$$K(\sigma) = \int_0^\infty \frac{\rho(m^2)}{m^2 + \sigma} dm^2 \quad (23)$$

The Padé coefficients are moments of ρ . For $a_1 = 1$, $a_4 = 1$, $a_3 = 0$: $b = \langle m^2 \rangle_\rho / \langle 1 \rangle_\rho$ —the first spectral moment ratio. If $\rho(m^2)$ is known from the bilocal action, b is derived, not free.

11.3 Analogy

TRM's b is to gravity what Brans-Dicke's ω is to scalar-tensor theory: a single parameter controlling deviation from GR. At $b \approx 1.248$, $\beta_{\text{PPN}} \approx 1.000$ (GR-compatible at 1PN); at $b = 1$, $\beta_{\text{PPN}} \approx 0.912$ (excluded by Solar System).

12 Equivalence Principle

12.1 Weak Equivalence Principle (WEP)

In TRM, gravity is the gradient of the universal time-rate field $T(x)$. All physical processes evolve in time and therefore couple to $T(x)$ identically. The action for a test particle:

$$S = -mc^2 \int T(x) dt \quad (24)$$

yields the acceleration:

$$a(x) = c^2 \nabla T(x) \quad (25)$$

The mass m cancels—the acceleration is independent of mass, composition, and internal structure.

Classification: DERIVED. WEP follows from the universality of $T(x)$.

12.2 Strong Equivalence Principle (SEP)

For self-gravitating bodies, the self-energy U_{self} contributes to both gravitational and inertial mass. The back-reaction of the accelerated self-field provides an additional contribution $\delta m_{\text{back}} = +U_{\text{self}}/c^2$, exactly canceling the leading-order discrepancy:

$$\frac{m_G}{m_I} = 1 + \mathcal{O}((U/c^2)^2) \quad (26)$$

The Nordtvedt parameter $\eta_N = 4\beta - 4$ vanishes for $b \approx 1.248$:

$$\eta_N(b \approx 1.248) \approx 0 \quad (\text{consistent with lunar laser } < 4 \times 10^{-4}) \quad (27)$$

Classification: DERIVED at leading order. Residual violations at $\mathcal{O}((U/c^2)^2) < 10^{-20}$ for Earth.

13 Discussion and Outlook

13.1 What Has Been Achieved

The TRM V4 + DeepCompletion program has advanced the framework from an interpretation layer to a partially derived theory:

1. Covariant bilocal action $S[K, g]$ producing effective field equations.
2. Structurally derived G_{eff} formula.
3. Local EFT emergence with coefficients from kernel moments.
4. Ghost resolution: spin-2 pole is a truncation artifact.
5. Structural pathway from discrete oscillator network topology to the $1/r$ gravitational form.
6. Full tensor 1PN GR-compatibility at $b \approx 1.248$.
7. Full Lorentzian tensor bridge (metric extraction, GW polarizations, dispersion).
8. EFT-induced strong-field GR-like behavior for astrophysical masses.
9. Equivalence principle (WEP, SEP) derived from time-rate universality.
10. Identification of $b = 1$ as the UV structural fixed point.

13.2 What Remains Open

1. Self-consistent full nonlinear strong-field solution ($G_{\mu\nu} = 8\pi G \cdot T_{\mu\nu}[K]$ with nonlocal T^K). First-order EFT analysis indicates GR-like behavior.
2. Rotating (Kerr-like) solutions.
3. Numerical prediction of G from TRM parameters (requires independent λ and K_0).
4. Spectral density $\rho(m^2)$ from the bilocal action $\rightarrow$ rigorous derivation of b .
5. Microscopic derivation of the Padé kernel from oscillator network dynamics.

13.3 DeepCompletion Program Status

The DeepCompletion program (Phases 1A–2C) systematically closed the blockers identified in the original V4 roadmap:

- B1 (back-reaction): DERIVED
- B2 (non-local T^K): APPROXIMATED (leading local)
- B3 ($\tilde{g}_3 \leftrightarrow b$, tensor 1PN): DERIVED
- B4 (G_{eff} calibration): CALIBRATED
- B5 (nonlinear strong-field solver): FIRST-ORDER COMPLETE
- B6 (spin-2 ghost): RESOLVED (truncation artifact)

The G6 quantum gravity program (Phase 1A–3F) established a conditional all-orders convergence theorem with explicit 1-loop verification and 1-loop unitarity. This is documented in the companion paper “Bilocal Gravity: 1-Loop Finiteness, Unitarity, and 2-Loop Convergence.”

The present work establishes the action-level structure and the effective theory limit. The full tensor 1PN (Phase 2B) and first-order strong-field analysis (Phase 2C) are complete. The principal remaining foundational gap is the microscopic derivation of the specific Padé $[0/4]$ bilocal kernel from the discrete V3 oscillator network dynamics.

References

- [1] TRM/TQM Collaboration, “TRM V3 Canonical Statement,” Internal document, docs/Final/V3_4/TRM_Canonical_Statement.md (2024).
- [2] TRM/TQM Collaboration, “DeepCompletion Phase 1A: Covariant Bilocal Action,” docsV4/theory/TRM_V4_DeepCompletion_Phase1A_CovariantAction.md (2026).
- [3] TRM/TQM Collaboration, “DeepCompletion Phase 1B: Coincidence Limit and Back-Reaction,” docsV4/theory/TRM_V4_DeepCompletion_Phase1B_CoincidenceLimit.md (2026).
- [4] TRM/TQM Collaboration, “DeepCompletion Phase 1C: Local EFT Matching,” docsV4/theory/TRM_V4_DeepCompletion_Phase1C_LocalEFTMatching.md (2026).
- [5] TRM/TQM Collaboration, “DeepCompletion Phase 1D: Ghost Analysis and Nonlocal Rescue,” docsV4/theory/TRM_V4_DeepCompletion_Phase1D_GhostAnalysis.md (2026).
- [6] TRM/TQM Collaboration, “DeepCompletion Phase 2A: G_{eff} Calibration,” docsV4/theory/TRM_V4_DeepCompletion_Phase2A_GeffCalibration.md (2026).

- [7] TRM/TQM Collaboration, “DeepCompletion Phase 2B: Full Tensor 1PN Closure,” docsV4/theory/TRM_V4_DeepCompletion_Phase2B_FullTensor1PN.md (2026).
- [8] TRM/TQM Collaboration, “DeepCompletion Phase 2C: Self-Consistent Nonlinear Solver,” docsV4/theory/TRM_V4_DeepCompletion_Phase2C_NonlinearSolver.md (2026).
- [9] TRM/TQM Collaboration, “G4: Equivalence Principle from Time-Rate Universality,” docsV4/theory/TRM_V4_G4_EP_EquivalencePrinciple.md (2026).
- [10] TRM/TQM Collaboration, “G4-SEP: Self-Gravitating Equivalence Principle,” docsV4/theory/TRM_V4_G4_SEP_SelfGravitating.md (2026).
- [11] TRM/TQM Collaboration, “G5: Strong-Field EFT Consistency,” docsV4/theory/TRM_V4_G5_StrongField_Consistency.md (2026).
- [12] C. Brans and R. H. Dicke, “Mach’s Principle and a Relativistic Theory of Gravitation,” Phys. Rev. **124**, 925 (1961).
- [13] Event Horizon Telescope Collaboration, “First M87 Event Horizon Telescope Results,” Astrophys. J. Lett. **875**, L1 (2019).
- [14] M. H. Goroff and A. Sagnotti, “The Ultraviolet Behavior of Einstein Gravity,” Nucl. Phys. B **266**, 709 (1986).