

# Environmental Toxicology (ET)

Course: ES 535

Dr. Md. Billal Hossain  
Ph.D. (JU)  
Honours and Masters (DU)  
Professional Masters (ITC, UT, Netherlands)

E-mail: [mdbhossain05@gmail.com](mailto:mdbhossain05@gmail.com)  
Cell: 88-01716157306

# Dose Response Relationship

- The dose – response relationship provides an estimation of the relationship between the dose of a chemical agent and incidence of effects in a population.
- The dose is the actual amount of a chemical that enters the body. The dose received may be due to either acute (short) or chronic (long-term) exposure.
- Approaches to characterizing dose – response relationships: include effect levels: such as lethal dose 50 (LD 50 ), lethal concentration 50 (LC 50 ), effective dose 50 (ED 50 ), No observed adverse effect levels (NOAELs), margins of safety.

# Dose Response Relationship


- In toxicology, the lethal dose (LD) is an indication of the lethal toxicity of a given substance or type of radiation
- Because resistance varies from one individual to another, the "lethal dose" represents a dose (usually recorded as dose per kilogram of subject body weight) at which a given percentage of subjects will die
- LD50 (abbreviation for "lethal dose, 50%"), LC50 (lethal concentration, 50%) of a toxin, radiation, or pathogen is the dose required to kill half the members of a tested population after a specified test duration.
- A lower LD50 is indicative of increased toxicity.

# Toxicity

## Lethal Dose-50% ( $LD_{50}$ )



- Lethal dose 50% ( $LD_{50}$ )-dose that is lethal to 50% of a population of test animals
  - Reported by mg of toxicant/kg of body weight
  - Smaller the  $LD_{50}$ =more toxicity of chemical
  - Greater the  $LD_{50}$ =less toxicity of chemical



# Limitations

- As a measure of toxicity, lethal dose is somewhat unreliable and results may vary greatly between testing facilities due to factors such as the
  - ✓ genetic characteristics of the sample population,
  - ✓ animal species tested,
  - ✓ environmental factors and
  - ✓ mode of administration
- There can be wide variability between species as well; what is relatively safe for rats may very well be extremely toxic for humans (cf. paracetamol toxicity), and vice versa.
- For example, chocolate, comparatively harmless to humans, is known to be toxic to many animals.
- When used to test venom from venomous creatures, such as snakes, LD50 results may be misleading due to the physiological differences between mice, rats, and humans.
- Many venomous snakes are specialized predators of mice, and their venom may be adapted specifically to incapacitate mice; and mongooses may be exceptionally resistant. While most mammals have a very similar physiology, LD50 results may or may not have equal bearing upon every mammal species, including humans.

# Toxic Action of Pollutants

- A pollutant can elicit **adverse effects** on the living processes of an organism when present at a **sufficiently high concentration or concentration high enough** to elicit physiological responses in organisms, potentially causing toxicity.
- To **exert damage** to an exposed organism, a pollutant must first enter the **host and reach its target site**.
- A complex pathway exists between the time of initial toxicant **exposure** and the manifestation of **damage** by the organism.
- **Focus on:** general ways in which environmental pollutants exert their actions on **plants, animals, and humans**.

# Plants: Sources, Composition, Uptake

## Sources of pollution:

- **Anthropogenic**: atmospheric pollution: motor vehicles, industrial activities, power generation, brick kilns (mainly in dry season in BD), construction activities, dust, and etc.
- About 58% of the fine particle concentration in Dhaka to the brick kiln industry, 10.4% to vehicles and 15.3% to dusts (CASE project, 2014)

## Composition:

- CO (vehicles), SO<sub>x</sub> (industries, power generation), PM, SO<sub>x</sub>, & VOC's (brick kilns), peroxyacetyl nitrate (PAN), and other photochemical oxidants

## Pollutant uptake:


- Terrestrial plants: exposed: Two main ways: One is exposure of **forage (vegetation)** to air pollutants; another is uptake of toxicants by **roots** growing in **contaminated soils**



## LIVING IN DHAKA

Water supplied to 57 areas was polluted because of faulty pipes


Wasa report in 2019



**AVERAGE TRAFFIC SPEED DECLINED FROM 21KMPH TO 7KMPH IN 10 YEARS**  
World Bank report 2017



**SECOND WORST CITY**  
World Air Quality Report 2021



**DHAKA RANKED 137 AMONG 140 CITIES**  
EIU's Global Liveability Index 2021



**DHAKA NOISIEST CITY IN THE WORLD**  
United Nations Environment Programme Report



## FALLING GROUNDWATER LEVEL

1980	9 metres
1990	22.5 metres
2000	41.5 metres
2020	74 metres

SOURCE: BWDB



# Dhaka Air Quality



## Daily Air Quality Index Report (AQI): Date\_ 08/02/2023 (DoE)



LOCATION	AQI	CATEGORY	RANGE
Dhaka <sup>b</sup>	170	UNHEALTHY	166-174
Chittagong <sup>c</sup>	181	UNHEALTHY	179-183
Gazipur <sup>c</sup>	212	VERY UNHEALTHY	<b>Approved AQI for BD</b> 0-50      Good 51-100    Moderate 101-150   Caution 151-200   Unhealthy 201-300   Very Unhealthy 301-500   Extremely Unhealthy
Narayanganj <sup>c</sup>	170	UNHEALTHY	
Sylhet <sup>c</sup>	182	UNHEALTHY	
Khulna <sup>c</sup>	180	UNHEALTHY	
Rajshahi <sup>c</sup>	286	VERY UNHEALTHY	
Barisal <sup>c</sup>	106	CAUTION	
SavarC	DNA	DNA	
Mymensingh <sup>c</sup>	211	VERY UNHEALTHY	
Rangpur <sup>c</sup>	228	VERY UNHEALTHY	
Cumilla <sup>c</sup>	158	UNHEALTHY	
Narsingdi <sup>c</sup>	132	CAUTION	

NOTE : (a),(b) and (c) Refer to the AQI Average in one city based on three, two and one CAMS respectively. In Bangladesh the AQI is based on 5 criteria pollutants; Particulate Matter (PM10 and PM2.5), NO<sub>2</sub>, CO, SO<sub>2</sub> and Ozone (O<sub>3</sub>)

# Plants: Uptake

- Vegetation growing near **industrial facilities**, such as smelters, aluminum refineries, coal-burning power plants, and brick kilns may absorb airborne pollutants **through the leaves** and become injured
- Important to understand the **uptake of the pollutants** by the plants
- Conductance through the **stomata**, which regulate the passage of ambient air into the cell is important
- The extent of uptake depends on the **chemical and physical properties** of the pollutant along the **gas-to-liquid diffusion pathway**




# Mechanisms of tree damage by toxicants



# Plants: Factors

- The flow of pollutant may be **restricted** by **leaf's physical structure**, or by **scavenging** chemical reactions within the leaf.
- Leaf orientation and morphology, including epidermal characteristics, and air movement across the leaf.
- Stomatal resistance (determined by: stomatal size and number, size of the stomatal **aperture**, and other anatomical characteristics).

# Guard Cells





# Plants: Factors

- **Stomatal opening:** regulated by: light, humidity, temperature, internal carbon dioxide content, water and nutrient status, particularly potassium ( $K^+$ ) ions transported into the guard cells.
- For most plants, the stomata are **open** during the daylight hours, and most damage from sulfur dioxide occurs then.
- Stomatal opening is induced by low  $CO_2$  concentrations, high light intensity and high humidity. By contrast closing is promoted by high  $CO_2$  concentrations, darkness, drought and the plant hormone ABA
- Stomatal opening is extremely important: little or no uptake may occur when the stomata is closed.
- Exposure of roots to toxicants in contaminated soils: waste sites: high levels of heavy metals such as Pb and Cd , often occur
- Metallic ions are more readily released, and thus more readily absorbed, when the soil is acidified by acid deposition.

## Plants: Transport

- Following uptake, a toxicant may undergo mixing with the surrounding medium of the plant, and then be transported to various organs and tissues.
  - Mixing involves microscopic movement of molecules and is accompanied by compensation of concentration differences
- Mode of transport:
- Generally, transport of chemicals in plants occurs passively by diffusion and flux.
  - Diffusion refers to movement across phase boundaries, from a high-concentration compartment to a low-concentration compartment. Flux, on the other hand, is due to the horizontal movement of the medium

# Plant Injury

- To cause **leaf injury**, an **air pollutant** needs to pass through the stomata of the epidermal tissue, as the epidermis is the first target for the pollutant
- In passing into the intercellular spaces, the pollutant may dissolve in the surface water of the leaf cells, affecting cellular pH
- The pollutant may then react with specific cellular constituents, such as, cytoplasmic membrane of the organelles, enzymes, thus affecting cell metabolism, causing plant injury
- Pollutant may then adversely affect cellular metabolism, resulting in plant injury
- Pollution injury is commonly divided into **acute** (destruction of tissues, collapsed leaf margin or other areas changes of leaves colour) and **chronic** (chlorophyll and carotenoids are destroyed) injury.

# Plant Injury: Example


- $\text{SO}_2$  : once absorbed into the leaf,  $\text{SO}_2$  can induce injuries to the ultrastructure of **chloroplasts** and **mitochondria**, which in turn disrupt photosynthesis or cellular energy metabolism
- Alternately,  $\text{SO}_2$  may oxidize and break apart the sulfur bonds in critical enzymes of the membrane, disrupting cellular function
- $\text{O}_3$ : may react with membrane material and induce **peroxidation** of the **lipid components**. This is followed by the formation of various forms of toxic substances, such as aldehydes, ketones, and free radicals
- Cellular enzyme inhibition is often observed when leaves are exposed to atmospheric pollutants

## Contaminated soils: Toxicant uptake

- Exposure of roots to toxicants in contaminated soils is another important process whereby toxicant uptake by plants occurs. Can absorb toxicants through the roots.
- In the contaminated sites, high levels of heavy metals, such as lead (Pb) and cadmium (Cd), chromium (Cr), and aluminum (Al) often occur
- Metallic ions are more readily released, and thus more readily absorbed, when soil is in lower pH condition
- These metal ions may directly damage roots by disrupting water and nutrient uptake, resulting in water deficit or nutrient deficiency
- Following uptake, a toxicant may undergo mixing with the surrounding medium of the plant, and then be transported to various organs and tissues.



# Mammalian Organisms: Exposure

- An environmental pollutant may enter an animal or human through a variety of pathways:
- Inhalation;
- Dermal contact;
- Eye contact; or
- Ingestion





# Process of poisoning in animals and humans



**DANGER**

**CORROSIVE  
CHEMICALS**



**AVOID CONTACT WITH  
EYES & SKIN**

Skin Contact/Absorption





Skin Contact/Absorption

# Mammalian Organisms: Uptake

- The immediate and long-term effects of a pollutant are directly related to its **mode of entry**
- The portals of entry for an atmospheric pollutant are **skin, eyes, lungs, and gastrointestinal tract**
- And the **hair follicles, sweat glands, and open wounds** are the possible entry sites where uptake from the skin may occur
- Uptake of toxicants by gastrointestinal tract may occur when consumed foods or beverages are contaminated by air pollutants, such as Pb, Cd, or sprayed pesticides

## Mammalian organisms: Uptake

- For a pollutant to be taken up into the body and finally carried to a cell, it must pass through several layers of biological membranes.
- These include not only the peripheral tissue membranes, but also the capillary and cell membranes.
- Therefore, the nature of the membranes and the chemical and physical properties of the toxicant in question are important factors affecting uptake.



## Mammalian organisms: Uptake

- The mechanisms by which chemical agents pass through the membrane include:
  - Filtration through spaces or pores in membranes
  - Passive diffusion through the spaces or pores, or by dissolving in the lipid material of the membrane
  - Facilitated transport, where a specialized protein molecule, called a carrier, carries a water-soluble substance across the membrane
  - Active transport, which requires both a carrier and energy



## Mammalian organisms: Transport

- After absorption, a toxicant may be bound to a blood protein, forming a complex, or it may exist in a free form
- Transport of a toxicant may occur through the bloodstream or lymphatic system
- The toxicant may then be distributed to various body tissues, including those of storage depots and sites of metabolism

## **Mammalian organisms: Storage**

- A toxicant may be stored in the liver, lungs, kidneys, bones, or adipose tissue
- These storage depots may or may not be the sites of toxic action.
- A toxicant may be stored in a depot temporarily and then released and translocated again.
- Similarly, a toxicant or its metabolite may be transported to a storage site and remain there for along period of time, even permanently

# Metabolism

- The set of chemical reactions that occur in living organisms in order to maintain life. These processes allow organisms to grow and reproduce, maintain their structures, and respond to their environments.
- Metabolism is usually divided into two categories: **Catabolism** breaks down organic matter, for example to harvest energy in cellular respiration, **Anabolism**, on the other hand, uses energy to construct components of cells such as proteins and nucleic acids.


# Metabolism of Toxicants

- Metabolism of toxicants may occur at the portals of entry, or in such organs as the skin, lungs, liver, kidney, and gastrointestinal tract
- **Liver** plays a central role in metabolism of environmental toxicants (xenobiotics)
- Liver contains a rich supply of nonspecific enzymes, enabling it to metabolize a broad spectrum of organic compounds

## Mammalian organisms: Excretion

- The final step in the pathway of a toxicant is its excretion from the body.
- Excretion may occur through the **lungs, kidneys, or gastrointestinal (GI)** tract.
- A toxicant may be excreted in its original form or as its metabolites, depending on its chemical property.
- Excretion is the most permanent means whereby toxic substances are removed from the body.

# Processes of poisoning in animals and humans





# Mechanism of Action


- Toxic action of pollutants involves: either compounds with **intrinsic toxicity** or **activated metabolites**
- The duration of toxic action depends on the characteristics of the toxicant and the physiological or biochemical functioning of the host organism
- The ways xenobiotics can induce adverse effects in living organisms include:
  - Disruption or destruction of cellular structure
  - Direct chemical combination with a cell constituent
  - Inhibition of enzymes
  - Initiation of a secondary action
  - Disruption of reproductive function

## Disruption or destruction of cellular structure

- A toxicant may induce an injurious effect on plant or animal tissues by disrupting or destroying the cellular structure
- Atmospheric pollutants, such as  $\text{SO}_2$ ,  $\text{NO}_2$ , and  $\text{O}_3$ , are **phytotoxic** – they can cause plant injuries.
- Studies show that low concentrations of  $\text{SO}_2$  can injure epidermal and guard cells, leading to enhanced stomatal conductance and greater entry of the pollutant into leaves
- In animals and humans Inhalation of sufficient quantities of  $\text{NO}_2$  and sulfuric acid mists can damage surface layers of the respiratory system
- High levels of  $\text{O}_3$  can cause problems with the lipid portion of the membrane, resulting in disruption of membrane structure.

# Chemical combination with a cell constituent


- A pollutant may combine with a cell constituent, forming a complex and disrupting cellular metabolism
- CO has the ability to bind to hemoglobin (Hb); after inhalation and diffusion into the blood, CO readily reacts with Hb to form carboxyhemoglobin (COHb).



- Presence of a large amount of COHb in the blood **disrupts the vital system** for exchange of  $\text{CO}_2$  and  $\text{O}_2$  between the blood and lungs and other body tissues. Interference with the functioning of hemoglobin by COHb accumulation is detrimental to health and can lead to death.

# Effect of Enzymes

- Reactions in living cell: Participation of enzymes as biological catalysts
- Almost all enzymes are proteins with a globular structure; few requires non protein components (**cofactors**) e.g., metal ions ( $K^+$ ,  $Na^+$ ,  $Cu^{2+}$ ,  $Fe^{2+}$ ,  $Fe^{3+}$ ,  $Mg^{2+}$ , organic molecules (coenzymes)
- However, the optimum activity of many enzymes depend on the presence of non protein substances called **cofactors**
- The molecular partnership of protein-cofactor is **holoenzyme** and exhibits maximum catalytic activity
- The protein component without its cofactor is termed an **apoenzyme** and exhibits **very low activity** or none at all



# Effect of Enzymes

- Ways in which environmental **pollutants may inactivate an enzyme** systems
- A pollutant may bind to the active site of an enzyme thus inactivating it
  - For example, a heavy metal such as Pb, Cd or Hg, after absorption into the body may attach itself to the thiol or sulfhydryl (-SH) group on a protein enzyme, forming a covalent bond with the sulfur atom. This will lead to the inactivation of the enzyme, with active site being blocked, the activity of the enzyme will be depressed or lost.

