Mahati Akella 22070521027

```
#PRINCIPAL COMPONENT ANALYSIS AND DATA Visualisation

#TO PERFORM DIMENSIONALITY REDUCTION USING PCA AND CREATE APPROPRIATE PLOTS TO ANALYZE COMPONETS  ON IRIS DATASET

import pandas as pd
import numpy as np
import matplotlib.pyplot as plt
import seaborn as sns
from sklearn.preprocessing import StandardScaler
from sklearn.decomposition import PCA


# Load dataset
df = pd.read_csv('Iris.csv')

# Display first few rows
df.head()
```

	Id	SepalLengthCm	SepalWidthCm	PetalLengthCm	PetalWidthCm	Species
0	1	5.1	3.5	1.4	0.2	Iris-setosa
1	2	4.9	3.0	1.4	0.2	Iris-setosa
2	3	4.7	3.2	1.3	0.2	Iris-setosa
3	4	4.6	3.1	1.5	0.2	Iris-setosa
4	5	5.0	3.6	1.4	0.2	Iris-setosa

Next steps: Generate code with df | View recommended plots | New interactive sheet

```
# Drop unnecessary columns like Id (if present)
if 'Id' in df.columns:
    df = df.drop(['Id'], axis=1)

# Separate features and target
features = df.columns[:-1]  # All columns except 'Species'
X = df[features]
y = df['Species']

# Standardize the features
scaler = StandardScaler()
X_scaled = scaler.fit_transform(X)


# Initialize PCA
pca = PCA(n_components=2)  # Reduce to 2 components
X_pca = pca.fit_transform(X_scaled)

# Create DataFrame for PCA result
pca_df = pd.DataFrame(data=X_pca, columns=['PC1', 'PC2'])
pca_df['Species'] = y


plt.figure(figsize=(8,6))
sns.scatterplot(data=pca_df, x='PC1', y='PC2', hue='Species', palette='Set2')
plt.title('PCA of Iris Dataset')
plt.xlabel('Principal Component 1')
plt.ylabel('Principal Component 2')
plt.grid(True)
plt.show()
```




```
print("Explained variance by each component:")
print(pca.explained_variance_ratio_)

# Bar plot of explained variance
plt.figure(figsize=(6,4))
plt.bar(['PC1', 'PC2'], pca.explained_variance_ratio_, color='skyblue')
plt.title('Explained Variance by Principal Components')
plt.ylabel('Variance Ratio')
plt.show()
```

```
Explained variance by each component:
[0.72770452 0.23030523]
```




```
# Try PCA with all components to see cumulative explained variance
pca_full = PCA()
pca_full.fit(X_scaled)

# Plot cumulative explained variance
plt.figure(figsize=(6,4))
plt.plot(np.cumsum(pca_full.explained_variance_ratio_), marker='o')
plt.title('Cumulative Explained Variance')
plt.xlabel('Number of Components')
plt.ylabel('Cumulative Variance Explained')
plt.grid(True)
plt.show()
```


Cumulative Explained Variance
Cumulative Variance Explained
1.00
0.95
0.90
0.85
0.80
0.75
0.0
0.5
1.0
1.5
2.0
2.5
3.0
Number of Components