

# Today: Neural Networks

## Problem: How to compute gradients?

$$s = f(x; W_1, W_2) = W_2 \max(0, W_1 x) \quad \text{Nonlinear score function}$$

$$L_i = \sum_{j \neq y_i} \max(0, s_j - s_{y_i} + 1) \quad \text{SVM Loss on predictions}$$

$$R(W) = \sum_k W_k^2 \quad \text{Regularization}$$

$$L = \frac{1}{N} \sum_{i=1}^N L_i + \lambda R(W_1) + \lambda R(W_2) \quad \text{Total loss: data loss + regularization}$$

If we can compute  $\frac{\partial L}{\partial W_1}, \frac{\partial L}{\partial W_2}$  then we can learn  $W_1$  and  $W_2$

## (Bad) Idea: Derive $\nabla_W L$ on paper

$$s = f(x; W) = Wx$$

$$L_i = \sum_{j \neq y_i} \max(0, s_j - s_{y_i} + 1)$$

$$= \sum_{j \neq y_i} \max(0, W_{j,:} \cdot x + W_{y_i,:} \cdot x + 1)$$

$$L = \frac{1}{N} \sum_{i=1}^N L_i + \lambda \sum_k W_k^2$$

$$= \frac{1}{N} \sum_{i=1}^N \sum_{j \neq y_i} \max(0, W_{j,:} \cdot x + W_{y_i,:} \cdot x + 1) + \lambda \sum_k W_k^2$$


$$\nabla_W L = \nabla_W \left( \frac{1}{N} \sum_{i=1}^N \sum_{j \neq y_i} \max(0, W_{j,:} \cdot x + W_{y_i,:} \cdot x + 1) + \lambda \sum_k W_k^2 \right)$$

**Problem:** Very tedious: Lots of matrix calculus, need lots of paper

**Problem:** What if we want to change loss? E.g. use softmax instead of SVM? Need to re-derive from scratch =(

**Problem:** Not feasible for very complex models!

## Better Idea: Computational graphs + Backpropagation



# Convolutional network (AlexNet)

input image

weights

loss




Figure copyright Alex Krizhevsky, Ilya Sutskever, and Geoffrey Hinton, 2012. Reproduced with permission.

# Neural Turing Machine

input image

loss




Figure reproduced with permission from a [Twitter post](#) by Andrej Karpathy.

# Neural Turing Machine




Figure reproduced with permission from a [Twitter post](#) by Andrej Karpathy.


Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$



Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$





Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream  
gradient

Local  
gradient

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream  
gradient

Local  
gradient

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial x}$$

Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

Upstream  
gradient

Local  
gradient

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$




$$\frac{\partial f}{\partial x}$$


Chain rule:


$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

Upstream  
gradient


Local  
gradient




















Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$



Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$

$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$




Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$




$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$

$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$

$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$

$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$


Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$




Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2 x_2)}}$$




$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$


Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!



Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!




Sigmoid local gradient:

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left( \frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left( \frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!



Sigmoid function

$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

Sigmoid


[upstream gradient] x [local gradient]  
 $[1.00] \times [(1 - 0.73) (0.73)] = 0.2$

Sigmoid local gradient:

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left( \frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left( \frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$


# Patterns in gradient flow

**add** gate: gradient distributor




# Patterns in gradient flow

**add** gate: gradient distributor




**mul** gate: “swap multiplier”




# Patterns in gradient flow


**add** gate: gradient distributor



**mul** gate: “swap multiplier”




**copy** gate: gradient adder






# Patterns in gradient flow


**add** gate: gradient distributor




**mul** gate: “swap multiplier”




**copy** gate: gradient adder



**max** gate: gradient router



# Backprop Implementation: “Flat” code



Forward pass:  
Compute output


```
def f(w0, x0, w1, x1, w2):
```

```
    s0 = w0 * x0
    s1 = w1 * x1
    s2 = s0 + s1
    s3 = s2 + w2
    L = sigmoid(s3)
```

Backward pass:  
Compute grads

```
    grad_L = 1.0
    grad_s3 = grad_L * (1 - L) * L
    grad_w2 = grad_s3
    grad_s2 = grad_s3
    grad_s0 = grad_s2
    grad_s1 = grad_s2
    grad_w1 = grad_s1 * x1
    grad_x1 = grad_s1 * w1
    grad_w0 = grad_s0 * x0
    grad_x0 = grad_s0 * w0
```

# Backprop Implementation: “Flat” code



Forward pass:  
Compute output

```
def f(w0, x0, w1, x1, w2):
```


```
    s0 = w0 * x0
    s1 = w1 * x1
    s2 = s0 + s1
    s3 = s2 + w2
    L = sigmoid(s3)
```

Base case

```
    grad_L = 1.0
```

```
    grad_s3 = grad_L * (1 - L) * L
    grad_w2 = grad_s3
    grad_s2 = grad_s3
    grad_s0 = grad_s2
    grad_s1 = grad_s2
    grad_w1 = grad_s1 * x1
    grad_x1 = grad_s1 * w1
    grad_w0 = grad_s0 * x0
    grad_x0 = grad_s0 * w0
```

# Backprop Implementation: “Flat” code



Forward pass:  
Compute output

Sigmoid

```
def f(w0, x0, w1, x1, w2):
```

```
    s0 = w0 * x0
```

```
    s1 = w1 * x1
```

```
    s2 = s0 + s1
```

```
    s3 = s2 + w2
```

```
    L = sigmoid(s3)
```

```
    grad_L = 1.0
```

```
    grad_s3 = grad_L * (1 - L) * L
```

```
    grad_w2 = grad_s3
```

```
    grad_s2 = grad_s3
```

```
    grad_s0 = grad_s2
```

```
    grad_s1 = grad_s2
```


```
    grad_w1 = grad_s1 * x1
```

```
    grad_x1 = grad_s1 * w1
```

```
    grad_w0 = grad_s0 * x0
```

```
    grad_x0 = grad_s0 * w0
```

# Backprop Implementation: “Flat” code




Forward pass:  
Compute output

```
def f(w0, x0, w1, x1, w2):
    s0 = w0 * x0
    s1 = w1 * x1
    s2 = s0 + s1
    s3 = s2 + w2
    L = sigmoid(s3)
```

Add gate

```
grad_L = 1.0
grad_s3 = grad_L * (1 - L) * L
grad_w2 = grad_s3
grad_s2 = grad_s3
grad_s0 = grad_s2
grad_s1 = grad_s2
grad_w1 = grad_s1 * x1
grad_x1 = grad_s1 * w1
grad_w0 = grad_s0 * x0
grad_x0 = grad_s0 * w0
```

# Backprop Implementation: “Flat” code




Forward pass:  
Compute output

```
def f(w0, x0, w1, x1, w2):
    s0 = w0 * x0
    s1 = w1 * x1
    s2 = s0 + s1
    s3 = s2 + w2
    L = sigmoid(s3)
```

Add gate

```
grad_L = 1.0
grad_s3 = grad_L * (1 - L) * L
grad_w2 = grad_s3
grad_s2 = grad_s3
grad_s0 = grad_s2
grad_s1 = grad_s2
grad_w1 = grad_s1 * x1
grad_x1 = grad_s1 * w1
grad_w0 = grad_s0 * x0
grad_x0 = grad_s0 * w0
```

# Backprop Implementation: “Flat” code



Forward pass:  
Compute output

```
def f(w0, x0, w1, x1, w2):
```

```
    s0 = w0 * x0
    s1 = w1 * x1
    s2 = s0 + s1
    s3 = s2 + w2
    L = sigmoid(s3)
```

```
    grad_L = 1.0
```

```
    grad_s3 = grad_L * (1 - L) * L
```

```
    grad_w2 = grad_s3
```

```
    grad_s2 = grad_s3
```

```
    grad_s0 = grad_s2
```

```
    grad_s1 = grad_s2
```

```
    grad_w1 = grad_s1 * x1
```


```
    grad_x1 = grad_s1 * w1
```

```
    grad_w0 = grad_s0 * x0
```

```
    grad_x0 = grad_s0 * w0
```

Multiply gate

# Backprop Implementation: “Flat” code



Forward pass:  
Compute output

```
def f(w0, x0, w1, x1, w2):
```

```
    s0 = w0 * x0
```

```
    s1 = w1 * x1
```

```
    s2 = s0 + s1
```

```
    s3 = s2 + w2
```

```
    L = sigmoid(s3)
```

```
grad_L = 1.0
```

```
grad_s3 = grad_L * (1 - L) * L
```

```
grad_w2 = grad_s3
```

```
grad_s2 = grad_s3
```

```
grad_s0 = grad_s2
```

```
grad_s1 = grad_s2
```

```
grad_w1 = grad_s1 * x1
```

```
grad_x1 = grad_s1 * w1
```

```
grad_w0 = grad_s0 * x0
```

```
grad_x0 = grad_s0 * w0
```

Multiply gate




# “Flat” Backprop: Do this for assignment 1!

Stage your forward/backward computation!

E.g. for the SVM:

```
# receive W (weights), X (data)
# forward pass (we have 8 lines)
scores = #...
margins = #...
data_loss = #...
reg_loss = #...
loss = data_loss + reg_loss
# backward pass (we have 5 lines)
dmargins = # ... (optionally, we go direct to dscores)
dscores = #...
dW = #...
```



# “Flat” Backprop: Do this for assignment 1!

E.g. for two-layer neural net:

```
# receive W1,W2,b1,b2 (weights/biases), X (data)
# forward pass:
h1 = #... function of X,W1,b1
scores = #... function of h1,W2,b2
loss = #... (several lines of code to evaluate Softmax loss)
# backward pass:
dscores = #...
dh1,dW2,db2 = #...
dW1,db1 = #...
```

# Recap: Vector derivatives

Scalar to Scalar

$$x \in \mathbb{R}, y \in \mathbb{R}$$

Regular derivative:

$$\frac{\partial y}{\partial x} \in \mathbb{R}$$

If  $x$  changes by a small amount, how much will  $y$  change?

Vector to Scalar

$$x \in \mathbb{R}^N, y \in \mathbb{R}$$

Derivative is **Gradient**:

$$\frac{\partial y}{\partial x} \in \mathbb{R}^N \quad \left( \frac{\partial y}{\partial x} \right)_n = \frac{\partial y}{\partial x_n}$$

For each element of  $x$ , if it changes by a small amount then how much will  $y$  change?

# Recap: Vector derivatives

## Scalar to Scalar

$$x \in \mathbb{R}, y \in \mathbb{R}$$

Regular derivative:

$$\frac{\partial y}{\partial x} \in \mathbb{R}$$

If  $x$  changes by a small amount, how much will  $y$  change?

## Vector to Scalar

$$x \in \mathbb{R}^N, y \in \mathbb{R}$$

Derivative is **Gradient**:

$$\frac{\partial y}{\partial x} \in \mathbb{R}^N \quad \left( \frac{\partial y}{\partial x} \right)_n = \frac{\partial y}{\partial x_n}$$

For each element of  $x$ , if it changes by a small amount then how much will  $y$  change?

## Vector to Vector


$$x \in \mathbb{R}^N, y \in \mathbb{R}^M$$

Derivative is **Jacobian**:


$$\frac{\partial y}{\partial x} \in \mathbb{R}^{N \times M} \quad \left( \frac{\partial y}{\partial x} \right)_{n,m} = \frac{\partial y_m}{\partial x_n}$$

For each element of  $x$ , if it changes by a small amount then how much will each element of  $y$  change?


## Backprop with Vectors




## Backprop with Vectors



## Backprop with Vectors




## Backprop with Vectors






# Backprop with Vectors



# Backprop with Vectors

4D input x:


$\begin{bmatrix} 1 \\ -2 \\ 3 \\ -1 \end{bmatrix}$



$f(x) = \max(0, x)$   
(*elementwise*)

4D output y:

$\begin{bmatrix} 1 \\ 0 \\ 3 \\ 0 \end{bmatrix}$



4D dL/dy:


$\begin{bmatrix} 4 \\ -1 \\ 5 \\ 9 \end{bmatrix}$

Upstream  
gradient

# Backprop with Vectors

4D input x:


$\begin{bmatrix} 1 \\ -2 \\ 3 \\ -1 \end{bmatrix}$



$f(x) = \max(0, x)$   
(*elementwise*)

4D output y:

$\begin{bmatrix} 1 \\ 0 \\ 3 \\ 0 \end{bmatrix}$




Jacobian  $dy/dx$

$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$


4D  $dL/dy$ :

$\begin{bmatrix} 4 \\ -1 \\ 5 \\ 9 \end{bmatrix}$


Upstream  
gradient




# Backprop with Vectors




# Backprop with Vectors



# Backprop with Vectors




# Backprop with Vectors



## Backprop with Matrices (or Tensors)

Loss L still a scalar!

$dL/dx$  always has the same shape as  $x$ !






# Backprop with Matrices (or Tensors)

Loss L still a scalar!

$dL/dx$  always has the same shape as  $x$ !



## Backprop with Matrices

x: [N×D]

[ 2 1 -3 ]

[ -3 4 2 ]

w: [D×M]

[ 3 2 1 -1 ]

[ 2 1 3 2 ]

[ 3 2 1 -2 ]

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

y: [N×M]

[13 9 -2 -6]

[ 5 2 17 1 ]

dL/dy: [N×M]

[ 2 3 -3 9 ]

[ -8 1 4 6 ]

Also see derivation in the course notes:

<http://cs231n.stanford.edu/handouts/linear-backprop.pdf>

## Backprop with Matrices

x: [N×D]  
[ 2 1 -3 ]  
[ -3 4 2 ]

w: [D×M]  
[ 3 2 1 -1 ]  
[ 2 1 3 2 ]  
[ 3 2 1 -2 ]

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

y: [N×M]  
[13 9 -2 -6]  
[ 5 2 17 1 ]

dL/dy: [N×M]  
[ 2 3 -3 9 ]  
[ -8 1 4 6 ]

**Jacobians:**  
dy/dx: [(N×D)×(N×M)]  
dy/dw: [(D×M)×(N×M)]

For a neural net we may have  
N=64, D=M=4096  
Each Jacobian takes 256 GB of memory!  
Must work with them implicitly!

## Backprop with Matrices

x: [N×D]  
[ 2 1 -3 ]  
[ -3 4 2 ]

w: [D×M]  
[ 3 2 1 -1 ]  
[ 2 1 3 2 ]  
[ 3 2 1 -2 ]

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

**Q:** What parts of y  
are affected by one  
element of x?

y: [N×M]  
[ 13 9 -2 -6 ]  
[ 5 2 17 1 ]

dL/dy: [N×M]  
[ 2 3 -3 9 ]  
[ -8 1 4 6 ]

## Backprop with Matrices

$x: [N \times D]$   
 $\begin{bmatrix} 2 & \boxed{1} & -3 \\ -3 & 4 & 2 \end{bmatrix}$   
 $w: [D \times M]$   
 $\begin{bmatrix} 3 & 2 & 1 & -1 \\ 2 & 1 & 3 & 2 \\ 3 & 2 & 1 & -2 \end{bmatrix}$

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

$y: [N \times M]$   
 $\begin{bmatrix} \boxed{13} & \boxed{9} & \boxed{-2} & \boxed{-6} \\ 5 & 2 & 17 & 1 \end{bmatrix}$

$dL/dy: [N \times M]$   
 $\begin{bmatrix} \boxed{2} & \boxed{3} & \boxed{-3} & \boxed{9} \\ -8 & 1 & 4 & 6 \end{bmatrix}$

**Q:** What parts of  $y$  are affected by one element of  $x$ ?

**A:**  $x_{n,d}$  affects the whole row  $y_{n,\cdot}$ .

$$\frac{\partial L}{\partial x_{n,d}} = \sum_m \frac{\partial L}{\partial y_{n,m}} \frac{\partial y_{n,m}}{\partial x_{n,d}}$$

## Backprop with Matrices

$x: [N \times D]$   
 $\begin{bmatrix} 2 & \boxed{1} & -3 \\ -3 & 4 & 2 \end{bmatrix}$   
 $w: [D \times M]$   
 $\begin{bmatrix} 3 & 2 & 1 & -1 \\ 2 & 1 & 3 & 2 \\ 3 & 2 & 1 & -2 \end{bmatrix}$

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

$y: [N \times M]$   
 $\begin{bmatrix} \boxed{13} & 9 & \boxed{-2} & -6 \\ 5 & 2 & 17 & 1 \end{bmatrix}$

$dL/dy: [N \times M]$   
 $\begin{bmatrix} \boxed{2} & 3 & -3 & 9 \\ -8 & 1 & 4 & 6 \end{bmatrix}$

**Q:** What parts of  $y$  are affected by one element of  $x$ ?

**A:**  $x_{n,d}$  affects the whole row  $y_{n,\cdot}$ .

**Q:** How much does  $x_{n,d}$  affect  $y_{n,m}$ ?

$$\frac{\partial L}{\partial x_{n,d}} = \sum_m \frac{\partial L}{\partial y_{n,m}} \frac{\partial y_{n,m}}{\partial x_{n,d}}$$

# Backprop with Matrices

x: [N×D]  
 $\begin{bmatrix} 2 & \boxed{1} & -3 \\ -3 & 4 & 2 \end{bmatrix}$

w: [D×M]  
 $\begin{bmatrix} 3 & 2 & 1 & -1 \\ 2 & 1 & \boxed{3} & 2 \\ 3 & 2 & 1 & -2 \end{bmatrix}$

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

y: [N×M]  
 $\begin{bmatrix} \boxed{13} & 9 & \boxed{-2} & -6 \\ 5 & 2 & 17 & 1 \end{bmatrix}$

dL/dy: [N×M]  
 $\begin{bmatrix} \boxed{2} & 3 & -3 & 9 \\ -8 & 1 & 4 & 6 \end{bmatrix}$

**Q:** What parts of y are affected by one element of x?

**A:**  $x_{n,d}$  affects the whole row  $y_{n,\cdot}$ .

**Q:** How much does  $x_{n,d}$  affect  $y_{n,m}$ ?

**A:**  $w_{d,m}$

$$\frac{\partial L}{\partial x_{n,d}} = \sum_m \frac{\partial L}{\partial y_{n,m}} \frac{\partial y_{n,m}}{\partial x_{n,d}} = \sum_m \frac{\partial L}{\partial y_{n,m}} w_{d,m}$$

# Backprop with Matrices

$$x: [N \times D]$$

$$\begin{bmatrix} 2 & \boxed{1} & -3 \\ -3 & 4 & 2 \end{bmatrix}$$

$$w: [D \times M]$$

$$\begin{bmatrix} 3 & 2 & 1 & -1 \\ 2 & 1 & \boxed{3} & 2 \\ 3 & 2 & 1 & -2 \end{bmatrix}$$

$[N \times D]$   $[N \times M]$   $[M \times D]$

$$\boxed{\frac{\partial L}{\partial x} = \left( \frac{\partial L}{\partial y} \right) w^T}$$

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

$$y: [N \times M]$$

$$\begin{bmatrix} \boxed{13} & \boxed{9} & \boxed{-2} & \boxed{-6} \\ 5 & 2 & 17 & 1 \end{bmatrix}$$

$$dL/dy: [N \times M]$$

$$\begin{bmatrix} \boxed{2} & \boxed{3} & \boxed{-3} & \boxed{9} \\ -8 & 1 & 4 & 6 \end{bmatrix}$$

**Q:** What parts of  $y$  are affected by one element of  $x$ ?

**A:**  $x_{n,d}$  affects the whole row  $y_{n,\cdot}$ .

**Q:** How much does  $x_{n,d}$  affect  $y_{n,m}$ ?

**A:**  $w_{d,m}$

$$\frac{\partial L}{\partial x_{n,d}} = \sum_m \frac{\partial L}{\partial y_{n,m}} \frac{\partial y_{n,m}}{\partial x_{n,d}} = \sum_m \frac{\partial L}{\partial y_{n,m}} w_{d,m}$$



## Backprop with Matrices

$$\begin{aligned}
 &x: [N \times D] \\
 &\begin{bmatrix} 2 & \boxed{1} & -3 \\ -3 & 4 & 2 \end{bmatrix} \\
 &w: [D \times M] \\
 &\begin{bmatrix} 3 & 2 & 1 & -1 \\ 2 & 1 & \boxed{3} & 2 \\ 3 & 2 & 1 & -2 \end{bmatrix}
 \end{aligned}$$

Matrix Multiply

$$y_{n,m} = \sum_d x_{n,d} w_{d,m}$$

$$\begin{aligned}
 &y: [N \times M] \\
 &\begin{bmatrix} \boxed{13} & \boxed{9} & \boxed{-2} & \boxed{-6} \\ 5 & 2 & 17 & 1 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 &dL/dy: [N \times M] \\
 &\begin{bmatrix} \boxed{2} & \boxed{3} & \boxed{-3} & \boxed{9} \\ -8 & 1 & 4 & 6 \end{bmatrix}
 \end{aligned}$$

By similar logic:

$$[N \times D] \quad [N \times M] \quad [M \times D]$$

$$\frac{\partial L}{\partial x} = \left( \frac{\partial L}{\partial y} \right) w^T$$

$$[D \times M] \quad [D \times N] \quad [N \times M]$$


$$\frac{\partial L}{\partial w} = x^T \left( \frac{\partial L}{\partial y} \right)$$

These formulas are easy to remember: they are the only way to make shapes match up!

## Summary for today:


- **(Fully-connected) Neural Networks** are stacks of linear functions and nonlinear activation functions; they have much more representational power than linear classifiers
- **backpropagation** = recursive application of the chain rule along a computational graph to compute the gradients of all inputs/parameters/intermediates
- implementations maintain a graph structure, where the nodes implement the **forward()** / **backward()** API
- **forward**: compute result of an operation and save any intermediates needed for gradient computation in memory
- **backward**: apply the chain rule to compute the gradient of the loss function with respect to the inputs

# Next Time: Convolutional Networks!




A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$\in \mathbb{R}^n \quad \in \mathbb{R}^{n \times n}$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$




A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial f}{\partial q_i} = 2q_i$$

$$\nabla_q f = 2q$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial f}{\partial q_i} = 2q_i$$

$$\nabla_q f = 2q$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$

$$\begin{bmatrix} 0.1 & 0.5 \\ -0.3 & 0.8 \end{bmatrix} W$$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial q_k}{\partial W_{i,j}} = \mathbf{1}_{k=i}x_j$$

$$\begin{aligned} \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k)(\mathbf{1}_{k=i}x_j) \\ &= 2q_i x_j \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial W_{i,j}} &= \mathbf{1}_{k=i} x_j \\ \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k) (\mathbf{1}_{k=i} x_j) \\ &= 2q_i x_j \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$


$$\begin{aligned} \frac{\partial q_k}{\partial W_{i,j}} &= \mathbf{1}_{k=i} x_j \\ \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k) (\mathbf{1}_{k=i} x_j) \\ &= 2q_i x_j \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$





A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial q_k}{\partial x_i} = W_{k,i}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial x_i} &= W_{k,i} \\ \frac{\partial f}{\partial x_i} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial x_i} \\ &= \sum_k 2q_k W_{k,i} \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



In discussion section: A matrix example...

$$z_1 = XW_1$$

$$h_1 = \text{ReLU}(z_1)$$

$$\hat{y} = h_1 W_2$$

$$L = ||\hat{y}||_2^2$$

$$\frac{\partial L}{\partial W_2} = ?$$

$$\frac{\partial L}{\partial W_1} = ?$$

