



27



M is no of symbol.

n is no bit in symbol.

in PSK $M = 2$, and $n = 1$ in QPSK $M = 4$ and $n = 2$ or
4 PSK.

Bandwidth in M-ary PSK

$$= \frac{2}{\log_2 M} f_b$$



* Phase shift for M-ary PSK

Consecutive Symbol = $\frac{2\pi}{M}$


* and symbol duration $T = nT_b$

January 2017						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

Notes:

28

General signal space diagram



29

Notes:

February 2017						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28				

2017



My Family

Monday

Continuous Phase

Week 6 / Day 30

is memory base Modulation Scheme, where phase to be continuous.

30

Frequency Shift Keying (CPFSK)



My Friends

Discontinuity in phase in QPSK during symbol change, which generate large spectral side lobe outside of main spectral band.

Here we define CPFSK signal

$$s(t) = \sqrt{\frac{2E_b}{T_b}} \cos[2\pi f_1 t + \theta(0)] \rightarrow '1'$$

$$= \sqrt{\frac{2E_b}{T_b}} \cos[2\pi f_2 t + \theta(0)] \rightarrow '0'$$



My Health

$\theta(0)$ is the value of Phase at $t=0$ (based on past history)



My Skills

Another way we may write

$$s(t) = \sqrt{\frac{2E_b}{T_b}} \cos(2\pi f_c t + \theta(t)) \quad \text{--- (3)}$$

$$\text{where } \theta(t) = \theta(0) + \frac{\pi h f t}{T_b} \quad 0 \leq t \leq T_b$$



My Money

As $\theta(t)$ is continuous f'n of t , $s(t)$ is continuous at all time.

+ for '1' sending
0 - for '0' sending

including interbit switching times.

January 2017						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

Notes:



My Family

Tuesday

Week 6 / Day 31

31

$s(t) = \sqrt{\frac{2E_b}{T_b}} \cos\left[2\pi f_c t \pm \frac{\pi h t}{T_b} + \theta(0)\right]$
 Compare with eq (144) and (244), we get (4)



My Friends

$$f_c + \frac{h}{2T_b} = f_1 \quad \text{for symbol '1'}$$

$$f_c - \frac{h}{2T_b} = f_2 \quad \text{for symbol '0'}$$

$$\therefore f_c = \frac{f_1 + f_2}{2}$$



My Health

$$h = \frac{2}{T_b} (f_1 - f_2) = \frac{f_1 - f_2}{\frac{1}{T_b} = f_b}$$

know as
deviation ration

now, $0 \leq t \leq T_b$

(normalised
freq diff.)



My Skills

$$\theta(t) = \theta(0) \pm \frac{\pi h t}{T_b} \quad \text{--- (5)}$$

$$\text{at } t = T_b \quad \theta(T_b) = \theta(0) \pm \frac{\pi h T_b}{T_b}$$

$$\therefore \theta(T_b) - \theta(0) = \pm \pi h \quad \text{--- (6)}$$



My Money

$$\text{for } T_b \leq t \leq 2T_b \quad \theta(t) = \theta(T_b) \pm \frac{\pi h}{T_b} t$$

Notes:

February 2017						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28				

Notes

Now, consider, $\theta(0) = 0$, $h = \frac{1}{2}$

eq (6) \Rightarrow Phase Change $\left[\begin{matrix} \theta(T_b) = \pm \frac{\pi}{2} \\ -\theta(0) = -\frac{\pi}{2} \end{matrix} \right] \left\{ \begin{matrix} +\frac{\pi}{2} \leftrightarrow '1' \\ -\frac{\pi}{2} \leftrightarrow '0' \end{matrix} \right.$

Now, t is even multiple of T_b

then
for $t = 0$ $\left(\pm \frac{\pi}{2T_b} t \right) \Rightarrow 0$

for $t = 2T_b$ $\hookrightarrow \pm \pi$

$\hookrightarrow \pm 2\pi = 0$

$t = 4T_b$


So, incremental angle is 0 or $\pm \pi$.

Now t is odd multiple of T_b

then
for $t = T_b$ $\pm \frac{\pi}{2T_b} t = \pm \frac{\pi}{2}$

$t = 3T_b$ $\hookrightarrow \pm \frac{3\pi}{2} =$

So incremental angle is $\pm \frac{\pi}{2}$



2017

Goals


Phase trellis diagram
Consider $\theta(0) = 0$



My Family




My Friends




from here we conclude previous
Statement.



My Health



My Skills



My Money



My Notes