

## Orthogonality b/w signal

Let us define concept of orthogonality of two signal  $f_1(t)$  and  $f_2(t)$ . We can approximate  $f_1(t)$  in terms of  $f_2(t)$


$$\text{as } f_1(t) = C_{12} f_2(t) + f_e(t) \quad t_1 < t \leq t_2$$

where error  $\Rightarrow f_e(t) = f_1(t) - C_{12} f_2(t)$

to minimize the error  $f_e(t)$  is integration over  $t_1$  to  $t_2$ , of mean square error.

$$E = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} [f_e(t)]^2 dt. \text{ where } C_{12} \text{ is the value which minimize the error.}$$

$$\frac{dE}{dC_{12}} = 0 \Rightarrow C_{12} = \frac{\int_{t_1}^{t_2} f_1(t) f_2(t) dt}{\int_{t_1}^{t_2} f_2^2(t) dt}$$



if

$C_{12} = 0$  then two signal are orthogonal,

$$\int_{t_1}^{t_2} f_1(t) f_2(t) dt = 0$$

Notes:

| March 2017 |    |    |    |    |    |    |
|------------|----|----|----|----|----|----|
| S          | M  | T  | W  | T  | F  | S  |
|            |    |    | 1  | 2  | 3  | 4  |
| 5          | 6  | 7  | 8  | 9  | 10 | 11 |
| 12         | 13 | 14 | 15 | 16 | 17 | 18 |
| 19         | 20 | 21 | 22 | 23 | 24 | 25 |
| 26         | 27 | 28 | 29 | 30 | 31 |    |



# orthogonal signal space



My Friends

let,  $x_1(t), x_2(t) \dots x_n(t)$   
mutually orthogonal signal.

$$\int_{t_1}^{t_2} x_j(t) x_k(t) dt = 0 \text{ when } j \neq k$$



My Health

Sunday

Week 7 / Day 43

let **12**

$$\int_{t_1}^{t_2} x_k^2(t) dt = k_k \quad j = k$$



My Skills

let a function  $f(t)$  can be approximated by the this orthogonal signal space by adding the components along mutually orthogonal signals.



My Money

$$f(t) = C_1 x_1(t) + C_2 x_2(t) + \dots + C_n x_n(t) + f_e(t)$$

$$f_e(t) = f(t) - \sum_{r=1}^n C_r x_r(t)$$

February 2017

| S  | M  | T  | W  | T  | F  | S  |
|----|----|----|----|----|----|----|
|    |    |    | 1  | 2  | 3  | 4  |
| 5  | 6  | 7  | 8  | 9  | 10 | 11 |
| 12 | 13 | 14 | 15 | 16 | 17 | 18 |
| 19 | 20 | 21 | 22 | 23 | 24 | 25 |
| 26 | 27 | 28 | 29 | 30 |    |    |

Notes:

Mean square error  $\epsilon = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} [f_e(t)]^2 dt$



My Family

Tuesday

Week 10 / Day 59

28

\*

The component which minimize the mean square error can be found



My Friends

$$\frac{dE}{dc_1} = \frac{dE}{dc_2} = \dots = \frac{dE}{dc_k} = 0$$

Let us consider  $\frac{dE}{dc_k} = 0$ .

as all term do not contain  $c_k$  is zero.



My Health

 $\Rightarrow$ 

$$c_k = \frac{\int_{t_1}^{t_2} f(t) x_k(t) dt}{\int_{t_1}^{t_2} x_k^2(t) dt}$$

$$\therefore \text{if } c_k = 0, \quad \int_{t_1}^{t_2} f(t) x_k(t) dt = 0$$



My Skills

 $k=1, 2, \dots$ 

$f(t)$  is. should be orthogonal to every  $x_k(t)$ ; then it called Orthogonal space



My Money





My Family

Monday

13

Week 8 / Day 44

orthogonal signal  $\Rightarrow \int_0^t \phi_1(t) \phi_2(t) dt = 0$ 

✓ orthonormal signal  $\Rightarrow \int_0^t \phi_i(t) dt = 1$   
 Basically energy of a signal unity.



My Friends

Geometrical representation of Signal.

Consider M no. of possible Tx signals  
 $\hookrightarrow x_1(t), x_2(t) \dots x_M(t)$



My Health

N no of orthonormal basis function  $\phi_j(t)$   
 $N \leq M$

We can represent any signal  $x_i(t)$ 

My Skills

as linear combination of orthonormal basis for:

$$\therefore x_i(t) = x_{i1} \phi_1(t) + x_{i2} \phi_2(t) + \dots + x_{iM} \phi_n(t)$$

$$\therefore x_i(t) = \sum_{j=1}^M x_{ij} \phi_j(t) \quad i=1, 2, \dots, M$$



My Money

↓  
Transmitted signal.

Notes:

| March 2017 |    |    |    |    |    |    |
|------------|----|----|----|----|----|----|
| S          | M  | T  | W  | T  | F  | S  |
|            |    |    | 1  | 2  | 3  | 4  |
| 5          | 6  | 7  | 8  | 9  | 10 | 11 |
| 12         | 13 | 14 | 15 | 16 | 17 | 18 |
| 19         | 20 | 21 | 22 | 23 | 24 | 25 |
| 26         | 27 | 28 | 29 | 30 | 31 |    |



# 2017

My Family


Tuesday

Week 8 / Day 45

## 14

Transmitter

My Friends



Receiver! Now the received  $x_i(t)$  multiply  $\phi_k(t)$  and Pass this through Integrator

My Health

$$\int_0^T x_i(t) \phi_k(t) dt$$

My Skills

$$\sum_{j=1}^M x_{ij} \int_0^T \phi_k(t) \phi_j(t) dt$$

now, if  $j \neq k$ , otherwise = 0

this become  $x_{ik}$

My Money

$$x_{ik} = \int_0^T x_i(t) \phi_k(t) dt$$

Notes:

| February 2017 |    |    |    |    |    |    |
|---------------|----|----|----|----|----|----|
| S             | M  | T  | W  | T  | F  | S  |
|               |    |    | 1  | 2  | 3  | 4  |
| 5             | 6  | 7  | 8  | 9  | 10 | 11 |
| 12            | 13 | 14 | 15 | 16 | 17 | 18 |
| 19            | 20 | 21 | 22 | 23 | 24 | 25 |
| 26            | 27 | 28 |    |    |    |    |






My Family

Wednesday

Week 8 / Day 46

15



My Friends

## Gram - Schmidt Orthogonalization Procedure



My Health

To find how many no. ~~orthogonal~~ <sup>orthonormal</sup> basis fun. is required.

Consider set of  $M$  energy signal  $x_1(t)$ ,  $x_2(t)$ , ...,  $x_M(t)$  are real.

$E =$  find energy of  $x_i(t) = \int_0^T x_i^2(t) dt$



My Skills

~~We assume~~ The first basis fun.  $\phi_1(t) = \frac{x_1(t)}{\sqrt{E}}$

normalised energy signal  $x_1(t)$  as orthonormal signal energy 1

For defining other basis fun., a new intermediate function is introduced,



My Money

Notes:

$$g_i(t) = x_i(t) - \sum_{j=1}^{i-1} x_{ij} \phi_j(t)$$

where  $x_{ij} = \int_0^T x_i(t) \phi_j(t) dt$

| March 2017 |    |    |    |    |    |    |
|------------|----|----|----|----|----|----|
| M          | T  | W  | T  | F  | S  |    |
|            |    | 1  | 2  | 3  | 4  |    |
| 5          | 6  | 7  | 8  | 9  | 10 | 11 |
| 12         | 13 | 14 | 15 | 16 | 17 | 18 |
| 19         | 20 | 21 | 22 | 23 | 24 | 25 |
| 26         | 27 | 28 | 29 | 30 | 31 |    |

$i = 1, \dots, M$   
 $g$  is new basis fun





16

from a given  $g_i(t)$  a new set of basis fun. are defined.



My Friends

$$\phi_i(t) = \frac{g_i(t)}{\sqrt{\int_0^T g_i^2(t) dt}} \quad i=1, 2, \dots, N$$




My Health

Prob

find orthonormal basis fun.




My Skills



Sol:  $\phi_1(t) = \frac{x_1(t)}{\sqrt{E_1}}$  where  $E_1 = \int_0^{T/3} x_1^2(t) dt = \frac{T}{3}$



My Money



| February 2017 |    |    |    |    |    |    |
|---------------|----|----|----|----|----|----|
| S             | M  | T  | W  | T  | F  | S  |
|               |    |    | 1  | 2  | 3  | 4  |
| 5             | 6  | 7  | 8  | 9  | 10 | 11 |
| 12            | 13 | 14 | 15 | 16 | 17 | 18 |
| 19            | 20 | 21 | 22 | 23 | 24 | 25 |
| 26            | 27 | 28 |    |    |    |    |

Notes:

$$= \sqrt{\frac{3}{T}} x_1(t)$$



My Family

Friday

Week 8 / Day 48

we know **17**

$$g_i(t) = x_i(t) = \sum_{j=1}^{i-1} x_{ij} \phi_j(t)$$



My Friends


for,  $i=2$ ,  $g_2(t) = x_2(t) = x_{21} \phi_1(t)$

$$x_{21} = \int_0^T x_2(t) \phi_1(t) dt$$




My Health

$$= \int_0^T \frac{1}{3} \sqrt{\frac{3}{T}} dt$$



$$= \sqrt{\frac{T}{3}}$$



My Skills


$$g_2(t) = 1 \quad 0 \leq t < \frac{T}{3}$$



$$\phi_2(t) = \frac{g_2(t)}{\sqrt{\int_{T/3}^{2T/3} 1 dt}} = \sqrt{\frac{3}{T}} g_2(t) \quad \frac{T}{3} \leq t < \frac{2T}{3}$$



My Money



Notes:

From here  $\phi_1(t)$  and  $\phi_2(t)$  are orthogonal

| March 2017 |    |    |    |    |    |    |
|------------|----|----|----|----|----|----|
| M          | T  | W  | T  | F  | S  | S  |
| 5          | 6  | 7  | 8  | 9  | 10 | 11 |
| 12         | 13 | 14 | 15 | 16 | 17 | 18 |
| 19         | 20 | 21 | 22 | 23 | 24 | 25 |
| 26         | 27 | 28 | 29 | 30 | 31 |    |





My Family

Saturday

i = 3

Week 8 / Day 49

18

$$g_3(t) = x_3(t) - \sum_{j=1}^2 x_{3j} \phi_j(t)$$

$$= g_3(t) = x_3(t) - x_{31} \phi_1(t) - x_{32} \phi_2(t)$$



My Friends

$$x_{31} = \int_0^T x_3(t) \phi_1(t) dt = 0$$



My Health

Sunday


Week 8 / Day 50

19

$$x_{32}(t) = \int_0^T x_3(t) \phi_2(t) dt$$



My Skills



$$= \int_{T/3}^{2T/3} \sqrt{\frac{T}{3}} dt$$


$$T/3 \leq t \leq 2T/3$$

$$x_{32}(t) \phi_2(t) = 1 \quad T/3 \leq t \leq 2T/3$$



My Money

$$g_3(t) = 1 \quad \frac{2T}{3} < t < T$$



| February 2017 |    |    |    |    |    |    |
|---------------|----|----|----|----|----|----|
| S             | M  | T  | W  | T  | F  | S  |
|               |    |    | 1  | 2  | 3  | 4  |
| 5             | 6  | 7  | 8  | 9  | 10 | 11 |
| 12            | 13 | 14 | 15 | 16 | 17 | 18 |
| 19            | 20 | 21 | 22 | 23 | 24 | 25 |
| 26            | 27 | 28 |    |    |    |    |

Notes:






My Family

Monday

20

Week 9 / Day 51



$$\phi_3(t) = \frac{g_3(t)}{\sqrt{\int_{2T/3}^T |g_3(t)|^2 dt}} = \sqrt{\frac{3}{T}} g_3(t)$$



My Friends

Now,  $L = 4$ ,  $g_4(t) = x_4(t) - \sum_{j=1}^3 x_{4j} \phi_j(t)$

$$\Rightarrow g_4(t) = x_4(t) - x_{41} \phi_1(t) - x_{42} \phi_2(t) - x_{43} \phi_3(t)$$

$$0 < t < T/3 \quad x_{41} = \frac{1}{\sqrt{3}} \int_0^{T/3} x_4(t) \phi_1(t) dt = \sqrt{\frac{T}{3}}$$

$$T/3 < t < 2T/3 \quad x_{42} = \frac{1}{\sqrt{3}} \int_{T/3}^{2T/3} x_4(t) \phi_2(t) dt = \sqrt{\frac{T}{3}}$$

$$2T/3 < t < T \quad x_{43} = \frac{1}{\sqrt{3}} \int_{2T/3}^T x_4(t) \phi_3(t) dt = \sqrt{\frac{T}{3}}$$



My Skills

$$\text{Now, } x_{41} \phi_1(t) = 1 \quad 0 < t < T/3$$

$$x_{42} \phi_2(t) = 1 \quad T/3 < t < 2T/3$$

$$x_{43} \phi_3(t) = 1 \quad 2T/3 < t < T$$

$$g_4(t) = 0 \quad 0 < t < T$$



My Money

$$\phi_4(t) = 0 \quad \text{no orthonormal fn!}$$

Notes:

| March 2017 |    |    |    |    |    |    |
|------------|----|----|----|----|----|----|
| S          | M  | T  | W  | T  | F  | S  |
|            |    |    | 1  | 2  | 3  | 4  |
| 5          | 6  | 7  | 8  | 9  | 10 | 11 |
| 12         | 13 | 14 | 15 | 16 | 17 | 18 |
| 19         | 20 | 21 | 22 | 23 | 24 | 25 |
| 26         | 27 | 28 | 29 | 30 | 31 |    |



# 2017



My Family

Tuesday

Week 9 / Day 52

## 21

So, 1st signal may represent only one basis function,  $x_1(t) = \sqrt{\frac{1}{3}} \phi_1(t)$



My Friends

2nd signal may represent two basis fun. and so on.

$$x_2(t) = \sqrt{\frac{1}{3}} \phi_1(t) + \sqrt{\frac{1}{3}} \phi_2(t)$$

$$x_3(t) = \sqrt{\frac{1}{3}} \phi_1(t) + \sqrt{\frac{1}{3}} \phi_2(t) + \sqrt{\frac{1}{3}} \phi_3(t)$$



My Health

$$x_4(t) = \sqrt{\frac{1}{3}} \phi_1(t) + \sqrt{\frac{1}{3}} \phi_2(t) + \sqrt{\frac{1}{3}} \phi_3(t)$$