

PMx Weekly Digest: May 30, 2026

Content Source: Full Text

Paper 1: Managing Drug–Drug Interactions Involving the Non-Prescription Opioid Loperamide

Bibliographic Information

- **Title:** Managing Drug–Drug Interactions Involving the Non-Prescription Opioid Loperamide Through Physiologically Based Pharmacokinetic Modeling
- **Authors:** Zhou Z, Anderson GR, Li M, Zamek-Gliszczyński MJ, Zhao P, Duan J, Paine MF
- **Journal:** CPT Pharmacometrics Syst Pharmacol. 2026 Feb 1. doi:10.1002/psp4.70148
- **DOI:** 10.1002/psp4.70148

Paper Type: POPK_PKPD

Quick Take

This study presents a novel, verified PBPK model for loperamide, a widely used nonprescription opioid with abuse potential and cardiotoxicity risk at high doses. The model accurately describes loperamide PK across 10 clinical studies and predicts drug–drug interactions with six CYP/P-gp inhibitors within 25 % of observed AUC ratios. Pharmacometricians involved in PBPK modeling, drug interaction assessment, or opioid safety will find this work valuable for understanding loperamide disposition and guiding risk mitigation strategies.

Executive Summary

Loperamide, a widely used nonprescription opioid, can cause life-threatening cardiac toxicity when taken at ultra-high doses, especially with inhibitors of CYP enzymes or P-glycoprotein. This study developed a mechanistic full-body PBPK model for loperamide using the Simcyp simulator, integrating experimentally determined plasma protein binding ($f_{u,plasma} = 0.03$) and blood-to-plasma ratio (0.71), literature physicochemical data, and retrograde analysis from clinical oral PK profiles. The model accurately predicted loperamide exposure at single doses of 4, 8, and 16 mg (AUC and C_{max} within 0.61- to 1.41-fold of observed values from 10 clinical studies) and successfully captured drug–drug interactions with quinidine, ritonavir, gemfibrozil, itraconazole, gemfibrozil+itraconazole, and abemaciclib (predicted AUC ratios within 0.78- to 1.09-fold of observed ratios). Notably, the model estimated loperamide oral bioavailability at ~19 %—tenfold higher than previously reported—and refined the fractional metabolic contributions of CYP3A4 (53 %), CYP2C8 (46 %), and CYP2D6 (1 %). This verified PBPK model provides a quantitative foundation for predicting untested DDI scenarios and supporting safety assessments of loperamide misuse.

Scientific Context & Motivation

Loperamide misuse at ultra-high doses (70–200 mg/day) has increased due to restricted access to prescription opioids, leading to case reports of cardiac dysrhythmias and deaths. Co-adminis-

tration of CYP3A4, CYP2C8, or P-gp inhibitors can further elevate loperamide systemic exposure, but clinical DDI studies at such doses are unethical. Fundamental knowledge gaps exist: the actual oral bioavailability of loperamide (previously cited as ≤ 0.02 without evidence), the fractional metabolic contributions of individual CYP enzymes, and the relative roles of intestinal vs. hepatic first-pass extraction. This study addresses these gaps by developing and verifying a mechanistic PBPK model that integrates in vitro, in silico, and clinical data, enabling quantitative prediction of exposures in clinically intractable scenarios.

✂ Methodological Snapshot

The study employed the Simcyp PBPK Simulator V21 to develop a full-body PBPK model for loperamide. Absorption was handled by the ADAM model, distribution by a full PBPK method with K_p scalar adjustment, and elimination by enzyme kinetics using retrograde estimation of CYP3A4, CYP2C8, and CYP2D6 intrinsic clearances from clinical oral PK data (CL/F derived from two-compartment analysis). Intestinal P-gp transport was optimized using the clinical quinidine-loperamide DDI study. Inhibitor models for quinidine, ritonavir, gemfibrozil, itraconazole, and abemaciclib were taken from the Simcyp library, with only the itraconazole K_i against P-gp adjusted. The model was verified by comparing simulated vs. observed concentration-time profiles and relevant PK metrics against twofold (base PK) and 1.25-fold (DDI AUCR) acceptance criteria.

☒ Structural Model Breakdown

The loperamide PBPK model is a whole-body, full PBPK model implemented in Simcyp V21. Absorption: The ADAM model simulates dissolution, absorption, and first-pass gut metabolism and efflux. Effective human jejunal permeability ($P_{eff_{man}} = 12 \times 10^{-4}$ cm/s) was predicted from MDCK II cell permeability (456×10^{-6} cm/s). The fraction absorbed (F_a) was optimized to 0.37 and the fraction escaping gut metabolism (F_g) to 0.93, yielding an oral bioavailability of 0.19. Distribution: Tissue:plasma partition coefficients (K_p) are predicted using Simcyp method 2 (based on logP, pKa, and tissue composition). A K_p scalar of 0.7 was applied to match the V_{ss} (9.6 L/kg) derived from two-compartment analysis of oral PK data. Elimination: Hepatic metabolism is modeled via enzyme kinetics for three CYP isoforms: CYP3A4 ($CL_{int} = 3.507$ μ L/min/pmol isoform), CYP2C8 (20.776 μ L/min/pmol), and CYP2D6 (1.027 μ L/min/pmol). These were calculated retrograde from a total apparent oral clearance ($CL_{tot}/F = 249$ L/h) and fractional metabolism ($fm_{3A4} = 0.53$, $fm_{2C8} = 0.46$, $fm_{2D6} = 0.01$). The active hepatic scalar was set to 1. Renal clearance was set to 0. Transport: Intestinal efflux by P-gp is described with an intrinsic clearance ($CL_{int,T}$) of 250 μ L/min and a relative activity factor (RAF/REF) of 1.5. Biliary clearance was not included.

Detailed Methodological Analysis

Modeling Approach

A whole-body full PBPK model was built in Simcyp V21. Absorption: Advanced Dissolution, Absorption, and Metabolism (ADAM) model with $Peff_{man}$ predicted from MDCK II permeability (456×10^{-6} cm/s). Distribution: Full PBPK model using method 2 with a Kp scalar (0.7) to achieve target V_{ss} . Elimination: Enzyme kinetics module with three CYP pathways (CYP3A4, CYP2C8, CYP2D6) using retrograde-estimated CL_{int} values. Transporter: P-gp efflux in the intestine ($CL_{int,T} = 250 \mu\text{L}/\text{min}$). Renal clearance set to zero. Inhibitor models (quinidine, ritonavir, gemfibrozil, itraconazole, abemaciclib) taken from the Simcyp library; K_i of itraconazole against P-gp was adjusted to $1.0 \mu\text{M}$ for the gemfibrozil+itraconazole simulation, and hydroxyitraconazole added as an additional inhibitor.

Data Sources

Ten clinical PK studies (one at 4 mg, four at 8 mg, five at 16 mg loperamide-HCl) were sourced from NDA reviews (Clinpharm Review 21,140 and 21,855) and published literature, digitized using WebPlotDigitizer. Seven DDI studies involving quinidine, ritonavir (two studies), gemfibrozil, itraconazole, gemfibrozil+itraconazole, and abemaciclib were identified from the Certara Drug Interaction Database and literature. Experimental data for fu_{plasma} (equilibrium dialysis) and CB/CP (in vitro blood partitioning) were generated in-house.

Estimation Methods

Retrograde analysis in Simcyp estimated intrinsic clearances for CYP3A4 ($3.507 \mu\text{L}/\text{min}/\text{pmol}$), CYP2C8 ($20.776 \mu\text{L}/\text{min}/\text{pmol}$), and CYP2D6 ($1.027 \mu\text{L}/\text{min}/\text{pmol}$) from a total apparent oral clearance ($CL_{tot}/F = 249 \text{L}/\text{h}$) and fractional contributions ($fm_{3A4} = 0.53$, $fm_{2C8} = 0.46$, $fm_{2D6} = 0.01$). The P-gp intestinal intrinsic clearance ($250 \mu\text{L}/\text{min}$) and absorption parameters ($F_a = 0.37$, $F_g = 0.93$) were manually optimized by fitting the quinidine DDI data. The Kp scalar (0.7) was adjusted to match the V_{ss} ($9.6 \text{L}/\text{kg}$) from two-compartment analysis in Phoenix WinNonlin.

Model Evaluation

Model verification compared simulated concentration–time profiles (10 trials with virtual healthy volunteers) against observed digitized data. Acceptance criteria: predicted AUC and C_{max} within twofold of observed for base PK, and predicted AUC ratios ($AUCR$) within 1.25-fold for DDI studies. Sensitivity analysis for P-gp CL_{int} was performed (Figure S1).

Covariate Analysis

No formal covariate analysis was performed. Virtual population was the Simcyp healthy adult library with default demographic variability. Dose-exposure proportionality was assessed using a power model ($y = ax^b$) on AUC and C_{max} across three dose levels (4, 8, 16 mg), with 95% confidence intervals for the exponent confirming dose proportionality.

Statistical Rigor Assessment

The study uses a deterministic PBPK framework rather than a full statistical estimation; most parameters are recovered via retrograde analysis and manual optimization, without formal confidence intervals. Model evaluation uses fold-difference criteria (2-fold for PK endpoints, 1.25-fold for DDI AUCR), which are standard in PBPK verification but do not provide a measure of statistical uncertainty. No formal goodness-of-fit plots, residual diagnostics, or bootstrap/uncertainty quantification were performed. The dose-proportionality assessment includes a 95 % confidence interval for the power model exponent, providing statistical support. Sample sizes in the clinical studies used for verification are typical ($n = 8\text{--}12$ per trial) but formal sample size justification is absent. Overall, the approach is rigorous within the PBPK modeling paradigm but lacks the formal statistical inference of population PK models.

Key Findings

The PBPK model successfully predicted loperamide PK at 4, 8, and 16 mg with AUC and C_{max} within 0.61–1.41-fold of observed values from 10 clinical studies. For DDI predictions with quinidine, ritonavir, gemfibrozil, itraconazole, gemfibrozil+itraconazole, and abemaciclib, the predicted AUC ratios were within 0.78–1.09-fold of observed ratios, satisfying the prespecified 0.75–1.25-fold acceptance criterion. The model recovered an oral bioavailability estimate of 0.19 (tenfold higher than the previously cited ≤ 0.02), and refined fractional metabolic contributions: 53 % CYP3A4, 46 % CYP2C8, and 1 % CYP2D6. Dose-exposure proportionality (exponent 95 % CI including 1) was confirmed across the therapeutic range. The intestinal P-gp intrinsic clearance was optimized to 250 $\mu\text{L}/\text{min}$.

Clinical & Regulatory Implications

The model can guide loperamide dosing decisions when co-administered with CYP3A4 or CYP2C8 inhibitors. For example, the gemfibrozil+itraconazole combination increased loperamide AUC 12.6-fold, indicating severe risk of toxicity; such combinations should be avoided at high loperamide doses. The model may support regulatory labeling updates for DDI warnings and help design risk mitigation strategies for loperamide misuse. The revised bioavailability estimate also implies that supratherapeutic doses may produce systemic exposures higher than previously assumed, reinforcing caution in patients taking any CYP3A4/2C8 or P-gp inhibitor.

Strengths & Limitations

Strengths

- Mechanistic full-body PBPK model incorporating ADAM absorption and P-gp transport.
- Experimental determination of $f_{u_{plasma}}$ (0.03) and CB/CP (0.71) under simulated physiological conditions.
- Model verification against 10 independent clinical PK studies and 7 DDI studies, covering a range of doses and inhibitor types.

- Recovery of previously unknown bioavailability estimate (~ 0.19) and refined CYP fractional contributions.
- Use of multiple DDI studies with inhibitors of varying specificity to triangulate metabolic pathways (CYP3A4, CYP2C8, CYP2D6).
- Practical clinical relevance: addresses a critical public health issue (loperamide misuse and cardiotoxicity).
- Detailed reporting of input parameters and assumptions in Table 2, enabling reproducibility and critique.

Limitations (Acknowledged by Authors)

- Lack of differentiation among various loperamide formulations (caplets, chewable tablets, soft gelatin capsules, Imodium A-D); all treated as one formulation.
- Itraconazole formulation choice (fasted solution rather than fed capsule) may explain overprediction of itraconazole-loperamide DDI (AUC overpredicted by 45 %).
- Model validated only up to 16 mg; linearity beyond this dose is unknown.
- No selective intestinal CYP3A inhibitor (e.g., grapefruit juice) was available in the Simcyp library to differentiate intestinal vs. hepatic CYP3A contribution.
- Biliary clearance not considered due to lack of data.
- Minor CYP2B6 pathway not incorporated.
- P-gp intestinal CL_{int} optimized based on a single DDI study (quinidine) and sensitivity analysis.

Limitations (Expert Review)

- Reliance on Simcyp library inhibitor models without independent verification of those models.
- Manual optimization of F_a , F_g , and P-gp CL_{int} against one DDI study; no formal estimation with confidence intervals.
- No formal uncertainty quantification or global sensitivity analysis for optimized parameters.
- Absorption parameters (F_a , F_g) are not independently verified; they were inferred from quinidine DDI data.
- The model does not evaluate the impact of food or formulation differences, which may affect absorption and DDI magnitude.
- No assessment of inter-individual variability in transporter expression beyond Simcyp defaults.
- The retrograde analysis assumes all clearance is metabolic and assigns fractions based on a single in vitro study; other pathways (e.g., direct glucuronidation) are ignored.

Generalizability

The model is verified for healthy adults at single doses up to 16 mg and for DDI with the specific inhibitors tested. Generalizability to supratherapeutic doses (> 16 mg), chronic dosing, special populations (elderly, hepatic/renal impairment, pediatric), or untested inhibitor combinations is unknown. However, the mechanistic structure provides a framework for future extension to these scenarios.

Key Equations

Dose-exposure proportionality model

$$y = a \times x^b$$

Power model used to assess dose proportionality of loperamide AUC and Cmax across 4–16 mg dose range. Exponent $b = 1$ indicates dose-proportional pharmacokinetics.

Figures & Tables

- **Figure 1:** Flowchart illustrating the iterative model development workflow, with blue boxes for model building data and green boxes for verification data.
 - *Significance:* Provides a clear overview of the modeling strategy, showing the sequence of base model development, verification with 10 PK studies, DDI model building, and final verification with 7 DDI studies.
- **Figure 2:** Observed (points) and PBPK model predicted (lines/shaded areas) loperamide concentration–time profiles when co-administered with itraconazole, gemfibrozil+itraconazole, quinidine, ritonavir (600 mg single dose and 200 mg BID), and abemaciclib.
 - *Significance:* Demonstrates the model’s ability to capture the time course of loperamide under various DDI conditions, including complex three-inhibitor (gemfibrozil+itraconazole+hydroxyitraconazole) scenarios.
- **Figure 3:** Observed and predicted loperamide concentration–time profiles at the three single-dose levels (4 mg, 8 mg, 16 mg) in the absence of inhibitors.
 - *Significance:* Visual confirmation of the base model’s accuracy across the full therapeutic dose range, with predictions falling within the 5th–95th percentile simulation intervals.
- **Figure 4:** Scatter plots of predicted vs. observed AUC ratio ($AUCR$) and C_{max} ratio ($C_{max}R$) for all DDI studies, with dashed lines showing 1.25-fold acceptance boundaries.
 - *Significance:* Concise quantitative summary of DDI prediction performance; all points fall within the prespecified acceptance criteria, validating the model’s utility for prospective simulations.
- **Table 1:** Comparison of observed vs. predicted AUC and C_{max} for the 10 clinical PK studies used for base model verification (4 mg, 8 mg, and 16 mg doses).
 - *Significance:* Provides a comprehensive fold-difference assessment, confirming that all predicted AUCs are within 0.84- to 1.41-fold and Cmax within 0.61- to 1.05-fold of observed values.
- **Table 2:** Complete list of input parameters for the loperamide PBPK model, including physicochemical properties, absorption (ADAM), distribution (full PBPK), elimination (enzyme kinetics), and P-gp transport parameters, with sources.
 - *Significance:* Essential for reproducibility; documents all model assumptions and parameter origins, including experimentally determined values ($f_{u_{plasma}}$, CB/CP) and optimized parameters (CYP CL_{int} values, P-gp CL_{int} , Kp scalar).
- **Table 3:** Predicted vs. observed AUC and C_{max} for each DDI study, with deviations and observed vs. predicted AUC ratios ($AUCR$).
 - *Significance:* Quantifies the model’s DDI prediction accuracy, showing that all predicted AUCRs are within 0.78–1.09-fold of observed values (meeting the 0.75- to 1.25-fold criterion).

- **Table S1:** Power model results (exponent b and 95 % CI) for dose-exposure proportionality assessment across 4–16 mg.
 - *Significance:* Statistically supports the assumption of dose-proportional PK over the therapeutic range, with 95 % CI including 1 for both AUC and C_{max} .
-

Code & Reproducibility Assessment

The model was developed using the commercial Simcyp PBPK Simulator V21 (Certara). All input parameters are listed in Table 2, and experimental methods for $f_{u_{plasma}}$ and CB/CP are described. No custom code or raw data files are publicly shared; the model is partially reproducible given access to Simcyp and the same clinical data.

Supplementary Materials

Supplementary materials include Figure S1 (local sensitivity analysis for P-gp intestinal CL_{int} , showing the impact on loperamide AUC ratio for the quinidine DDI) and Table S1 (power model results for dose proportionality assessment, with exponent b and 95 % CI for AUC and C_{max}).

Future Directions

Several important extensions are identified: (1) incorporate saturable metabolism and transport (V_{max} , K_m) to assess nonlinear PK at supratherapeutic doses (> 16 mg); (2) adapt the model to special populations such as the elderly, patients with hepatic impairment, or paediatric patients; (3) link the PBPK model to a cardiac toxicity PD model (e.g., QTc prolongation) to predict concentration-response for dysrhythmia risk; (4) include a selective intestinal CYP3A inhibitor (e.g., grapefruit juice) study to differentiate gut vs. liver contribution to first-pass metabolism; (5) incorporate biliary clearance if experimental data become available; and (6) perform formal uncertainty quantification and global sensitivity analysis on optimized parameters.

Expert Commentary

This work convincingly demonstrates the power of PBPK modeling to recover fundamental PK parameters—especially oral bioavailability and fractional metabolism—that are nearly impossible to obtain experimentally for drugs lacking an intravenous formulation. The tenfold revision of loperamide bioavailability (from ≤ 0.02 to 0.19) has profound implications for interpreting the magnitude of DDIs and safety margins. A methodological highlight is the use of multiple clinical DDI studies with inhibitors of different specificity to triangulate CYP contributions, a best practice in PBPK model refinement. However, the manual optimization of P-gp CL_{int} and absorption parameters (F_a , F_g) against a single DDI study introduces subjectivity; future work should incorporate formal sensitivity analysis and uncertainty quantification. The reliance on Simcyp library inhibitor models without independent verification is common but limits the ability to attribute prediction errors. Overall, this is a well-executed, clinically relevant study that addresses an urgent public health question.

Bottom Line

This novel, verified whole-body PBPK model for loperamide accurately describes its pharmacokinetics across 4–16 mg doses and reliably predicts clinically significant drug–drug interactions with CYP3A4, CYP2C8, and P-gp inhibitors. The model revises loperamide’s oral bioavailability upward to $\sim 19\%$ (from previous estimates of $\leq 2\%$) and establishes fractional metabolic contributions of 53 % CYP3A4, 46 % CYP2C8, and 1 % CYP2D6. Pharmacometricians can use this model to simulate untested DDI scenarios (including high-dose misuse) and to inform risk mitigation strategies for cardiac toxicity without exposing subjects to dangerous exposures.

☒ Figures




Figure 1: Workflow for loperamide model development. Blue boxes indicate the data used for model development, and green boxes indicate the data used for model verification




Figure 2: Observed and PBPK model-predicted loperamide concentration–time profiles in the presence of different inhibitors: (A) 4 mg loperamide•HCl + itraconazole (100 mg




Figure 3: Observed and PBPK model-predicted loperamide concentration–time profiles at three loperamide•HCl doses (A, 4 mg; B, 8 mg; C, 16 mg). Grey shaded areas represent




Figure 4: Predicted vs. observed (A) AUCR (AUC of loperamide in the presence and absence of inhibitor) and (B) $C_{\max}R$ ($C_{\max}$ of loperamide in the presence and absence of inhi