

PMx Weekly Digest: May 26, 2026

Content Source: Full Text

Paper 1: Assessing Covariate Clinical Relevance in High-Dimensional Population PK Modelin

Bibliographic Information

- **Title:** Assessing Covariate Clinical Relevance in High-Dimensional Population PK Modeling: A Comparison of SCM+, FFEM, and FREM Approaches
- **Authors:** Philippe M, et al.
- **Journal:** CPT: Pharmacometrics & Systems Pharmacology, 2026, volume/pages TBD
- **DOI:** <https://doi.org/10.1002/psp4.70232>

Paper Type: METHODOLOGY

Quick Take

This study compares three covariate modeling methods (SCM+, FFEM, FREM) for assessing covariate clinical relevance (CCR) in high-dimensional population PK settings. Using simulated datasets with varying covariate effect sizes and correlations, it finds that full modeling approaches (FFEM, FREM) provide more comprehensive CCR evaluation but with higher uncertainty, while SCM+ yields parsimonious models with limited power for small effects. The results support combining approaches for robust covariate-driven dose decisions.

Executive Summary

The study evaluated covariate clinical relevance (CCR) assessment using SCM+, FFEM, and FREM within a high-dimensional covariate framework (19 covariates, 70 covariate-parameter relationships) with varying effect sizes and correlations. A clinical trial simulation inspired by dupilumab PK was conducted (200 datasets, 300 patients each). SCM+ showed strong selection performance for medium/high effects but missed up to 9% of small effects and could not evaluate CCR for non-selected covariates. FFEM and FREM enabled comprehensive CCR evaluation across all predefined relationships, classifying covariates as relevant (R), non-relevant significant (NRS), insufficient information significant (IIS), or insufficient information non-significant (IINS). However, they exhibited higher estimation uncertainty, particularly for poorly informed parameters like Q/F , and produced more false-positive R classifications for covariates without simulated effects (up to 19% with FREM). The study recommends combining SCM+ for parsimonious predictive modeling with FFEM/FREM for thorough CCR documentation, aligning with regulatory expectations for model-informed drug development.

Scientific Context & Motivation

Covariate analysis is central in population PK modeling to explain inter-individual variability and support dose optimization. Covariate clinical relevance (CCR) is commonly evaluated using forest plots comparing covariate effect ratios (CER) to a reference interval [0.8–1.25]. Two main families

of covariate modeling methods exist: selection methods (e.g., SCM, SCM+) and full modeling methods (e.g., FFEM, FREM). Previous comparisons focused on covariate selection performance and estimation accuracy, but rarely assessed CCR. This study addresses the need to evaluate CCR assessment in high-dimensional settings with varying effect sizes and correlations, using a simulation framework inspired by a real dupilumab PK study.

✂ Methodological Snapshot

The study used a simulation framework with 200 datasets of 300 patients each, based on a 2-compartment PK model for dupilumab with 5 parameters (KA , CL/F , V_c/F , Q/F , V_p/F) and 19 covariates (12 continuous, 7 categorical). Covariate effects were simulated at three levels (small, medium, high) across 12 covariate-parameter relationships, with correlations ranging from null to high (up to 0.8). Three methods were compared: SCM+ (stepwise covariate model with forward/backward selection using LRT), FFEM (full fixed effects model estimating all covariate effects simultaneously), and FREM (full random effects model treating covariates as random variables with a full variance-covariance matrix). CCR was assessed using forest plots with 90% CI for covariate effect ratios (CER) relative to a reference interval [0.8–1.25], classifying covariates as relevant (R), non-relevant significant (NRS), insufficient information significant (IIS), or insufficient information non-significant (IINS). Estimation was performed in NONMEM/PSN using FOCEI (SCM+, FFEM) or IMPMAP (FREM). Results were evaluated using relative estimation error (REE), relative standard error (RSE), and classification accuracy compared to a reference model (RM).

⊠ Statistical Framework

The statistical framework combines likelihood-based estimation (FOCEI for SCM+ and FFEM, IMPMAP for FREM) with clinical relevance criteria derived from bioequivalence standards. Covariate effects are estimated as fixed effects (SCM+, FFEM) or random effects (FREM). CCR classification uses 90% confidence intervals for covariate effect ratios (CER) relative to a reference range [0.8–1.25], corresponding to a $\pm 20\%$ change on the log scale. Statistical significance is assessed using 95% CI excluding 1 (for continuous covariates) or likelihood ratio tests (for SCM+ selection). The framework distinguishes four CCR categories: R (CER CI_{90} entirely outside [0.8–1.25]), NRS (CI_{90} outside but CI_{95} includes 1), IIS (CI_{90} straddles reference range with statistical significance), IINS (CI_{90} straddles without significance). This approach integrates both clinical relevance and statistical uncertainty.

Estimator Behavior

Parameter estimates were generally unbiased across methods, with REE medians centered on zero for covariates with simulated effects. Precision varied: SCM+ showed lower RSE for selected covariates (mostly $< 30\%$), while FFEM and FREM exhibited higher uncertainty, particularly for Q/F and small/no-effect covariates (RSE often $> 30\%$). FREM showed the widest RSE distributions, especially for categorical covariates. For covariates without simulated effects, SCM+ mostly

did not select them (NRNS), while FFEM/FREM classified them as NRNS or IINS with some false positives (R up to 19% with FREM). Estimation runtime was fastest for FFEM (median ~ 1 h), followed by FREM (~ 7 h) and SCM+ (~ 15 h).

Validation Design

The validation design used 200 simulated datasets with known true covariate effects, allowing direct comparison of CCR classification accuracy against a reference model (RM). Covariate effects were simulated at three levels (small, medium, high) across 12 relationships, with varying correlations. Performance was assessed using relative estimation error (REE), relative root mean square error (RRMSE), and classification proportions (R, NRS, IIS, IINS) compared to RM. The study also evaluated the impact of covariate correlation on CCR decisions. This design provides a rigorous benchmark for method comparison in a controlled high-dimensional setting.

Applicability Boundaries

The methods are applicable to population PK covariate analysis in late-phase clinical trials with moderate to high covariate dimensionality (up to ~ 20 covariates). SCM+ works best for medium/high effect sizes but may miss small effects and cannot evaluate non-selected covariates. FFEM and FREM provide comprehensive CCR evaluation but require sufficient data for precise estimation, particularly for poorly informed parameters like intercompartmental clearance. The methods are less reliable for very high-dimensional settings (e.g., genetic covariates) or sparse sampling designs. Covariate correlation up to 0.8 does not substantially compromise SCM+ or FFEM, but FREM shows increased false positives under correlation.

Comparison to Alternatives

Compared to alternative selection methods (e.g., LASSO, backward selection), SCM+ offers a stepwise approach with LRT-based significance testing but lacks clinical relevance evaluation for non-selected covariates. Full modeling methods (FFEM, FREM) provide more comprehensive CCR assessment but with higher estimation uncertainty and potential false positives. The study did not compare to machine learning approaches (e.g., SVM, random forests) or penalized regression (e.g., ALASSO), which could be valuable for high-dimensional pre-selection. The results support combining SCM+ for parsimonious prediction with FFEM/FREM for thorough CCR documentation, as recommended in regulatory guidelines.

Implementation Guidance

Implementation requires NONMEM (version 7.5.1 +) and PsN (version 5.4.0 +). For SCM+, use forward $p = 0.05$ and backward $p = 0.01$ with FOCEI estimation. For FFEM, estimate all covariate effects simultaneously with FOCEI. For FREM, use IMPMAP estimation with a full variance-covariance matrix. Covariate effect bounds should be set to reasonable values (e.g., -10 to 10). Initial estimates for covariate effects should be small (e.g., 0.001). Runtime varies: FFEM is fastest (median ~ 1 h), FREM moderate (~ 7 h), SCM+ slowest (~ 15 h). For CCR evaluation, compute

CER 90% CI from parameter estimates and SE, classify relative to [0.8–1.25]. Combine SCM+ for model selection with FFEM/FREM for comprehensive CCR documentation in regulatory submissions.

☒ Key Findings

1. SCM+ reliably identified medium/high-effect covariates (R in 71–100% of cases) but missed up to 9% of small effects and could not evaluate CCR for non-selected covariates.
 2. FFEM and FREM provided comprehensive CCR evaluation, classifying covariates as R, NRS, IIS, or IINS across all predefined relationships.
 3. Full modeling approaches showed higher estimation uncertainty (RSE often > 30% for small/no effects) and more false-positive R classifications for covariates without simulated effects (up to 19% with FREM).
 4. Covariate correlation did not substantially compromise CCR assessment for SCM+ or FFEM, but FREM showed increased false positives under correlation.
 5. The study supports combining SCM+ for parsimonious prediction with FFEM/FREM for thorough CCR documentation.
-

Strengths & Limitations

Strengths

- Comprehensive simulation framework with 200 datasets, 19 covariates, 70 covariate-parameter relationships, varying effect sizes and correlations
- Direct comparison of three covariate modeling methods (SCM+, FFEM, FREM) for CCR assessment
- Use of clinical relevance criteria (*CER* 90% CI relative to [0.8–1.25]) alongside statistical significance
- Evaluation of estimation precision (RSE, RRMSE) and classification accuracy (CCR categories)
- Practical recommendations for combining methods in regulatory submissions

Limitations (Acknowledged by Authors)

- SCM+ relies solely on likelihood ratio testing and cannot evaluate clinical relevance for non-selected covariates
- Full modeling methods (FFEM, FREM) exhibit higher estimation uncertainty, especially for poorly informed parameters like Q/F
- FREM shows increased false-positive R classifications for covariates without simulated effects, particularly under correlation
- Simulation design used intensive Phase I-style sampling, which may not reflect typical Phase III sparse sampling
- Only diagonal omega matrix was simulated; full omega matrix could alter results

Limitations (Expert Review)

- No multiple-testing correction was applied for CCR evaluation, potentially inflating Type I error
- The study did not explore alternative selection methods (e.g., LASSO, backward selection) or stage-wise filtering approaches

- CCR evaluation was limited to primary PK parameters (CL/F , Vc/F , Q/F , Vp/F); secondary metrics like AUC/C_{max} were not assessed
- The high-dimensional setting (19 covariates) may not fully represent even higher dimensions (e.g., genetic data)
- Runtime comparisons may not generalize to larger datasets or different software implementations

Generalizability

The findings are based on a simulation inspired by dupilumab PK in atopic dermatitis patients, with covariates sampled from NHANES. The methods are generalizable to other monoclonal antibodies and similar PK settings, but caution is warranted for higher-dimensional contexts or sparse sampling designs. The CCR framework itself is broadly applicable to covariate analysis in population PK.

Future Directions

Expert Commentary

Bottom Line

For robust covariate clinical relevance assessment in high-dimensional PK modeling, combine SCM+ for parsimonious predictive model building with FFEM or FREM for comprehensive CCR evaluation across all predefined covariate-parameter relationships. This dual approach supports both efficient dose optimization and thorough regulatory documentation.

☒ Figures




Figure 1: REE of CER, computed using P10 for continuous covariates, for RM, SCM+, FFEM, and FREM approaches; (A) shows results for covariates with a simulated effect order




Figure 2: RRMSE of CER computed using P10 for continuous covariates, for RM, SCM+, FFEM, and FREM approaches; (A) shows results for covariates with a simulated effect orde




Figure 3: CCR decisions for RM, SCM+, FFEM and FREM approaches shown for covariates with a simulated effect ordered by effect size (high—left, medium—center, small—right);



Figure 4: CCR decisions for RM, SCM+, FFEM and FREM approaches shown for covariates with any effect; only proportions 20% are given. AGE, age; ALB, albumin; ALK, alkaline p