PMx Weekly Digest: February 26, 2026

Paper 1: Risk Assessment for Drug-Induced Hyperbilirubinemia: A Mechanistic Approach

Bibliographic Information

- **Title**: Risk Assessment for Drug-Induced Hyperbilirubinemia: A Mechanistic Approach
- **Authors**: Kumar R, Kumar A, Matsuura Y, Weiss J, et al.
- **Journal**: CPT: Pharmacometrics & Systems Pharmacology, 2026
- **DOI**: https://doi.org/10.1002/psp4.70037

Quick Take

A mechanistic systems toxicology model dissecting the interplay between drug inhibition and genetic polymorphisms (UGT1A1, OATP1B1, MRP2) in hyperbilirubinemia. Essential reading for safety scientists and QSP modelers seeking to distinguish benign transporter-mediated bilirubin elevation from true liver injury.

Executive Summary

This study presents a quantitative systems pharmacology (QSP) model of bilirubin metabolism to assess the risk of drug-induced hyperbilirubinemia (DIH). By integrating physiological bilirubin turnover with drug-specific in vitro inhibition data (Ki), the authors successfully modeled the impact of key transporters (OATP1B1, MRP2) and enzymes (UGT1A1). The model was validated against clinical data from genetic disorders (Gilbert's, Dubin-Johnson syndromes) and specific drug case studies (nilotinib, atazanavir). A critical finding is the synergistic sensitivity of OATP1B1 and UGT1A1; reduced UGT1A1 activity significantly amplifies the hyperbilirubinemic effect of OATP1B1 inhibition. This framework offers a mechanistic rationale for interpreting safety signals in early clinical development, particularly for drugs exhibiting off-target transporter effects.

⚡ Methodological Snapshot

The authors developed a deterministic ordinary differential equation (ODE) model using the Cellworks biosimulation platform. The model structure comprises four compartments: blood, liver, bile, and intestine. It accounts for unconjugated bilirubin (UCB) synthesis, albumin binding, hepatic uptake via OATP1B1, conjugation by UGT1A1, and elimination of conjugated bilirubin (CB) via MRP2 (biliary) and MRP3 (sinusoidal). The model was parameterized using literature physiological values and in vitro kinetic data, then validated by simulating disease states (hemolysis, genetic mutations) and clinical drug scenarios.

🏗 Structural Model Breakdown

The model tracks bilirubin flux through four compartments: Blood, Liver, Bile, and Intestine. Key species include Free Unconjugated Bilirubin (FUCB), Albumin-bound UCB, Bilirubin Monoglucuronide (BMG), and Diglucuronide (BDG).

Key processes: 1. **Uptake:** Diffusion and OATP1B1-mediated transport from blood to liver. 2. **Metabolism:** UGT1A1-mediated conjugation of UCB to BMG and BDG. 3. **Elimination:** MRP2-mediated secretion into bile; MRP3-mediated efflux to blood; renal and fecal elimination. 4. **Recirculation:** Enterohepatic circulation of UCB.

Detailed Methodological Analysis

Modeling Approach

Mechanistic ODE-based systems toxicology model developed on the proprietary Cellworks platform. It simulates steady-state and dynamic bilirubin concentrations based on enzyme/transporter kinetics.

Data Sources

Physiological parameters from literature; in vitro inhibition constants (Ki) for nilotinib, nelfinavir, atazanavir, and TAK-875; clinical validation data from healthy volunteers, patients with genetic syndromes (Gilbert's, Dubin-Johnson), and specific drug trials.

Estimation Methods

Parameters were derived from literature and in vitro experiments (bottom-up approach). Sensitivity analysis was performed using steady-state continuation analysis to assess the impact of parameter perturbations.

Model Evaluation

Predictive visual checks comparing simulated steady-state concentrations against observed values in healthy and disease populations. Case study validation comparing predicted vs. observed bilirubin time-courses for nilotinib and atazanavir.

Covariate Analysis

Genetic polymorphisms were modeled mechanistically by reducing the Vmax of specific enzymes/transporters (e.g., reducing UGT1A1 activity by 30-70% to simulate Gilbert's syndrome).

Scientific Context & Motivation

Hyperbilirubinemia is a standard biomarker for drug-induced liver injury (DILI), yet it is confounded by genetic polymorphisms (e.g., Gilbert's syndrome) and benign transporter inhibition. Distinguishing between pathological liver dysfunction and benign accumulation due to transporter blockade is a significant challenge in drug development. Existing models often treat these processes in isolation. This study addresses the gap by creating a holistic systems model that accounts for the dynamic interplay between hepatic uptake, conjugation, and elimination, providing a platform to quantify the contribution of specific genetic variants and drug interactions to observed bilirubin levels.

Statistical Rigor Assessment

As a deterministic systems model, traditional statistical rigor (SE, shrinkage) is less relevant than physiological plausibility. The authors rigorously validated the model against distinct physiological states (hemolysis, genetic defects) before applying it to drug scenarios. Sensitivity analyses were comprehensive, exploring the interaction space between multiple transporters.

Key Findings

1. **Genetic Validation:** The model accurately reproduced bilirubin levels for Gilbert's syndrome (14-42 M UCB) and Dubin-Johnson syndrome (40 M conjugated bilirubin).
2. **OATP1B1 Impact:** A 90% reduction in OATP1B1 activity increased UCB by 1.58-fold and conjugated bilirubin by 2.2-fold.
3. **Synergistic Sensitivity:** The sensitivity of blood bilirubin to OATP1B1 inhibition increases dramatically when UGT1A1 activity is low. With 99% UGT1A1 reduction, OATP1B1 inhibition causes a ~195-fold increase in UCB vs 2-fold in wild-type.
4. **Drug Case Studies:**
   - **Nilotinib:** Hyperbilirubinemia linked to UGT1A1 inhibition; model captured dose-dependency.
   - **Atazanavir:** Predicted dose-dependent increase in UCB consistent with clinical data.
   - **TAK-875:** Demonstrated that uncertainty in tissue accumulation (free drug concentration) is a major driver of risk; 10x tissue accumulation predicted a 24-fold increase in conjugated bilirubin.

Clinical & Regulatory Implications

The model supports the use of bilirubin as a biomarker for OATP1B1 and UGT1A1 activity. It suggests that patients with UGT1A1 polymorphisms (e.g., Gilbert's) are disproportionately sensitive to drugs inhibiting OATP1B1. The framework can be used to set exclusion criteria or dose adjustments in clinical trials for drugs with known transporter liabilities.

Strengths & Limitations

Strengths

- Mechanistic integration of multiple transporters (OATP, MRP2) and enzymes (UGT1A1).
- Successful validation against distinct genetic disease phenotypes.
- Demonstration of non-linear interactions between pathway components (e.g., UGT1A1 status altering OATP1B1 sensitivity).
- Practical application to withdrawn drugs (TAK-875) to elucidate toxicity mechanisms.

Limitations (Acknowledged by Authors)

- Empirical modeling of free intracellular drug concentrations is limited.
- Does not account for hepatocellular diseases like autoimmune hepatitis.
- Lack of direct human tissue sampling data for validation of intracellular concentrations.

Limitations (Expert Review)

- Reliance on proprietary software (Cellworks) may hinder external reproducibility.
- Assumes linear scaling of in vitro Ki values to in vivo inhibition without accounting for complex transporter kinetics (e.g., time-dependent inhibition) in detail.

Generalizability

Highly generalizable for small molecule drugs where in vitro transporter inhibition data is available. Applicable across therapeutic areas for DILI risk assessment.

Figures & Tables

- **Table 1**: Comparison of model-simulated steady-state bilirubin concentrations against observed values for Control, Gilbert's syndrome, Dubin-Johnson syndrome, and Hemolytic disorders.
  - *Significance*: Validates the physiological structure of the model by proving it can reproduce distinct pathological states solely by modifying specific parameters.
- **Figure 3**: Sensitivity analysis surface plots showing the impact of varying MRP2 and OATP1B1 activity on Total Unconjugated Bilirubin (TUCB) at different levels of UGT1A1 activity.
  - *Significance*: Crucial visualization of the non-linear interaction between pathways; shows that OATP1B1 sensitivity is dependent on UGT1A1 baseline function.
- **Figure 4**: Simulation of Nilotinib-induced hyperbilirubinemia, comparing simulated vs observed Total Bilirubin across different dosing regimens.
  - *Significance*: Demonstrates the model's clinical utility in predicting adverse events for a specific kinase inhibitor with known UGT1A1 liability.
- **Figure 6**: Impact of TAK-875 tissue accumulation (1x, 5x, 10x) on predicted bilirubin levels.
  - *Significance*: Highlights the critical sensitivity of the model to intracellular drug concentration assumptions, explaining potential discrepancies between in vitro safety and clinical toxicity.

Supplementary Materials

File S1 contains model details, key assumptions, parameterization methodologies, and specific parameter values. Table S1 provides the specific parameter values used. Tables S6 and S7 provide drug-specific parameters (plasma concentrations, Ki values).

Future Directions

Future work should incorporate immune-mediated hepatocellular injury mechanisms to broaden the scope beyond transporter-mediated hyperbilirubinemia. Better methods for predicting intracellular free drug concentrations (e.g., PBPK integration) would refine the risk assessment for highly protein-bound drugs like TAK-875.

Expert Commentary

This paper represents a solid advancement in 'Systems Toxicology.' The most valuable insight is the quantitative demonstration of the 'multiple hit' hypothesis—that a drug with mild OATP1B1 inhibition might be safe in a wild-type population but cause significant signals in patients with reduced UGT1A1 activity. This underscores the importance of pharmacogenetic stratification in early phase trials. The analysis of TAK-875 is particularly instructive for modelers, highlighting how sensitive toxicity predictions are to the 'Kp,uu' (unbound partition coefficient) assumption. It serves as a reminder that without accurate intracellular concentrations, predicting transporter-mediated DILI remains challenging.

Bottom Line

Kumar et al. provide a validated mechanistic tool for de-risking hyperbilirubinemia signals in drug development. By quantifying the interplay between OATP1B1, MRP2, and UGT1A1, the model helps pharmacometricians distinguish between benign transporter inhibition and potential liver injury, while highlighting the critical influence of patient genetics (e.g., Gilbert's syndrome) on safety margins.

📊 Figures



Figure 1: FIGURE 1: Schematic bilirubin metabolism model network. The model includes four compartments: Blood, liver, bile, and intestine. Bilirubin is synthesized as free UCB in th





Figure 2: FIGURE 2: Simulated steady-state levels of unconjugated and conjugated bilirubin in blood; (A) effect of albumin–bilirubin KD on total and free blood unconjugated bilirubi



Figure 3: FIGURE 3: Sensitivity analysis of MRP2 and OATP1B1 at different UGT1A1 activity levels on total blood UCB. (A) results of steady-state continuation analysis by varying MRP





Figure 4: FIGURE 4: Nilotinib-induced hyperbilirubinemia. (A) Comparison of simulated and observed changes in total bilirubin for different trough plasma concentrations of nilotinib



Figure 5: FIGURE 5: Model evaluation of drug-induced hyperbilirubinemia with clinical doses of different drugs. (A) Nelfinavir; (B) Atazanavir; (C) TAK-875. The fold change was calc



(B)



Figure 6: FIGURE 6: Effect of free TAK-875 concentration at different levels of tissue accumulation (1×, 5×, and 10× compared with the plasma concentration). (A) Total blood unconju