

PMx Weekly Digest: June 17, 2026

Content Source: Full Text

Paper 1: Cellular Heterogeneity in Drug Uptake Amplifies Pharmacodynamic Variability: A S

Bibliographic Information

- **Title:** Cellular Heterogeneity in Drug Uptake Amplifies Pharmacodynamic Variability: A Stochastic PK-PD Analysis
- **Authors:** Nhung Hong-Thi Duong, Tuan N. Do, Linh T. Nguyen, Thao T.-N. Nguyen, Khoa Q. Pham
- **Journal:** CPT: Pharmacometrics & Systems Pharmacology, 2026, DOI: 10.1002/psp4.70282
- **DOI:** <https://doi.org/10.1002/psp4.70282>

Paper Type: POPK_PKPD

Quick Take

This paper presents a stochastic multiscale PK-PD framework that incorporates cellular heterogeneity in drug transporter expression. Using cobimetinib as an exemplar, the authors demonstrate that such heterogeneity, when coupled with nonlinear pharmacodynamics, can amplify outcome variability by up to 1000-fold and produce treatment failure in 36% of simulated populations despite adequate systemic exposure. The work challenges traditional mean-field PK-PD models and advocates for stochastic cell-level modeling in drug development.

Executive Summary

Traditional PK-PD models assume cellular homogeneity, but clinical observations show substantial response variability even with similar plasma exposure. This study hypothesizes that cellular heterogeneity in drug transporter expression, combined with nonlinear dose-response relationships, amplifies microscopic variability into population-level outcome variability. Using a multiscale stochastic framework with cobimetinib as a model drug, the authors show that log-normally distributed transporter expression leads to broad extinction time distributions, with a 22-fold variance amplification and 36% treatment failure under intermittent dosing. Mean-field models systematically underpredict extinction times by 60-81%. Sensitivity analysis identifies transporter expression coefficient of variation as the key determinant of outcome variability. The findings support incorporating stochastic cell-level modeling into PK-PD frameworks and suggest that single-cell transporter measurements could stratify patients.

Scientific Context & Motivation

Classical PK-PD models use population-averaged approaches that address inter-patient variability but ignore intra-tumor cellular heterogeneity in drug transporter expression. Single-cell measurements reveal coefficients of variation of 30-80% for transporter expression, yet current models treat all cells as identical pharmacological units. This gap leaves unexplained a substantial

component of observed clinical variability. The paper addresses how cellular-level heterogeneity in drug uptake, mediated by transporter expression variability, propagates through nonlinear dose-response relationships to determine population-level outcomes. It builds on recent stochastic pharmacodynamic models but focuses specifically on the amplification mechanism arising from transporter heterogeneity.

✂ Methodological Snapshot

The authors developed a multiscale stochastic PK-PD framework. Systemic pharmacokinetics are modeled deterministically (assumed uniform plasma exposure within a patient). Cellular pharmacokinetics are modeled with stochastic differential equations where each cell has a log-normally distributed number of influx transporters, leading to heterogeneous intracellular drug concentrations. Pharmacodynamics follow a Hill function for cell death rate. The framework is implemented in Julia using Euler-Maruyama discretization. Five experiments were conducted: (1) amplification scaling law validation, (2) extinction time distribution characterization, (3) mean-field model comparison, (4) population size scaling, and (5) parameter sensitivity analysis.

☒ Structural Model Breakdown

The model has two hierarchical levels: (1) Systemic PK: Deterministic plasma concentration profile for cobimetinib under 21/7 regimen (not explicitly modeled but assumed known). (2) Cellular PK: Each cell i has a stochastic intracellular concentration $C_i(t)$ governed by $dC_i/dt = k_{in}T_iC_{plasma}(t) - k_{out}C_i(t)$, where T_i is the number of influx transporters (log-normally distributed across cells). (3) Cellular PD: Death rate for cell i is given by $k_{max}C_i^\gamma/(EC_{50}^\gamma + C_i^\gamma)$. Population extinction occurs when all cells have died. The amplification factor A is defined as the ratio of variance in death rate to variance in transporter expression, theoretically $A = \left(\frac{\gamma EC_{50}^\gamma C_{med}^{\gamma-1}}{(EC_{50}^\gamma + C_{med}^\gamma)^2} \right)^2 C_{med}^2$ for small heterogeneity.

Detailed Methodological Analysis

Modeling Approach

Multiscale stochastic PK-PD framework: deterministic systemic PK (not explicitly detailed, but assumed to provide plasma concentration-time profiles for the 21/7 dosing regimen of cobimetinib), coupled with cellular-level stochastic differential equations for intracellular drug concentration. Transporter expression per cell is modeled as log-normal with $CV = 53\%$ (baseline). Cell death rate follows a Hill function of intracellular concentration. Three model variants: Model H (homogeneous), Model PK (heterogeneous transporters but averaged death rate), Model PKPD (full heterogeneity). Implemented in Julia 1.10 with Euler-Maruyama discretization (time step 0.01 h).

Data Sources

All parameters derived from published literature and regulatory documents for cobimetinib without free fitting. IC_{50} values from literature (range 0.001-39.3 μ M across cell lines; baseline

2.0 μM). Hill coefficient set to 1 (competitive inhibition). Transporter expression CV from single-cell measurements (50-70%; baseline 53%). Dosing regimen: 21/7 schedule (21 days on, 7 days off) as per clinical protocol.

Estimation Methods

No parameter estimation; all parameters fixed from literature. The study is simulation-based, not estimation-based.

Model Evaluation

Model variants compared via extinction time distributions using Kolmogorov-Smirnov tests, F-tests for variance equality, and quantile spreads. Extended simulations (180 days) with 500 replicates. Mixture model comparison using Bayesian Information Criterion (BIC) for multimodal distributions. Sensitivity analysis across 72 parameter combinations.

Covariate Analysis

Systematic sensitivity analysis varying Hill coefficient (1-4), transporter CV (10-70%), dose (0.5-2.0 $\times$ clinical dose), and EC_{50} (0.5-4.0 μM). Transporter CV identified as the strongest predictor of treatment failure. Amplification factor characterized as function of these parameters.

Statistical Rigor Assessment

The study uses appropriate statistical tests for comparing distributions (KS test, F-test). The use of BIC for mixture model selection is standard. Sensitivity analysis covers a wide parameter space. However, no formal power analysis or confidence intervals are provided for the simulation results. The number of replicates (1000 for main experiments, 500 for extended) is adequate for characterizing distributions. The analysis is descriptive rather than inferential, which is appropriate for a proof-of-principle study.

✕ Key Findings

The study demonstrates that cellular transporter heterogeneity amplifies pharmacodynamic variance up to 1000-fold through nonlinear dose-response, creating cycle-structured extinction time distributions under intermittent dosing. Under the 21/7 regimen, 36% of heterogeneous populations failed to achieve extinction within 56 days, compared to 0% for homogeneous populations. The median extinction time increased from 261 h (homogeneous) to 875 h (heterogeneous), a 3.4-fold prolongation, and variance increased 22-fold. Extended 180-day simulations showed 4.8% of populations remained viable, with extinction times spanning 17.5 to 118.9 days. Mean-field models underpredicted median extinction times by 60-81%. Treatment failure increased with population size, while the amplification factor remained approximately constant. Transporter CV emerged as the strongest predictor of treatment failure across all parameter combinations.

🔗 Clinical & Regulatory Implications

The findings suggest that tumors with high transporter heterogeneity may exhibit lower complete response rates and wider progression-free survival distributions. The 36% treatment failure rate under adequate systemic exposure implies that therapeutic drug monitoring alone may not reduce variability. The cycle-structured extinction times under intermittent dosing suggest that response assessment timing matters. Combination therapies using drugs with uncorrelated transporter dependencies could reduce sanctuary cell formation. Single-cell measurements of transporter expression could stratify patients: low-heterogeneity tumors may have high complete response rates, while high-heterogeneity tumors may benefit from alternative dosing schedules or combinations.

Strengths & Limitations

Strengths

- Novel multiscale stochastic framework coupling deterministic systemic PK with cellular-level SDEs
- All parameters derived from literature without free fitting, ensuring mechanistic plausibility
- Systematic exploration of amplification mechanism through five computational experiments
- Clear demonstration of mean-field model inadequacy with quantitative error metrics
- Comprehensive sensitivity analysis across 72 parameter combinations
- Clinically relevant insights for dosing schedule design and combination therapy

Limitations (Acknowledged by Authors)

- Focus on influx transporter heterogeneity, excluding efflux transporters and metabolic enzymes
- Cells treated as independent, neglecting paracrine signaling, spatial structure, and microenvironment
- Parameters anchored to preclinical literature, not calibrated against matched experimental data
- Model does not include acquired resistance mechanisms
- Proof-of-principle framework not intended for quantitative clinical inference

Limitations (Expert Review)

- Systemic PK model not explicitly described; assumed deterministic but details lacking
- No formal model selection or validation against real clinical data
- Extrapolation to clinical tumor sizes (10^9 cells) is qualitative; finite-size scaling may not hold
- Death process modeled as deterministic after concentration threshold; stochastic cell death could add further variability
- No consideration of drug binding, intracellular trafficking, or target engagement dynamics

Generalizability

The findings are demonstrated for cobimetinib, a MEK inhibitor, but the mechanism is general for drugs with steep dose-response curves and transporter-mediated uptake. The qualitative conclusions likely apply to other targeted therapies, but quantitative predictions require drug-specific parameterization. The framework can be extended to other drugs and transporters.

Key Equations

Cellular Pharmacokinetic SDE

$$\frac{dC_i}{dt} = k_{in} T_i C_p(t) - k_{out} C_i$$

Stochastic differential equation for intracellular drug concentration in cell i , where T_i is the number of influx transporters and $C_p(t)$ is the plasma concentration.

Hill Function for Cell Death Rate

$$\mu_i = \mu_{max} \frac{C_i^\gamma}{EC_{50}^\gamma + C_i^\gamma}$$

Death rate of cell i as a function of its intracellular concentration C_i , with Hill coefficient γ and half-maximal effect concentration EC_{50} .

First-Order Amplification Factor

$$A = \left(\frac{\gamma EC_{50}^\gamma C_{med}^{\gamma-1}}{(EC_{50}^\gamma + C_{med}^\gamma)^2} \right)^2 C_{med}^2$$

Theoretical amplification factor relating variance in death rate to variance in transporter expression, derived from first-order Taylor expansion around median concentration C_{med} .

Log-Normal Transporter Distribution

$$T_i \sim \text{LogNormal}(\mu_T, \sigma_T^2), CV = \sqrt{e^{\sigma_T^2} - 1}$$

Distribution of transporter counts across cells, parameterized by mean and coefficient of variation (CV).

Figures & Tables

- **Figure 1:** Amplification factor scaling with Hill coefficient and transporter heterogeneity.
 - *Significance:* Validates the amplification mechanism and shows that amplification peaks at intermediate heterogeneity and saturates at high CV.
- **Figure 2:** Extinction time distributions across model variants (H, PK, PKPD).
 - *Significance:* Demonstrates the dramatic broadening of extinction times in the full heterogeneity model (PKPD) compared to homogeneous (H) and PK-only models.
- **Figure 3:** Extended 180-day simulation extinction time distributions.
 - *Significance:* Shows that a fraction of heterogeneous populations remain viable beyond 180 days, with multimodal structure aligned to dosing cycles.
- **Figure 4:** Mean-field model inadequacy: comparison of extinction times predicted by mean-field variants vs. stochastic simulations.

- *Significance*: Quantifies the systematic underprediction (60-81% error) of mean-field models, highlighting their fundamental inadequacy for heterogeneous systems.
 - **Table 1**: Quantile characterization of extinction times for Model H, PK, and PKPD.
 - *Significance*: Provides quantitative comparison of distributional broadening, showing that PKPD/H ratios range from 1.65 to 3.48 across quantiles.
-

Future Directions

The authors suggest several testable predictions: (1) In vitro treatment of heterogeneous cell populations should yield cycle-structured extinction time distributions. (2) Clonogenic survival assays should show increased failure rates with larger initial populations. (3) Pre-treatment transporter expression heterogeneity should predict response variability better than mean expression. (4) Continuous vs. intermittent dosing should produce different extinction time distributions. Future work should extend the framework to include efflux transporters, spatial drug gradients, proliferation, and adaptive resistance. Experimental validation in matched cell lines and animal models is needed before clinical translation.


Expert Commentary

This paper addresses a critical gap in pharmacometrics by explicitly modeling cellular heterogeneity in drug uptake. The amplification mechanism is elegantly demonstrated and has profound implications for how we interpret variability in clinical trials. The failure of mean-field models is particularly striking and should serve as a cautionary tale for modelers who rely on population-averaged approaches without considering underlying heterogeneity. The sensitivity analysis revealing that transporter CV is the key driver of outcome variability provides a clear target for experimental measurement. However, the proof-of-principle nature means that quantitative predictions should not be overinterpreted. The framework is a valuable addition to the stochastic pharmacodynamics toolkit and could be extended to other therapeutic areas such as antibiotics and antivirals where cellular uptake heterogeneity is relevant.

Bottom Line

This proof-of-principle study provides compelling evidence that cellular heterogeneity in drug transporter expression, when combined with nonlinear pharmacodynamics, can dramatically amplify outcome variability and produce treatment failure even with adequate systemic exposure. The work highlights the limitations of traditional mean-field PK-PD models and advocates for incorporating stochastic cell-level modeling into drug development. For practicing pharmacometricians, the key takeaway is that intra-tumor heterogeneity in drug uptake should be considered as a source of unexplained variability, and that single-cell measurements of transporter expression could improve patient stratification and treatment optimization.

☒ Figures



(A) Log-log scatter plot of A versus n^2 at $\sigma_T = 0.5$

(B) Heatmap of $\log_{10}$ -transformed amplification factor A

Figure 1: Amplification factor scaling with Hill coefficient and transporter heterogeneity.




Figure 2: Extinction time distributions across model variants.




Figure 3: Extended simulations extinction time distributions.




Figure 4: Mean-field model inadequacy.