

Rotator Cuff Injury

The rotator cuff is a group of muscles and tendons that surround the shoulder joint, keeping the head of your upper arm bone firmly within the shallow socket of the shoulder. A rotator cuff injury can cause a dull ache in the shoulder, which often worsens when you try to sleep on the involved side.

Rotator cuff injuries occur most often in people who repeatedly perform overhead motions in their jobs or sports. Examples include painters, carpenters, and people who play baseball or tennis. The risk of rotator cuff injury also increases with age.

Many people recover from rotator cuff disease with physical therapy exercises that improve flexibility and strength of the muscles surrounding the shoulder joint.

Sometimes, rotator cuff tears may occur as a result of a single injury. In those circumstances, medical care should be provided as soon as possible. Extensive rotator cuff tears may require surgical repair, transfer of alternative tendons or joint replacement.







Symptoms

- **The pain associated with a rotator cuff injury**
- **Be described as a dull ache deep in the shoulder**
- **Disturb sleep, particularly if you lie on the affected shoulder**
- **Make it difficult to comb your hair or reach behind your back**
- **Be accompanied by arm weakness**

Treatment

Conservative treatments — such as rest, ice and physical therapy — sometimes are all that's needed to recover from a rotator cuff injury. If your injury is severe and involves a complete tear of the muscle or tendon, you might need surgery.

Injections

If conservative treatments haven't reduced your pain, your doctor might recommend a steroid injection into your

shoulder joint, especially if the pain is interfering with your sleep, daily activities or exercise. While such shots are often temporarily helpful, they should be used judiciously, as they can contribute to the weakening of the tendon.

Physical Therapy

Physical therapy is usually one of the first treatments your doctor may suggest. Exercises tailored to the specific location of your rotator cuff injury can help restore flexibility and strength to your shoulder. Physical therapy is also an important part of the recovery process after rotator cuff surgery.

Surgery

Many different types of surgeries are available for rotator cuff injuries, including:

- Arthroscopic tendon repair. In this procedure, surgeons insert a tiny camera (arthroscope) and tools through small incisions to reattach the torn tendon to the bone.**
- Open tendon repair. In some situations, an open tendon repair may be a better option. In these types of surgeries, your surgeon works through a larger incision to reattach the damaged tendon to the bone. Compared to**

arthroscopic procedures, open tendon repairs typically heal in the same length of time but recovery may be more uncomfortable.

- **Tendon transfer. If the torn tendon is too damaged to be reattached to the arm bone, surgeons may decide to use a nearby tendon as a replacement.**
- **Shoulder replacement. Massive rotator cuff injuries may require shoulder replacement surgery. To improve the artificial joint's stability, an innovative procedure (reverse shoulder arthroplasty) installs the ball part of the artificial joint onto the shoulder blade and the socket part onto the arm bone**

إصابة الكفة المدورة

الكفة المدورة عبارة عن مجموعة من العضلات والأوتار التي تحيط بمفصل الكتف ، مما يحافظ على رأس عظم الذراع بقوة داخل التجويف الضحل للكتف. يمكن أن تسبب إصابة الكفة المدورة ألمًا خفيفًا في الكتف ، والذي غالبًا ما يزداد سوءًا عند محاولة النوم على الجانب المصاب

تحدث إصابات الكفة المدورة في أغلب الأحيان عند الأشخاص الذين يؤدون حركات علوية بشكل متكرر في وظائفهم أو ممارسة الرياضة. تشمل الأمثلة الرسامين والنجارين والأشخاص الذين يلعبون البيسبول أو التنس. يزداد خطر إصابة الكفة المدورة أيضًا مع تقدم العمر

يتعافى العديد من الأشخاص من مرض الكفة المدورة من خلال تمارين العلاج الطبيعي التي تعمل على تحسين مرونة وقوة العضلات المحيطة بمفصل الكتف

في بعض الأحيان ، قد تحدث تمزقات الكفة المدورة نتيجة لإصابة واحدة. في مثل هذه الظروف ، يجب تقديم الرعاية الطبية في أسرع وقت ممكن. قد تتطلب تمزقات الكفة المدورة الواسعة إصلاحًا جراحيًا أو نقل أوتار بديلة أو استبدال مفصل







الأعراض المعروفة

- الألم المصاحب لإصابة الكفة المدورة •
- أن يوصف بأنه ألم خفيف عميق في الكتف •
- اضطراب النوم ، خاصة إذا كنت مستلقيًا على الكتف المصاب •
- يصعب تمشييط شعرك أو الوصول إلى خلف ظهرك •
- أن يكون مصحوبا بضعف في الذراع •

ما هي طرق العلاج؟

العلاجات التحفظية - مثل الراحة والتلج والعلاج الطبيعي - تكون أحياناً كل ما يلزم للتعافي من إصابة الكفة المدورة. إذا كانت إصابتك شديدة وتنطوي على تمزق كامل للعضلة أو الوتر ، فقد تحتاج إلى جراحة.

الحقن

إذا لم تخفف العلاجات التحفظية من الألم ، فقد يوصي طبيبك بحقن الستيرويد في مفصل الكتف ، خاصةً إذا كان الألم يتعارض مع نومك أو أنشطتك اليومية أو ممارسة الرياضة. في حين أن مثل هذه الحقن غالباً ما تكون مفيدة بشكل مؤقت ، إلا أنه يجب استخدامها بحكمة ، لأنها يمكن أن تسهم في إضعاف الوتر.

العلاج الطبيعي

عادةً ما يكون العلاج الطبيعي أحد العلاجات الأولى التي قد يقترحها طبيبك. يمكن أن تساعد التمارين المصممة وفقًا للموقع المحدد لإصابة الكفة المدورة في استعادة المرونة والقوة لكتفك. يعد العلاج الطبيعي أيضًا جزءًا مهمًا من عملية التعافي بعد جراحة الكفة المدورة.

• عملية جراحية يتم فيها إزالة الجزء التالف من المريء وتوصيل الجزء المتبقي بمعدتك.

تتوفر أنواع مختلفة من العمليات الجراحية لإصابات الكفة المدورة ، بما في ذلك

- في هذا الإجراء ، يدخل الجراحون كاميرا صغيرة (منظار المفصل) .إصلاح الأوتار بالمنظار . وأدوات من خلال شقوق صغيرة لإعادة وصل الوتر الممزق بالعظم.
- في بعض الحالات ، قد يكون إصلاح الوتر المفتوح خيارًا أفضل. في إصلاح الوتر المفتوح . هذه الأنواع من العمليات الجراحية ، يعمل الجراح من خلال شق أكبر لإعادة ربط الوتر التالف بالعظام. بالمقارنة مع إجراءات التنظير المفصلي ، عادةً ما تلتئم إصلاحات الأوتار المفتوحة في نفس المدة الزمنية ولكن التعافي قد يكون أكثر إزعاجًا.
- إذا كان الوتر الممزق تالفًا للغاية بحيث لا يمكن إعادة توصيله بعظم الذراع ، .نقل الأوتار . فقد يقرر الجراحون استخدام وتر قريب كبديل.
- قد تتطلب إصابات الكفة المدورة الضخمة جراحة استبدال الكتف. لتحسين . استبدال الكتف . ثبات المفصل الاصطناعي ، يقوم إجراء مبتكر (تقويم مفصل الكتف العكسي) بتثبيت الجزء الكروي من المفصل الاصطناعي على لوح الكتف وجزء التجويف على عظم الذراع