

Ovarian Cyst


Ovarian cysts are fluid-filled sacs or pockets in an ovary or on its surface. Women have two ovaries — each about the size and shape of an almond — on each side of the uterus. Eggs (ova), which develop and mature in the ovaries, are released in monthly cycles during the childbearing years.





Many women have ovarian cysts at some time. Most ovarian cysts present little or no discomfort and are harmless. The majority disappears without treatment within a few months.

However, ovarian cysts — especially those that have ruptured — can cause serious symptoms. To protect your health, get regular pelvic exams and know the symptoms that can signal a potentially serious problem.



Symptoms

Most cysts don't cause symptoms and go away on their own. However, a large ovarian cyst can cause:

- **Pelvic pain — a dull or sharp ache in the lower abdomen on the side of the cyst**
- **Fullness or heaviness in your abdomen**
- **Bloating**



When to see a doctor?

Seek immediate medical attention if you have:

- **Sudden, severe abdominal or pelvic pain**
- **Pain with fever or vomiting**

If you have these signs and symptoms or those of shock — cold, clammy skin; rapid breathing; and lightheadedness or weakness — see a doctor right away.

Causes

Most ovarian cysts develop as a result of your menstrual cycle (functional cysts). Other types of cysts are much less common.

Functional cysts

Your ovaries normally grow cyst-like structures called follicles each month. Follicles produce the hormones estrogen and progesterone and release an egg when you ovulate.

If a normal monthly follicle keeps growing, it's known as a functional cyst. There are two types of functional cysts:

- Follicular cyst. Around the midpoint of your menstrual cycle, an egg bursts out of its follicle and travels down the fallopian tube. A follicular cyst begins when the follicle doesn't rupture or release its egg, but continues to grow.**
- Corpus luteum cyst. When a follicle releases its egg, it begins producing estrogen and progesterone for conception. This follicle is now called the corpus luteum. Sometimes, fluid accumulates inside the follicle, causing the corpus luteum to grow into a cyst.**
- Functional cysts are usually harmless, rarely cause pain, and often disappear on their own within two or three menstrual cycles.**

Other cysts

Types of cysts not related to the normal function of your menstrual cycle include:

- **Dermoid cysts.** Also called teratomas, these can contain tissue, such as hair, skin or teeth, because they form from embryonic cells. They're rarely cancerous.
- **Cystadenomas.** These develop on the surface of an ovary and might be filled with a watery or a mucous material.
- **Endometriomas.** These develop as a result of a condition in which uterine endometrial cells grow outside your uterus (endometriosis). Some of the tissue can attach to your ovary and form a growth.

Risk factors

Your risk of developing an ovarian cyst is heightened by:

- **Hormonal problems.** These include taking the fertility drug clomiphene (Clomid), which is used to cause you to ovulate.
- **Pregnancy.** Sometimes, the cyst that forms when you ovulate stays on your ovary throughout your pregnancy.
- **Endometriosis.** This condition causes uterine endometrial cells to grow outside your uterus. Some of the tissue can attach to your ovary and form a growth.

- **A severe pelvic infection.** If the infection spreads to the ovaries, it can cause cysts.
- **A previous ovarian cyst.** If you've had one, you're likely to develop more.

Complications

Some women develop fewer common types of cysts that a doctor finds during a pelvic exam. Cystic ovarian masses that develop after menopause might be cancerous (malignant). That's why it's important to have regular pelvic exams.

Infrequent complications associated with ovarian cysts include:

- **Ovarian torsion.** Cysts that enlarge can cause the ovary to move, increasing the chance of painful twisting of your ovary (ovarian torsion). Symptoms can include an abrupt onset of severe pelvic pain, nausea and vomiting. Ovarian torsion can also decrease or stop blood flow to the ovaries.
- **Rupture.** A cyst that ruptures can cause severe pain and internal bleeding. The larger the cyst, the greater the risk of rupture. Vigorous activity that affects the pelvis, such as vaginal intercourse, also increases the risk.

Prevention

Although there's no way to prevent ovarian cysts, regular pelvic examinations help ensure that changes in your ovaries are diagnosed as early as possible. Be alert to changes in your monthly cycle, including unusual menstrual symptoms, especially ones that persist for more than a few cycles. Talk to your doctor about changes that concern you.

Diagnosis

A cyst on your ovary can be found during a pelvic exam. Depending on its size and whether it's fluid filled, solid or mixed, your doctor likely will recommend tests to determine its type and whether you need treatment. Possible tests include:

- **Pregnancy test.** A positive test might suggest that you have a corpus luteum cyst.
- **Pelvic ultrasound.** A wandlike device (transducer) sends and receives high-frequency sound waves (ultrasound) to create an image of your uterus and ovaries on a video screen. Your doctor analyzes the image to confirm the presence of a cyst, help identify its location and determine whether it's solid, filled with fluid or mixed.
- **Laparoscopy.** Using a laparoscope — a slim, lighted instrument inserted into your abdomen through a small

incision — your doctor can see your ovaries and remove the ovarian cyst. This is a surgical procedure that requires anesthesia.

- **CA 125 blood test. Blood levels of a protein called cancer antigen 125 (CA 125) often are elevated in women with ovarian cancer. If your cyst is partially solid and you're at high risk of ovarian cancer, your doctor might order this test.**

Elevated CA 125 levels can also occur in noncancerous conditions, such as endometriosis, uterine fibroids and pelvic inflammatory disease.

Treatment

Treatment depends on your age, the type and size of your cyst, and your symptoms. Your doctor might suggest:

- **Watchful waiting. In many cases you can wait and be re-examined to see if the cyst goes away within a few months. This is typically an option — regardless of your age — if you have no symptoms and an ultrasound shows you have a simple, small, fluid-filled cyst.**

Your doctor will likely recommend that you get follow-up pelvic ultrasounds at intervals to see if your cyst changes in size.


- **Medication.** Your doctor might recommend hormonal contraceptives, such as birth control pills, to keep ovarian cysts from recurring. However, birth control pills won't shrink an existing cyst.
- **Surgery.** Your doctor might suggest removing a cyst that is large, doesn't look like a functional cyst, is growing, continues through two or three menstrual cycles, or causes pain.

Some cysts can be removed without removing the ovary (ovarian cystectomy). In some cases, your doctor might suggest removing the affected ovary and leaving the other intact (oophorectomy).

If a cystic mass is cancerous, your doctor will likely refer you to a gynecologic cancer specialist. You might need to have your uterus, ovaries and fallopian tubes removed (total hysterectomy) and possibly chemotherapy or radiation. Your doctor is also likely to recommend surgery when an ovarian cyst develops after menopause.


كيس المبيض

أكياس المبيض هي أكياس أو جيوب مملوءة بالسوائل في المبيض أو على سطحه. لدى النساء مبيضان - كل منهما بحجم وشكل اللوز - على كل جانب من الرحم. البويضات ، التي تنمو وتنضج في المبايض ، يتم إطلاقها في دورات شهرية خلال سنوات الإنجاب.



تعاين العديد من النساء من تكيسات المبيض في وقت ما. معظم تكيسات المبيض تسبب القليل من الانزعاج أو لا تسبب أي إزعاج وهي غير ضارة. تختفي الغالبية دون علاج في غضون بضعة أشهر.

ومع ذلك ، يمكن أن تسبب تكيسات المبيض - خاصة تلك التي تمزق - أعراضاً خطيرة. لحماية صحتك ، قم بإجراء فحوصات الحوض بانتظام وتعرف على الأعراض التي يمكن أن تشير إلى مشكلة خطيرة محتملة.



الأعراض المعروفة

لا تسبب معظم التكيّسات أعراضاً وتختفي من تلقاء نفسها. ومع ذلك ، يمكن أن يتسبب كيس المبيض الكبير في:

- ألم الحوض - ألم خفيف أو حاد في أسفل البطن على جانب الكيس .

- امتلاء أو ثقل في بطنك
- الانتفاخ •



متى ترى الطبيب؟

اطلب عناية طبية فورية إذا كان لديك

- ألم مفاجئ وشديد في البطن أو الحوض
- ألم مصحوب بحمى أو قيء

إذا كانت لديك هذه العلامات والأعراض أو أعراض الصدمة - جلد بارد ورطب ؛ تنفس سريع؛ والدوار
أو الضعف - راجع الطبيب على الفور

الأسباب وراء سرطان المريء

تتطور معظم تكيسات المبيض كنتيجة لدورة الطمث (الخراجات الوظيفية). الأنواع الأخرى من
الخراجات أقل شيوعاً.

الخراجات الوظيفية

عادةً ما ينمو مبيضك هياكل شبيهة بالأكياس تسمى الجُريبات كل شهر. تنتج الجريبات هرموني
الإستروجين والبروجسترون وتطلق البويضة عند الإباضة.

إذا استمرت الجريبات الشهرية الطبيعية في النمو ، فإنها تُعرف باسم كيس وظيفي. هناك نوعان من التكيسات الوظيفية:

- في منتصف الدورة الشهرية ، تخرج البويضة من جريبها وتنتقل عبر قناة .كيس مسامي فالوب. يبدأ الكيس الجريبي عندما لا يتمزق الجريب أو يطلق البويضة ، ولكنه يستمر في النمو.
- عندما يطلق الجريب بيضته ، يبدأ في إنتاج هرمون الاستروجين .كيس الجسم الأصفر والبروجسترون للحمل. يسمى هذا الجريب الآن الجسم الأصفر. في بعض الأحيان ، يتراكم السائل داخل الجريب ، مما يتسبب في نمو الجسم الأصفر إلى كيس.
- عادة ما تكون الأكياس الوظيفية غير ضارة ونادرًا ما تسبب الألم وغالبًا ما تختفي من تلقاء نفسها في غضون دورتين أو ثلاث دورات شهرية.

الخراجات الأخرى

تشمل أنواع التكيسات غير المرتبطة بالوظيفة الطبيعية لدورة الطمث ما يلي:

- تسمى أيضًا التراتومة ، ويمكن أن تحتوي على أنسجة ، مثل الشعر أو الجلد .كيسات الجلد .أو الأسنان ، لأنها تتكون من خلايا جنينية. نادرا ما تكون سرطانية.
- تتطور هذه على سطح المبيض وقد تمتلئ بمادة مائية أو مخاطية .أورام المثانة .
- تتطور هذه نتيجة حالة تنمو فيها خلايا بطانة الرحم خارج الرحم (الانتباز .أورام بطانة الرحم .البطاني الرحمي). يمكن أن تلتصق بعض الأنسجة بالمبيض وتشكل نموًا

عوامل الخطر

يزداد خطر إصابتك بكيس المبيض من خلال

- وتشمل هذه تناول عقار كلوميفين (كلوميد) للخصوبة ، والذي يستخدم .مشاكل هرمونية .لأحداث التبويض
- في بعض الأحيان ، يبقى الكيس الذي يتشكل عند التبويض على مبيضك طوال فترة .حمل .الحمل
- تتسبب هذه الحالة في نمو خلايا بطانة الرحم خارج الرحم. يمكن أن تلتصق بطانة الرحم • .بعض الأنسجة بالمبيض وتشكل نموًا
- .إذا انتشرت العدوى إلى المبايض ، فقد تتسبب في حدوث تكيسات .عدوى شديدة في الحوض
- .إذا كان لديك واحد ، فمن المحتمل أن تتطور أكثر .كيس مبيض سابق

المضاعفات

تظهر لدى بعض النساء عدد أقل من أنواع التكيسات الشائعة التي يكتشفها الطبيب أثناء فحص الحوض. قد تكون كتل المبيض الكيسية التي تظهر بعد انقطاع الطمث سرطانية (خبيثة). لهذا السبب من المهم إجراء فحوصات الحوض بانتظام

:تشمل المضاعفات النادرة المرتبطة بكيسات المبيض ما يلي

- يمكن أن تتسبب الأكياس التي تتضخم في تحرك المبيض ، مما يزيد من .التواء المبيض فرصة التواء المبيض المؤلم (التواء المبيض). يمكن أن تشمل الأعراض الظهور المفاجئ لآلام الحوض الشديدة والغثيان والقيء. يمكن أن يؤدي التواء المبيض أيضًا إلى انخفاض أو إيقاف تدفق الدم إلى المبايض
- يمكن أن يسبب الكيس الذي يتمزق ألمًا شديدًا ونزيفًا داخليًا. كلما زاد حجم الكيس ، .تمزق زاد خطر التمزق. كما أن النشاط القوي الذي يؤثر على الحوض ، مثل الجماع المهبلي ، يزيد أيضًا من المخاطر

هل يوجد طرق للوقاية؟

على الرغم من عدم وجود طريقة للوقاية من تكيسات المبيض ، فإن فحوصات الحوض المنتظمة تساعد في ضمان تشخيص التغيرات في المبايض في أقرب وقت ممكن. كوني متيقظة للتغيرات في دورتك الشهرية ، بما في ذلك أعراض الدورة الشهرية غير العادية ، خاصة تلك التي تستمر لأكثر من بضع دورات. تحدث إلى طبيبك حول التغيرات التي تقلقك.

تشخيص

يمكن العثور على كيس على مبيضك أثناء فحص الحوض. اعتمادًا على حجمه وما إذا كان ممتلئًا بالسوائل أو صلبًا أو مختلطًا ، فمن المرجح أن يوصي طبيبك بإجراء اختبارات لتحديد نوعه وما إذا كنت بحاجة إلى العلاج. تشمل الاختبارات المحتملة ما يلي:

- قد يشير الاختبار الإيجابي إلى إصابتك بكيس الجسم الأصفر. اختبار الحمل.
- جهاز يشبه العصا (محول الطاقة) يرسل ويستقبل موجات. الحوض بالموجات فوق الصوتية صوتية عالية التردد (الموجات فوق الصوتية) لإنشاء صورة للرحم والمبيضين على شاشة فيديو. يقوم طبيبك بتحليل الصورة للتأكد من وجود كيس ، ويساعد في تحديد موقعه وتحديد ما إذا كان صلبًا أو مملئًا بالسوائل أو مختلطًا.
- باستخدام منظار البطن - وهو أداة رفيعة ومضاعة يتم إدخالها في البطن من منظار البطن • خلال شق صغير - يمكن لطبيبك رؤية المبيضين وإزالة كيس المبيض. هذا إجراء جراحي يتطلب التخدير.
- **CA 125** غالبًا ما ترتفع مستويات بروتين يسمى مستضد السرطان **CA 125** فحص الدم في الدم لدى النساء المصابات بسرطان المبيض. إذا كان التكيس صلبًا جزئيًا وكنت **125** أكثر عرضة للإصابة بسرطان المبيض ، فقد يطلب طبيبك هذا الاختبار.

المرتفعة أيضًا في الحالات غير السرطانية ، مثل الانتباذ البطاني **CA 125** يمكن أن تحدث مستويات الرحمي والأورام الليفية الرحمية ومرض التهاب الحوض.

ما هي طرق العلاج؟

يعتمد العلاج على عمرك ونوع وحجم التكييس والأعراض. قد يقترح طبيبك ما يلي:

- في كثير من الحالات ، يمكنك الانتظار وإعادة الفحص لمعرفة ما إذا كان .الانتظار اليقظ • . الكيس قد اختفى في غضون بضعة أشهر. عادةً ما يكون هذا خيارًا - بغض النظر عن عمرك - إذا لم تكن لديك أعراض وأظهر الفحص بالموجات فوق الصوتية أن لديك كيسًا بسيطًا وصغيرًا مليئًا بالسوائل.

من المحتمل أن يوصي طبيبك بإجراء متابعة بالموجات فوق الصوتية للحوض على فترات لمعرفة ما إذا كان حجم التكييس لديك يتغير.

- قد يوصي طبيبك بوسائل منع الحمل الهرمونية ، مثل حبوب منع الحمل ، لمنع .دواء • . تكيسات المبيض من التكرار. ومع ذلك ، لن تقلص حبوب منع الحمل التكيسات الموجودة.
- قد يقترح طبيبك إزالة كيس كبير ، لا يبدو ككيس وظيفي ، ينمو ، يستمر خلال .الجراحة • . دورتين أو ثلاث دورات حيض ، أو يسبب الألم.

يمكن إزالة بعض الأكياس دون إزالة المبيض (استئصال المئانة المبيض). في بعض الحالات ، قد يقترح طبيبك إزالة المبيض المصاب وترك الآخر سليمًا (استئصال المبيض).

إذا كانت الكتلة الكيسية سرطانية ، فمن المحتمل أن يحيلك طبيبك إلى أخصائي سرطان الجهاز التناسلي للمرأة. قد تحتاجين إلى إزالة الرحم والمبيضين وقتاتي فالوب (استئصال الرحم الكلي) وربما العلاج الكيميائي أو الإشعاعي. من المرجح أيضًا أن يوصي طبيبك بإجراء عملية جراحية عند ظهور كيس مبيض بعد انقطاع الطمث.