1. The probability of a leap year selected at random contain 53						
Sunday is:						
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365			
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at						
random. The prol	bability of draw	ving a black ball	is:			
• •	` '	(c) 0/5				
3. The probabilit	y that it will rai	in tomorrow is 0	.85. What is the			
probability that it						
* *			(d) none of these			
•	•		ed from the numbers			
(1, 2, 3,,15	•		4.0			
	* *	(c) 2/15	• •			
5. What are the						
		(c) 8				
-	-	number selecte	d at random from the			
numbers (1,2,3, .		() 10/0-	(1)			
• •		• '	(d) none of these			
7. The sum of th						
		0 (d) none				
_	_	are given; choos	e the correct answer			
for that which is			(1)			
			(d) none of these.			
		ultaneously, tha	n the probability of			
getting at least ty		/-\ 1/	(-1) 1 (0			
		(c) ½				
10. A letter is ch						
*A55A55INA I II	JN. I ne proi	pability that the	letter chosen has:			
(a) 6/13	(D) // 13	(C) I	(d) none of these.			
11 A diag is thre	wn Find the nu	cobobility of gott	ing on oven number			
	-	· · · · · · · · · · · · · · · · · · ·	ing an even number.			
(A) 2/3	(D) I	(C) 5/6	D) 1/2			
12. Two coins are thrown at the same time. Find the probability of getting both heads.						
(A) 3/4 (B) 1/4		(D) 0				
13. Two dice are thrown simultaneously. The probability of getting a						

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	! /9			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4	(B) 27/50	(C) 1/4	(D) 29/100			
of drawing blue balls in	a blue ball is do n a bag is:	ouble that of a	red ball, th	If the probability en the number of			
(A) 5	(B) 10	C) 15	(D) 20				
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:							
(A) 143/150	(B) 147/	(C)	1/25	(D) 1/50			
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100							
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366							
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$							

same birthd	ay is the same b		both will have the (D) 1/133225			
(A) 304/303	(D) 31/30	(C) 1/303	(D) 1/ 133223			
2. Then the	per x is chosen at probability that x B) 2/5 (C) 3/	x²< 2 is?	numbers -2, -1, 0 , 1,			
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21			
	<u>-</u>	n in 5 one day mate viation is	ches are 50, 70, 82,			
a) 25 70						
a) 20.79	b) 25.49	c) 25.29	d) 25.69			
28. Find me	dian and mode o	of the messages re	ceived on 9			
28. Find me	dian and mode of days 15, 11, 9,	,	ceived on 9			
28. Find me consecutive a) 13, 15	dian and mode of days 15, 11, 9, 5 b) 13, 18 s tossed up 4 tin	of the messages re 5, 18, 4, 18, 13, 17. c) 18, 15	ceived on 9			
28. Find me consecutive a) 13, 15 29. A coin is 3 cases isa) 1/2	dian and mode of days 15, 11, 9, 5 b) 13, 18 s tossed up 4 tin b) $\frac{1}{3}$	of the messages re 5, 18, 4, 18, 13, 17. c) 18, 15 mes. The probabilit	ceived on 9 d) 13, 16 by that tails turn up in d) $\frac{1}{6}$			
28. Find me consecutive a) 13, 15 29. A coin is 3 cases isa) 1/2	dian and mode of days 15, 11, 9, b) 13, 18 s tossed up 4 tin b) $\frac{1}{3}$ ariate between 0	of the messages re 5, 18, 4, 18, 13, 17. c) 18, 15 mes. The probabilit c) $^1/_4$ and 3. The value o	ceived on 9 d) 13, 16 by that tails turn up in d) $\frac{1}{6}$			

respectively. Let Z= 5X-2Y. The variance of Z is?

32.Out of to probability?	•	alues, which	one is not poss	sible in
	b) ∑ x d) P(P(x) = 3 x) = -0.5		
33.If E(x) =	2 and E(z) = 4 b) 6	4, then E(z - c) 0		sufficient data
34.The cov	ariance of two	independer	nt random varial	ole is
a) 1	b) 0	c) - 1	d) Ur	ndefined
35.If Σ P(x) a) 0) = k² - 8 then b) 1	, the value o <mark>c) 3</mark>		sufficient data
	0.5 and x = 4, b) 0.5		? d) 2	
37.In a disc is always?	rete probabili	ty distributio	on, the sum of al	l probabilities
a) 0	b) Infinite	c) 1	d) Un	defined
38.If the pr	obability of hi	tting the tar	get is 0.4, find n	nean and
	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16
-	% and if 10 bo	mbs are dro	pped from a place opped, find mear 0.4, 0.16	
a) 2	e mean of toss b) 4 s the mean and	c) 8	d) 1 or standard norn	nal distribution?

c) 5

d) 7

a) 3

b) 4

				•		l variance nd varianc		
42. a) E		e of a rand b) E()	lom varia (2)		•		. d) (E(X))2	
	43.Mean of a random variable X is given by a) E(X)							
	44.Mean of a constant 'a' is a) 0 b) a c) a/2 d) 1							
45.V a) 0	ariance/	of a cons b) a	stant 'a' is	c) a/2		d) 1		
46.Find the mean and variance of X?								
г 	Х	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		
	a) 2, 4/3 b) 3, 4/3 c) 2, 2/3 d) 3, 2/3							

47. Find the expectation of a random variable X?

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2