The Psion Hypothesis

A Framework for the Origin of Matter, Antimatter, Dark Matter, and Gravity

October 2026

PH_Foundations v86

Karlos Marden Maia

Independent Researcher, Marietta, Georgia

karlosmardenmaia@gmail.com

[T1] Exact algebraic theorem or exact Gel'fand–Yaglom functional-determinant calculation. No numerical verification required.

[T2] Structurally derived and numerically verified. Mechanism identified; derivation complete at leading order.

[T2+] Structurally closed: mechanism identified, scheme-equivalence complete. One explicit algebraic step remains for T1 closure.

[T3] Pipeline defined, calculation open. Strategy identified; numerical result not yet obtained.

[T4] Conjectural: mechanism identified but not derived. Dimensional estimates only.

[SC] Self-consistency input: the empirical anchor. Two exist: GN (dimensional; mass–gravity–cosmology) and $\Delta m^2{}_{21}/m_e^2$ (adimensional; neutrino-sector scale, §11).

T1 and T2 results are the quantitative backbone of the PH.

All numerical verifications and derivation scripts: psion-proofs [2].

MSC: 81V05, 81V17, 83C65 | Keywords: psion, Z6 symmetry, composite spectrum, cosmological constant, baryogenesis, dark matter, Newton's constant

Abstract

The Psion Hypothesis (PH) derives the Standard Model's structural constants and the mass–gravity–cosmology observables (CKM/PMNS mixing and individual quark masses are not derived) from a single fermionic field ψ — the psion — with Z6 = Z3 × Z2 internal symmetry and spin j = 5/2. The framework has one dimensional empirical anchor in the mass–gravity–cosmology sector: Newton's constant GN (CODATA 2018: G = 6.67430 × 10^{-11} m^3 kg^{-1} s^{-2}, i.e. GN = 6.70883 × 10^{-39} GeV^{-2} in natural units). This is the uniquely appropriate anchor: the causal hierarchy

$$L(Z6, g^2, j) \to \text{condensate} \text{—Sakharov}\to \text{GN [SC]} \to \Lambda\text{meta} \to \text{me}$$

places GN two causal levels above me; anchoring to me would be causally circular. From the single dimensional SC input GN, the condensation scale follows: $\Lambda\text{meta} = \sqrt{(K(\xi)/GN)} \cdot \exp(-S/2g^2) = 1.2609 \times 10^{12}$ GeV ([T2]), from which the electron mass is a genuine cascade output:

$$\text{me} = \sqrt{(K(\xi)/GN)} \cdot \exp(-S/2g^2 - Ne/be) = 0.5089 \text{ MeV (0.42\% error)}.$$

Here Ne = 10 and be = 325/1152 are exact theorems of Z6 [T1], and $K(\xi) = 1.16335$ is the Gel'fand–Yaglom prefactor [T1]. This result is falsifiable: $\delta\text{me/me} = -\frac{1}{2}\,\delta\text{GN/GN}$ (prediction F2, testable via variation-of-constants cosmology).

A single condensate action S = 6Ne/be = 212.68 simultaneously yields: $\rho\Lambda = 2.492 \times 10^{-47}$ GeV^4 (0.93%, [T2+]); GN recovered at 0.83% self-consistency ([T2]); and the master invariant $\rho\Lambda GN^2 = 1.121 \times 10^{-123}$ predicted vs 1.132×10^{-123} observed (0.93%, [T2+] — Λmeta cancels algebraically and the prediction is exactly GN-independent). The invariant admits the closed structural form $[Nc/(27\pi^4)]\cdot K(\xi)^2\cdot e\text{^}(-S(1+3/\pi^2))$ with S = Nc·Ne/be, and is linear in $\hbar$ — the unique dimensionless completion is $\rho\Lambda GN^2\hbar/c^7$ — so that at fixed classical scales $\hbar \to 0$ sends $\rho\Lambda \to 0$: the accelerated expansion is a first-order quantum effect, with the falsifiable co-drift law $\delta\rho\Lambda/\rho\Lambda = -2\,\delta\text{GN/GN}$ (F17, §5.1).

Exact theorems establish: $N\nu = 3$ active neutrino species; Ng = 8 gluons; Ngen = 3 charged-lepton generations (the fourth is kinematically forbidden within Z6); and the absolute stability of 39 dark-matter states at body numbers $N \in \{9, 12, 15, 18, 21, 27\}$, with termination proved to N = 48. Further structural theorems include the unique-electron theorem (exactly one observable $Q = -1$ three-body state), the charge–chirality identity Q = nR − 1, and Theorem DM-S (dark composites are fermions at odd body number; the dark sector is Pauli-structured — forced-singlet s-wave states at N = 9, 12 and psion nuclei from N = 15 — while the observable sector is exactly the Pauli-trivial sector). The emergent-gravity sector passes its consistency gauntlet: the condensation channel is a Lorentz scalar (local Lorentz invariance at mean-field grade; c_gw = c_γ by structure), and the Z6 domain-wall problem resolves by infrared wall tension ($\sigma^{1/3} = 0.42$ MeV, 46 orders below the naive Λmeta^3) with complete chiral-bias annihilation in exact class ratios 5 : 8 : 9. Psion number is exactly conserved, so B + L is fixed and only B − L can change, the reverse of the Standard Model; the earlier derivation of $\eta B = 6.02 \times 10^{-10}$ is withdrawn (§13.3), and level bifurcation during the formation of composites after reheating is identified as a candidate mechanism, conditional on a conserved baryon label (P14-B, §13.7). Neff = 3.054 is retained with a status note (§13.5). Two new forces are predicted: the proto-colour force (X17 singlet, 3.59%, [T3]) and the Chiral-Flip force (CF-CC: structure [T1], rate bounded by $|O|^2 \leq 0.370$). Seventeen falsifiable predictions span four timescales; hard T1 falsifiers include any detection of a fourth charged lepton, neutrino, or gluon. The framework has two empirical (SC) inputs, GN and Δm^2_{21} (the neutrino-sector scale, §11); open problem P2/P12 ([T2]) seeks to close GN from 1 to 0 by solving the fixed-point condition $\Phi(\Lambda) = \Lambda$ of the two-loop NJL kernel, while Δm^2_{21} has a separate, unpromoted T4 closure candidate (P26-C, §11). This paper is the third of a four-act programme: from the Keplerian susceptibility G′r = Gp, through the phenomenological Relational Gravitational Dynamics (RGD) kernel, to the present Lagrangian, and finally to the derivation of the RGD kernel from this Lagrangian and

its resolution of the dark-matter/MOND controversy [12, 13].

Contents

1 The Primordial Field: The Psion

2 The Lagrangian and Cascade Action

3 The Cosmological Constant

4 Three Predictions from One Action

5 Master Invariant

6 Composite Spectrum from Z6

7 Physical Interpretation

8 Kinematic Stability: Theorem KS

9 Falsifiable Predictions — Complete Registry

10 Lepton Mass Spectrum

11 Neutrino Masses

12 Cosmic Accountability

13 Baryogenesis: Status after the P14 Audit

14 Dark Matter: Identity, Mass, and Relic Density

15 Open Problems

16 The Psion in Hawking Radiation

17 Exotic Signatures and Experimental Context

18 Discussion

19 Conclusions

A Complete Inventory of PH Results

B Derivation of the Kink Fluctuation Operator

C Exact Identities Used in Theorem 3.1

D Theorem P16: Renormalisation of the Z6 Wall Determinant

E Unit Conversion: $\rho\Lambda$ in GeV^4

F Principal Conclusions and Spectrum Analysis

G PH Proofs Repository

H The Psion Diagram: Complete Calculus for Composite Classification

I The Proto-Quark: PH Hadronic Constituents and Their Relation to SM Quarks

J The Chirality Asymmetry $\varepsilon\psi = 25/169$: Results

K Gravitational Sector: Condensate Response

L Method Audit Record: Eliminated Routes and Negative Findings

References

1 The Primordial Field: The Psion

One fermionic field ψ — a field whose quanta carry half-integer spin — with Z6 = Z3 × Z2 internal symmetry generates all observed particles as composites. The cyclic group Z6 of order 6 decomposes as the product of Z3 (proto-colour sector) and Z2 (chirality). The quantum of this field is the psion.

Each psion state k ∈ {0, . . . , 5} carries three labels: (a) proto-colour — a Z3 label {red, green, blue}, distinct from QCD colour; (b) chirality — L (left-handed, k ≤ 2) or R (right-handed, k ≥ 3); (c) magnetic quantum number mJ(k) = k − 5/2 ∈ {−5/2, −3/2, −1/2, +1/2, +3/2, +5/2}.

The electric charge is Qk = (4/9) Re($\omega^{k \bmod 3}$) − (1/2)(−1)^⌊k/3⌋ + 1/6, $\omega = e^{2\pi i/3}$.

k	Proto-colour	Chirality	mJ(k)	Qk
0	red	L	−5/2	+1/9
1	green	L	−3/2	−5/9
2	blue	L	−1/2	−5/9
3	red	R	+1/2	+10/9
4	green	R	+3/2	+4/9
5	blue	R	+5/2	+4/9

Table 1: The six psion states.

Here SU(2)meta is a global symmetry group of the psion Lagrangian, unrelated to the electroweak SU(2) of the Standard Model. In the charge formula, the three terms encode the Z3 proto-colour contribution, the Z2 chirality contribution, and the universal offset fixing ΣQk = 1 (Theorem V.1, App. J). The six states span the spin-5/2 representation of SU(2)meta: $J^2 = j(j + 1) = (5/2)(7/2) = 35/4$.

2 The Lagrangian and Cascade Action

2.1 Lagrangian

$$L = \bar{\psi}\, i\gamma^{\mu}\partial_{\mu}\psi - GUV(\bar{\psi}\psi)^2 - (\bar{\psi}J^{a}\psi)(\bar{\psi}\Sigma^{a}\psi) \qquad (1)$$

The first term is the standard Dirac kinetic term; the second is a contact (four-fermion) interaction with UV coupling GUV controlling the self-interaction strength at high energies; Ja and Σa are the SU(2)meta generators acting on the left and right sectors. The relative coefficient −1 is fixed by Z6 invariance — no free parameter.

2.2 UV Fixed Point

|β0| = NDirac + 2/Nf = 13/3; At the ultraviolet fixed point the beta function vanishes; the resulting UV fixed-point coupling $g^2 = 2\pi^2/3$, αmeta = π/6.

2.3 Condensate Action

$$S = 6Ne/be = 212.677,\ be = b0/(ZNP\ Nc) = 325/1152 \qquad (2)$$

where b0 = 13/6, ZNP = 32/25, Nc = 6. Three projections of S:

$$e^{-S} \to \rho\Lambda \qquad (3)$$

$$e^{-S/g^2} \to GN \qquad (4)$$

$$e\text{^}(-S(1+2/g^2)) \to \rho\Lambda GN^2 \qquad (5)$$

Their ratio $e\text{^}(S(1-3/(2\pi^2))) \approx 10^{78}$ resolves the hierarchy problem.

3 The Cosmological Constant

3.1 Observed Value (Planck 2018)

$$\rho\Lambda_obs = \Omega\Lambda \cdot (3H0^2c^2)/(8\pi GN) \cdot (\hbar c)^3 \text{ GeV} = 2.515 \times 10^{-47} \text{ GeV}^4.$$

3.2 Theorem: C = Nc/(27π^4)

Theorem 3.1 ([T2+]). $\rho\Lambda = (Nc/27\pi^4) \cdot \Lambda meta^4 \cdot e^{-S}$.

Proof sketch: (1) Z6 kink; (2) n = 2 Pöschl–Teller operator; (3) det′/det = 1/48 [T1]; (4) σdw/N0 = 1 [T1]; (5) Z3 Faddeev–Popov factor = 27. Full derivation: Appendix C. □

3.3 Numerical Verification

$\rho\Lambda_pred = 2.492 \times 10^{-47}$ GeV^4 (error: 0.93% against $\rho\Lambda$,obs = 2.515×10^{-47}).

4 Three Predictions from One Action

4.1 From GN to Λmeta to Observables

Causal Hierarchy and the SC Input Rationale. The PH has one dimensional empirical anchor: GN (Newton's constant, CODATA 2018). The neutrino sector carries one additional adimensional datum, $\Delta m^2_{21}/me^2$ (scale role; §11), not yet derived.

L(Z6, g^2, j) [pure theory] $\to$ condensate —Sakharov→ GN [SC] [anchor] $\to$ Λmeta = $\sqrt{}$(K(ξ)/GN) $e^{-S/2g^2}$ [derived scale] $\to$ me = 0.5089 MeV [genuine prediction]

Why GN, not me? GN sits two causal levels above me in the hierarchy. Using me as anchor is causally circular: it defines Λmeta via a quantity the theory is supposed to derive from Λmeta. With GN as anchor, me becomes a first-principles output with a falsifiable coupling: δme/me = $-\frac{1}{2}$ δGN/GN.

From the single dimensional SC input GN = 6.70883×10^{-39} GeV^{-2}, the condensation scale is:

$$\Lambda meta = \sqrt{}(K(\xi)/GN) \cdot \exp(-S/2g^2) = 1.2609 \times 10^{12} \text{ GeV. [SC]} \to \text{[T2]} \qquad (6)$$

Remark 4.2 (Λmeta is not an input). Λmeta is a fixed function of the single dimensional anchor, Λmeta = f(GN), with f built entirely from derived constants (K(ξ) [T1], S = 6Ne/be [T1/T2], $g^2 = 2\pi^2/3$ [T1]). It carries no independent empirical content and no continuous freedom: a second dimensional degree of freedom would break the parameter-free closure of the adimensional invariants $\rho\Lambda GN^2$ and GN me^2 (eq. 9; Table 4, F1–F2), which close at the 0.9% level with nothing to adjust. The Sakharov relation GN = K(ξ) $\Lambda meta^{-2}$ e^{-S/g^2} is this same definition inverted — one equation, not two constraints. The residual theoretical softness is localized and declared: be enters the exponent at [T2] (open problem P1), and its status is tracked by the Class A/B partition of Table 7.

The logical output cascade:

GN [SC] (dimensional empirical anchor)

↓

Λmeta = $\sqrt{(K(\xi)/GN)}\ e^{-S/2g^2}$ (derived scale, [T2])

↓

me (0.42%) · $\rho\Lambda$ (0.93%) · GN [self-consistency] (0.83%) · mDM (T2) · Neff (<1σ) · lepton spectrum (5–9%, P8b)

Open Problem 1 (P2/P12 [T2] — eliminating the last input). *Solving $\Phi(\Lambda) = \Lambda$ of the two-loop NJL kernel without observational input would deliver $L \to \Phi(\Lambda) = \Lambda \to \Lambda meta \to GN \to me$, reducing SC inputs from 1 to 0 and making both GN and me first-principles predictions.*

4.2 Dark Energy

$\rho\Lambda = C\ \Lambda meta^4\ e^{-S} = 2.492 \times 10^{-47}$ GeV^4 (0.93%, [T2+]).

4.3 Newton's Constant (Self-Consistency Check)

$$GN = K(\xi)\ \Lambda meta^{-2} \exp(-S/g^2),\ K(\xi) = 1.16335. \qquad (7)$$

Predicted: 6.653×10^{-39} GeV^{-2}; observed: 6.70883×10^{-39} GeV^{-2} (CODATA 2018, natural units); error: 0.83%, [T2]. GN emerges via the Sakharov induced-gravity mechanism [3] — a ground-state tensorial susceptibility of the condensate, not a mediator particle.

Theorem 4.1 (Falsifiable coupling, [T2]). *Differentiating the cascade law me = $\sqrt{(K(\xi)/GN)}\ exp(-S/2g^2 - Ne/be)$:*

$$\delta me/me = -\tfrac{1}{2} \cdot \delta GN/GN. \qquad (8)$$

Parameter-free; testable via variation-of-constants cosmology (prediction F2).

5 Master Invariant

$$\rho\Lambda GN^2 = (Nc/27\pi^4)\ K(\xi)^2 \exp[\ -6(\pi^2 + 3)Ne\ /\ (\pi^2\ be)\ ]. \qquad (9)$$

Predicted: 1.1214×10^{-123} (exactly GN-independent); observed: $2.515 \times 10^{-47} \times GN^2 = 1.1320 \times 10^{-123}$; error: 0.93% [T2+]. Λmeta cancels algebraically.

Remark 5.1 (Irreducibility of exactly one dimensional input). The invariant web ($\rho\Lambda GN^2$, GN·me^2, eq. 9) shows that Λmeta and GN cancel in every adimensional combination: all ratios close internally. Conversely, any dimensionful output requires exactly one dimensional input — a scale-free kernel cannot fix a scale. One dimensional anchor is therefore a structural floor of any theory producing dimensionful observables, not a deficiency of this one.

5.1 Closed Form, $\hbar$-Linearity, and Falsification Handles

Composing the registered chain ($GN\Lambda meta^2 = K(\xi)e^{-S/g^2}$; me = Λmeta $e^{-Ne/be}$) with the wall coefficient $C = Nc/(27\pi^4)$ yields the master invariant in closed structural form: $\rho\Lambda GN^2 = C{\cdot}K(\xi)^2{\cdot}e\text{^}(-S(1 + 3/\pi^2))$, with S = 6Ne/be = Nc·Ne/be = 69120/325 = 212.677 — the condensate action is exactly Nc cascade depths

(the registered $S/g^2 = 32.323$ reproduces this to 10^{-6}). Every factor is structural — {Nc, π, K(ξ), Ne, be} — with no dimensional input, which is why this combination is the invariant. Ledger of the 123 decades: ln C = -6.08; 2 ln K = +0.30; $-2S/g^2 = -64.65$; $-S = -212.68$. Intermediate identity, confronted and resolved: $\rho\Lambda = C{\cdot}me^6/\Lambda meta^2$ is the exact rearrangement of Theorem 3.1 ($\rho\Lambda = C{\cdot}\Lambda meta^4{\cdot}e^{-S}$) under the definitional $S = 6Ne/be$, since $e^{-S} = (me/\Lambda meta)^6$ — an identity, not a coincidence; the closed form above therefore inherits [T2+] from its components, and the $\leq 0.1\%$ criterion reduces to an input-rounding check.

$\hbar$-linearity. Restoring SI units, the unique dimensionless combination of {$\rho\Lambda$, GN, $\hbar$, c} is $\rho\Lambda GN^2\hbar/c^7$: the exponents (2, 1, -7) are forced by the three dimension equations, so the invariant is linear in $\hbar$ by necessity. Within the framework the linearity is structural: $\rho\Lambda$ is a one-loop functional determinant (Gel'fand–Yaglom) on the condensate background — an O($\hbar$) quantity. Stated without metaphor: at fixed GN, c, and the classical structure of the condensate, $\hbar \to 0$ sends $\rho\Lambda \to 0$ — the de Sitter component of the expansion is a first-order quantum effect. The standard cosmological-constant embarrassment presumes an additive sum of zero-point energies against a UV cutoff; here the invariant is a multiplicative semiclassical factor, e^($-$S(1+3/π^2)) with $S \approx 213$ — instanton-grade suppression, not a 123-decimal cancellation. Loop-order attributions to the dimensionful factors separately are convention-dependent and carried as interpretation; the $\hbar$-linearity of the invariant is exact.

Falsification handles (three, independent). (i) The numerical identity: with paper-precision inputs the closed form must land within $\leq 0.1\%$ (leading order exact); a persistent residual at O(1/S) = 0.47% is registered as the two-loop term of the $\hbar$-linear structure — with sign and size — not as failure; a residual $\gg$ 1/S demotes the identities to coincidence. (ii) The co-drift law: because $\rho\Lambda GN^2$ equals a pure structural number, any temporal or environmental drift of GN forces $\delta\rho\Lambda/\rho\Lambda = -2\ \delta GN/GN$ — jointly with F2 ($\delta me/me = -\frac{1}{2}\ \delta GN/GN$): one drift parameter, three observables (GN, me, $\rho\Lambda$), two exact ratios ($-\frac{1}{2}$, -2); a single inconsistent pair falsifies the web (F17). (iii) Any claim that the expansion is tied to $\hbar$ carries meaning only through these dimensionless statements; a drift of $\hbar$ alone is a unit convention, not an observable.

6 Composite Spectrum from Z6

Key definitions. A composite of N psions is a bound state of exactly N psion quanta; N is the body number. Constituents are indexed by k and written as a tuple, e.g. (0, 1, 2); chirality is abbreviated by an L/R prefix (L: $k \leq 2$, R: $k \geq 3$). A composite is stable if it is the lightest state in its (I, Q, J) sector. Observable composites satisfy $I \equiv (\Sigma ki) \bmod 6 \in \{0, 3\}$.

Two-body. $|Z3|^2 - 1 = 8$ colour-charged bilinears $\to$ eight gluons (Ng = 8, [T1]). The Z6-singlet spin-1 massless bilinear $\to$ photon.

Three-body (leptons and neutrinos).

Configuration	I	Q	ΣmJ	Identity
LLL (0, 1, 2)	3	-1	-9/2	e^- (unique Q = -1, Theorem 6.2)
LLR (0, 1, 5)	0	0	-3/2	n / $\nu\tau$ (dual-sector)
LLR (0, 2, 4)	0	0	-3/2	$\nu\mu$ / Σ^0-like (dual-sector)
LRR (0, 4, 5)	3	+1	3/2	Proton p (charge check: 1/9 + 4/9 + 4/9 = +1)
LLR (1, 2, 3)	0	0	-3/2	νe / Λ^0-like (dual-sector; generation ordering open)
LRR (1, 3, 5)	3	+1	3/2	Σ^+-like
LRR (2, 3, 4)	3	+1	3/2	Σ^+-like (p-alternative configuration)
RRR (3, 4, 5)	0	+2	9/2	Q = +2 exotic (BSM)

Table 2: The eight observable three-body composites (generated census; `gen_composite_tables.py`).

Theorem 6.2 (Unique electron, [T1]). *Among all C(6,3) = 20 three-body combinations, exactly one observable configuration carries Q = −1: the all-left state (0, 1, 2).*

Charged-lepton generations are therefore not distinct k-triples but the eigenvalue ladder of the leptonic transfer matrix on this single charge sector — mass ratios are $\lambda i/\lambda j$ (A.2.5) — with |mJ_R| ∈ {0, 3/2, 5/2} as excitation labels (Table 5). Neutrinos are the three neutral configurations; $\nu e \leftrightarrow (1, 2, 3)$ (generation ordering open, ν-mass sector).

Theorem 6.3 (Charge–chirality identity, [T1]). *For every observable three-body composite, Q = nR − 1, where nR counts right-handed constituents.*

The per-psion form is Theorem V.1 (Qk+3 = Qk + 1). Corollary: the electron is the unique all-L state — the chirality vacuum of the census; the same L-preference measured by $\varepsilon\psi$ (App. J) marks the electron as the extremal composite.

Standard Model comparison. In the SM, electric charge is a U(1) input, the Gell-Mann–Nishijima relation expresses Q through two further inputs (isospin, hypercharge), and the existence of exactly one stable charged lepton is an empirical fact. In the PH all three are counting theorems: charge measures chirality content, and electron uniqueness follows from Pauli exclusion (which restricts the ground-state census to distinct-k triples) combined with Z6 charge arithmetic (which admits exactly one Q = −1 triple).

Z_N structure. The deep form of Theorem 6.2 is universal: for every anchor group Z2M (M odd) the all-L extremal state is unique, with Qmin = −(M−1)/2. The literal form is a Z6 discriminant: the number of Q = −1 observables is C(M, (M−3)/2) = 1, 5, 21, 84 for M = 3, 5, 7, 9 (exact Z10 census in [2]). Two further facts select M = 3: the charge table is rational only for M = 3 among odd M ≥ 3 (the real cyclotomic field $Q(\zeta M)+$ is rational iff M ∈ {1, 2, 3, 4, 6}), and integer charge quantization of singlets already fails at M = 5 (fractional fifths in the exact Z10 census). Their elevation to a selection principle for Z6 is registered as conjecture C-Q-rat ([T3], [2]).

6.1 Pauli Structure: the SM as the Pauli-Trivial Sector

The Dirac kinetic term of eq. (1) fixes the bookkeeping: the psion carries spacetime spin-1/2 and a six-valued internal SU(2)meta label (the internal spin-5/2 multiplet), giving exactly twelve single-particle states — a counting resolved through the U-count adherence programme, whose registered decider (the first-principles derivation of the relic prefactor) fired on it, with the review of its two residuals recorded in [2]. Two Pauli thresholds follow. From N ≥ 7, label repetition is forced and the auto-singlet lemma ([T1]) applies: two fermions on the same label in the same orbital are a forced spin singlet. From N ≥ 13, full state repetition is forced and spatial antisymmetrization begins: psion nuclei proper start at N = 15, while N = 9 and 12 are s-wave states. For N = 9 the ground state is decided by a uniqueness theorem ([T1]): among the 32 neutral singlet candidates, exactly one is a pure s-wave configuration (nk ≤ 2) — (2, 2, 2, 1, 1, 1), three forced singlets plus a free-spin triplet on the distinct labels {3, 4, 5}, with mean-field JΣ energy maximally anti-aligned ($\langle Jz\rangle\langle \Sigma z\rangle = -81/2$). The free-spin triplet is the same configuration class as the three-body leptons (distinct labels, s-wave, three free spins); the observed lepton spin J = 1/2 anchors the spin selector, giving Jgs = 1/2 and g = 2 — the derived pair $(2, 2\pi/3)$ of the relic prefactor (§14.1). Conversely, every observable SM composite is Pauli-trivial: bilinears are psion–antipsion pairs, and all N = 3 identifications are distinct-k. The SM shadow of §F.3 acquires a structural reading: the Standard Model is exactly the sector of the spectrum that requires no Pauli structure. The historical resonance is inverted with respect to the quark model: there, the Pauli crisis of the Δ++ forced the invention of colour; here, the observable sector never meets the crisis, and the Pauli pressure discharges entirely into the dark sector — as forced singlets at N = 9, 12 and nuclear structure from N = 15. Stability is unaffected: shell corrections below 13% of the N = 9 cascade exponent

cannot alter any Theorem KS fragmentation bound.

Theorem 6.1 (Nν = 3, [T1]). *All C(6,3) = 20 three-body Z6-colour-singlet combinations yield exactly three neutral stable states — the three active neutrino generations.*

Dark matter. 39 stable Z6-singlet states at N ∈ {9, 12, 15, 18, 21, 27}. mDM = 2Λmeta/NLandau = 7.205 × 10^{10} GeV, NLandau = 35.

7 Physical Interpretation

S ≈ 212.7 resolves the hierarchy problem: e^(S(1−3/(2π^2))) = e^180.4 ≈ 10^{78}.

7.1 The Spin-2 State hAB and the Graviton (PC1, [T2])

GN arises via Sakharov induced-gravity (ground-state susceptibility of the condensate). hAB is a massive spin-2 composite at ∼0.613 GeV, distinct from the GR graviton. It satisfies the Weinberg–Witten theorem and decays to 2γ above threshold. Full derivation: Appendix J.

8 Kinematic Stability: Theorem KS

The referee objection that "Z6-algebraic stability does not imply kinematic stability" is addressed here by a direct calculation across the complete Z6 composite spectrum.

8.1 The Cascade Mass Function

The mass of a state at cascade level ℓ is given by the cascade formula:

$$\mathrm{m}(\ell) = \Lambda\mathrm{meta}\,\exp(-\ell/\mathrm{be}),\ \ell \in \mathbb{N}^{+}. \qquad (10)$$

This is a strictly decreasing function of ℓ: each rung down the cascade suppresses the mass by a factor exp(−1/be) = exp(−1152/325) ≈ 1/34.6.

Cascade level is not body number. The level ℓ is the ladder rung, an independent quantum number; it is not a function of the body number N. Three anchors fix this: h_{AB} (N = 2, ℓ = 8), X17 (N = 2, ℓ = 9) and the electron (N = 3, ℓ = 10) — two states of equal N at different levels, and a heavier N at a lighter level (repository [2], `scripts/cascade_level_anchors.py`). Identifications at leading order: m(8) ≈ 610 MeV (nucleon and h_{AB} scale), m(9) = 17.62 MeV (X17), m(10) = 0.5089 MeV (electron). The stable dark singlets of §14 are labelled by body number N ∈ {9, 12, 15, 18, 21, 27}; their mass is *not* m(N) but the Landau-pole value m_{DM} = 2Λmeta/35 (§14.2), at the top of the cascade.

Status note (v86, WP-KILL). Taking $\ell_{\mathrm{eff}}(m) = 10 - \ln(m/0.5089\,\mathrm{MeV})/\ln 34.6$ with the electron as anchor, 25 known hadron ground states (light, strange, charm, bottom) fall between $\ell_{\mathrm{eff}} = 7.2$ and 8.4; 9 of 25 lie within 0.15 of an integer rung, against a mean of 7.5 for a randomly offset grid ($p = 0.34$). Known hadrons therefore show no alignment with the electron-anchored ladder; their clustering follows the QCD scale, and the nucleon sits 0.12 rung from $\ell = 8$, the pion 0.42 from it. The ladder is kept as a structural statement for states not yet observed, not as a description of the known hadron spectrum. The masses in this test were entered from memory of the PDG tables and are to be verified, but the result is far from the decision threshold. In the same registered test the static-hexagon radius of the Σ^-, $R = +0.846$ against 0.862 ± 0.297 (SELEX),

was consistent; 17% of random charge assignments would also fall inside 1σ, so this is consistency, not confirmation. Verdict of the registered rule: undetermined (not refuted, not supported).

8.2 Theorem KS: Kinematic Stability

Theorem 8.1 (KS — Kinematic Stability, [T1]). *For any Z6 composite of body number $N \geq 2$ and any partition $N = N1 + N2 + \cdots + Nk$ with $k \geq 2$, $Ni \geq 1$:*

$$\Sigma\, m(Ni)/m(N) = \Sigma \exp((N - Ni)/be) \geq k \exp((k-1)/be) \geq 2 \exp(1/be) \approx 69.3. \qquad (11)$$

In particular, $\Sigma\, m(Ni) > m(N)$ for ALL fragmentation decays, universally.

Proof. Since $N - Ni = \Sigma_{j\neq i} Nj \geq k - 1$ (each $Nj \geq 1$), each term $\exp((N - Ni)/be) \geq \exp((k-1)/be)$. Summing k terms gives $\Sigma \exp((N - Ni)/be) \geq k \exp((k-1)/be)$, which for $k \geq 2$ and $Ni \geq 1$ satisfies $\geq 2 \exp(1/be) \approx 69.3 \gg 1$. □

Cascade level	m(ℓ)	Min k = 2 ratio	Min k = 3 ratio
$\ell = 2$	1.05×10^9 GeV	69.3	—
$\ell = 8$ (nucleon / h_{AB} scale)	610 MeV	2.88×10^6	1.82×10^9
$\ell = 10$ (electron)	0.509 MeV	9.96×10^7	1.21×10^{11}
$\ell = 9$ (X17)	17.6 MeV	5.12×10^7	5.17×10^9
$\ell = 12$	4.25×10^{-7} GeV	3.45×10^9	6.20×10^{12}
$\ell = 27$	3.44×10^{-30} GeV	3.67×10^{21}	1.54×10^{28}

Table 3: Minimum ratio $\Sigma m(\ell_i)/m(\ell)$ of the cascade mass function over partitions $\ell = \Sigma \ell_i$ into k parts (the minimum is the balanced partition; — : no k-part partition exists). Rows are cascade levels, not dark-matter states (dark-matter mass: §14.2). Physical scope: see the status note on Theorem KS. Erratum: earlier prints gave 3597, 5.60×10^{11} and 8.05×10^{13} for the k = 3 entries at ℓ = 2, 8, 10; the first was the bound $3e^{2/be}$, the other two were overstated. Inequality (11) holds for all.

Status note (under review). Theorem KS, as stated, treats the mass as a function of body number and partitions body number, $N = \Sigma N_i$. Fragmentation conserves psion (body) number, but the cascade level is independent of body number (§8.1), and no rule is yet derived that fixes the levels of the fragments from the level of the parent. The inequality (11) remains exact as arithmetic on the mass function; its application to physical fragmentation decays, and hence Corollary 8.2, is suspended pending that rule. Dark-matter stability (§14.1) does not depend on it: it rests on the exhaustive Z6 fragmentation census (Theorem DM.1, `proofs/P_dark_matter.py`) and on psion-number conservation. The module `proofs/P_kinematic_stability.py` verifies inequality (11) on the level partitions only and registers this gap as a scope guard. **v86:** the missing rule is identified as a conserved baryon (rung) label required by nucleon stability (§13.4); it is open problem P14-B.

8.3 Corollary: Z6 is Necessary and Sufficient

Corollary 8.2 ([T1]). *If Z6 algebra forbids all decay channels of a composite of body number N (topological stability), then the composite is fully dynamically stable: all fragmentation decays — whether Z6-allowed or not — are kinematically forbidden by Theorem 8.1.*

Therefore: Z6 algebraic stability is both necessary and sufficient for full dynamical stability, including 4-momentum conservation.

Scope and caveats. Theorem KS covers fragmentation decays ($N \to \Sigma Ni$, $k \geq 2$). It does not cover:

1. Radiative transitions ($N \to N + \gamma$, same body number): excluded because m(N) is the ground-state mass by definition of the cascade. No lower-energy state of the same N exists within the PH spectrum.

2. Weak rearrangements ($\Delta N = 0$, e.g. β decay): covered separately by Z6 charge conservation and the selection rules of App. H. Rearrangement decays ($\Delta N = 0$, e.g. neutron β-decay: (0, 1, 5) $\to$ (0, 1, 2) + (0, 4, 5) + $\bar{\nu}$) are covered by Z6 charge selection rules (App. H) and are outside the scope of Theorem KS, which addresses fragmentation only.

Anti-matter. Anti-composites have the same body number N and the same cascade mass m(N) by CPT symmetry. Theorem KS applies to anti-composites by charge conjugation.

T-level. Theorem KS is [T1]: the proof is an algebraic identity on the cascade mass function. The only T2 input is $m(N) = \Lambda_{meta} \exp(-N/b_e)$ itself (which inherits T2 from the leading-order cascade). Verification in companion repository [2], module `proofs/P_kinematic_stability.py` (11 claims; 33 tests in `tests/test_kinematic_stability.py`): inequality (11) on all 1,295,920 partitions of every level $2 \leq \ell \leq 50$, the balanced-partition minimum, Table 3, the level/body-number scope guard, and the delegated DM.1 census. Its physical application is subject to the status note above.

9 Falsifiable Predictions — Complete Registry

ID	Prediction	Value / Error	T-level	E	Experiment
F1	$\rho_\Lambda G N^2$ (Λ_{meta} cancels)	1.121×10^{-123} | 0.93%	[T2+]	E3	Precision cosmology
F2	$G N \times m_e^2$ ($\Rightarrow \delta m_e/m_e = -\frac{1}{2}\,\delta G N/G N$)	1.737×10^{-45} | 0.83%	[T2]	E3/E1	CODATA; clock drift
F6	$N_\nu = 3$ (counting theorem)	3 exact	[T1]	E2	LEP; FCC-ee
F9	$N_g = 8$ ($\lvert Z_3\rvert^2 - 1$)	8 exact	[T1]	E2	Lattice QCD
F7	$N_{eff} = 3.054$	3.054 | $<1\sigma$	[T2]	E3	CMB-S4 (§9.1: non-discriminant vs SM)
F12	$\eta_B \leftrightarrow N_{eff}$ via $\varepsilon_\psi = 0.1479$	suspended (v86, §13.3)	—	—	CMB-S4 + BBN
F5a	No dark-matter state outside the 39 singlets at m_{DM}: any direct-detection DM signal at $m \neq m_{DM}$ falsifies	exclusion	[T2]	E2	XENONnT; LZ; PandaX
F5b	$\sigma_{DM-n}(m_{DM}) \sim 10^{-54}$ cm^2 (magnetic dipole), conditional on zero proto-colour coupling of dark singlets	null at all foreseeable exposures	[T4]	E1	XENONnT; LZ; XLZD
F8	UHECR excess above 10^{22} eV	above GZK	[T2]	E1	Auger; GRAND
F13	γ-line at 72 EeV	72 EeV	[T2]	E1	AugerPrime
F3	ν_τ/ν_e asymmetry A_{te} of the astrophysical flux, conditional on P-PMNS (§17.3); +7% would require $\sin^2\theta_{23} \approx 0.574$ with $\delta_{CP} \approx \pi$	not derived	[T4]	E1	IceCube-Gen2; DUNE, Hyper-K (θ_{23}, δ_{CP})
F10	GN phase transition at $T = \Lambda_{meta}$	GN changes $10^{15}\times$	[T2]	E1	CMB-S4 tensors
F4	Proto-quark charges at $Q^2 \sim \Lambda_{meta}^2$	DIS deviation	[T3]	E1	Future DIS
F14	Cstable(N) $\in \{29, 30, 31\}$	$\sim$30/level	[T3]	E2	Computational (internal)
F11	DIS $\Sigma e^2 q = 47/27$ at $Q^2 \sim \Lambda_{meta}^2$	47/27	[T4]	E1	Future DIS ($\gtrsim \Lambda_{meta}$)
F15	X17 singlet at 17.62 MeV †	17.62 MeV | 3.59%	[T3]	E4	ATOMKI vs MEG II/PADME †
F16	hAB: γ branching = 5 : 3	5/3 ratio	[T2]	E1	Belle II
F17	Co-drift $\delta\rho_\Lambda/\rho_\Lambda = -2\,\delta G N/G N$ (joint with F2: one drift parameter, ratios $-\frac{1}{2}$ and -2)	exact ratios	[T2]	E1	GN-drift (LLR/BBN/CMB) $\times$ w(z) constancy

Table 4: All 17 falsifiable predictions of the PH (18 rows: F5 is split into F5a and F5b), with T-level (derivation status) and E-class (evidential status).

E-classification. T-levels grade how a prediction is derived; E-classes grade what its agreement is worth.

E1 — novel, forward-looking, discriminating: made before the measurement, distinguishable from rival frameworks, data pending on a schedule; the only class with conversion power. E2 — hard falsifier: forbidden or fixed by theorem; passing is shared with existing knowledge (zero conversion power), but violation kills the framework with no parameter to adjust — the PH forbids by T1 theorem what the Standard Model merely does not contain. E3 — postdiction of established data: a consistency scoreboard; necessary, never sufficient. E4 — postdiction of a disputed anomaly: the weakest class; carried for completeness under quarantine (K.8 discipline). Current census: E1 × 10, E2 × 4, E3 × 3, E4 × 1 (F5 is split into F5a, E2, and F5b, E1). The framework's scientific weight rests on the E2 rigidity today and on the E1 set tomorrow; the E3 set, however precise, cannot convert.

† *X17 (status mid-2026): MEG II excludes the Atomki ^{8}Be observation at 94% CL (2025); PADME reports an independent 2.5σ excess at 16.90 MeV — mutually conflicting. If the PADME value consolidates, the F15 prediction (17.62) moves further from data (3.6% → 4.3%). MEG II (PSI), JINR (Dubna), and NA64 (CERN) found no signal at the Atomki claimed mass and coupling. Experimental status disputed; registered as T3 candidate pending independent confirmation.*

F6 and F9 are T1 hard falsifiers — any detection of a fourth neutrino or ninth gluon immediately refutes the framework. F7 and F12 are structurally linked through $\varepsilon\psi = 0.1479$; this two-observable, one-parameter test is uniquely distinct from leptogenesis. The CF-CC structural prediction is discussed in Appendix K §K.8.

P8b dependency. None of F1–F17 above rests on the lepton-mass ratios m_μ/m_e, m_τ/m_e or their P8b refinement (§12 Table 5, §A.3.1): those ratios are reported as structural results but are not themselves registered in this falsifiability census. Consequently, closure or non-closure of P8b Passo A.3 ([T2], App. A) does not add or remove any entry from Table 4; it bears only on the precision claimed for the lepton-mass sector (Appendix L.4.4–L.4.5), not on the falsifiability registry.

9.1 Notes on Selected Predictions

F7 and F12 (Neff–ηB structural link). *Status (v86): F12 is suspended, because the η_B chain is withdrawn (§13.3); F7 is retained with the status note of §13.5.* Both Neff = 3.054 and $\eta\mathrm{B} = 6.02 \times 10^{-10}$ arise from the single parameter $\varepsilon\psi = 0.1479$ fixed by SU(2)meta geometry (§12). This structural link — one parameter, two independent observables — is uniquely testable by CMB-S4 combined with BBN likelihoods and cannot be reproduced by leptogenesis, which requires right-handed neutrinos absent from the PH spectrum. Honesty note: the PH−SM discriminant in Neff is $3.054 - 3.044 = 0.010$, below even the CMB-S4 target $\sigma(\mathrm{Neff}) \approx 0.03$; F12 can falsify PH and the SM jointly against a third value, but cannot separate them — the $\varepsilon\psi$ two-observable link, not Neff alone, is the discriminating structure.

F2 (variation-of-constants co-drift, honesty note). The exact ratio $\delta m_e/m_e = -\frac{1}{2}\,\delta G_N/G_N$ is a genuine structural prediction (zero free parameters once Λmeta is fixed), but it is not by itself PH-specific: any scalar-tensor gravity in which G_N and m_e share a common dilaton-like scalar predicts *some* fixed linear relation between the two drifts, generically. What is PH-specific is the particular coefficient $-\frac{1}{2}$ (forced by the cascade's Λmeta-dependence, eq. 6) and its joint closure with F17 ($\delta\rho_\Lambda/\rho_\Lambda = -2\,\delta G_N/G_N$, same single drift parameter, three observables, two fixed ratios). A measured pair of ratios inconsistent with $(-\frac{1}{2}, -2)$ falsifies PH specifically; a measured pair consistent with some linear relation only confirms the general scalar-tensor class, not PH against its rivals. F2 in isolation is therefore weaker evidence than Table 4's E1 label suggests; the discriminating power is in the F2–F17 joint constraint, not F2 alone.

F5 (direct detection): rationale for the F5a/F5b split and kill criterion, registered ex ante (2 Oct 2026). *Why split.* Earlier releases carried F5 as $\sigma_{\mathrm{DM-N}} < 10^{-50}$ cm^2 at [T2], E2. No derivation of that number exists in this paper or in the companion repository [2], so neither label is earned: [T2] requires a derived and verified mechanism, and E2 requires a prohibition fixed by theorem. The number is retired. What carries falsification power is structural and is kept as **F5a**: Theorem DM.1 ([T1]) and the canonical mass (§14.2,

[T2]) leave no dark-matter state outside the 39 singlets at m_{DM}, so a signal at any other mass is a component the PH spectrum lacks and contradicts the Ω_{DM} accounting of §14.3. It is a hard prohibition with nothing to adjust: [T2], E2. The rate is separated as **F5b** at [T4] (mechanism identified, dimensional estimates only). For the Landau-pole state (Q = 0, charged constituents, size $\sim 1/m_{DM}$): magnetic-dipole scattering $\sim 10^{-54}$ cm^2, charge-radius scattering $\sim 10^{-75}$ cm^2. Xenon sensitivity scales linearly with mass at high mass and reaches only $\sim 10^{-39}$ cm^2 at m_{DM} (rough extrapolation), so F5b predicts a null result with ~ 15 orders of margin, with one exception: a coupling of dark singlets to the proto-colour mediator (the X17 family, §A.5) of nucleon-like strength gives $\gtrsim 10^{-38}$ cm^2. *Promotion path.* [T3] once the proto-colour form factor of the N = 9 singlet is set up as a calculation (tied to P25, the Z6 composite gauge vertex); [T2] once that coupling is derived or shown to vanish (candidate argument to test: proto-colour singlets decouple at tree level, as neutral hadrons do from the photon). *Kinematic fact.* Under the canonical mass assignment (§14.2) all 39 stable dark singlets sit at m_{DM}; no stable neutral singlet exists at $N \leq 6$ (Theorem DM.1), and no other PH state is a halo candidate. The maximum xenon recoil at m_{DM} is ~ 1.7 MeV, so kinematics never excludes a signal: the rate does. At $m_{DM} = 7.2 \times 10^{10}$ GeV the local number density is $\sim 4 \times 10^{-12}$ cm^{-3}, so any detectable rate in a tonne-scale detector requires $\sigma \gtrsim 10^{-39}$ cm^2, some fifteen orders above the F5b estimate. *Kill criterion.* The PH dark sector is falsified by a dark-matter nuclear-recoil signal that (i) reaches $\geq 5\sigma$ global significance in one experiment, (ii) is confirmed at $\geq 3\sigma$ by an independent detector or independent exposure, and (iii) is not attributable to known backgrounds (neutrons, CEνNS, detector artefacts) by the collaborations' own analyses. Mass and operator do not matter: a signal at the Landau-pole mass violates F5b, and a signal at any other mass violates F5a. Not falsifying: single events, sub-5σ excesses, electron-recoil-only excesses. *Current status.* The LZ event of September 2026 (one nuclear recoil at 248 $\pm$ 23 (stat) $\pm$ 23 (sys) keV, 2.6σ global, 3.4σ local; arXiv:2609.02823) is below threshold and is carried as a watch item under E4 quarantine.

F10 (gravitational phase transition). Before the condensate forms (T > Λmeta), GN(T) $\sim 12\pi^2/(Nc\Lambda meta^2)$ $\sim 10^{15}$ GN_obs. The Z6 condensation at T = Λmeta is a first-order gravitational phase transition reducing GN by 15 orders of magnitude. This imprints on primordial tensor spectra and is in principle detectable via the CMB-S4 tensor-to-scalar ratio.

F11 (DIS structure function at Λmeta, [T4]). $\Sigma e^2 q = 47/27$. At $Q^2 \ll \Lambda meta^2$, PH and QCD give identical DIS predictions. At $Q^2 \sim \Lambda meta^2 \sim 10^{24}$ GeV^2, the proto-quark charges +1/9, +4/9, +5/9 become individually resolvable, shifting the charge-weighted structure function sum from the SM value 1.00 to $\Sigma e^2 q = (1 + 16 + 25)/81 \times 3 = 47/27 \approx 1.74$. Beyond current technology but precisely computable.

G1–G3 (gravitational sector, App. K). The driving frequency of [G3] is fdrive $\in$ [0.38, 0.43] GHz with Edrive $\approx 1.6 \times 10^{-6}$ eV, fixed by self-consistency of the susceptibility relation (CN-1). The factor 2 in νJ follows from order-parameter bilinearity (Lemma J.1, [T2+]). A quality factor Q $\sim 10^{10}$ (typical of Josephson junctions) would reduce the threshold driving energy density to $\sim 10^{24}$ GeV^4/m^3 — beyond current technology but computable. Classifications: [G1][T2+], [G2][T2] conditional on CEC, [G3][T3]. Full treatment: App. K.

10 Lepton Mass Spectrum

Theorem 10.1 (Ne = 10, [T1]). *Ne = Nγ + Ng + Nφ = 1 + 8 + 1 = 10.*

Leading-order lepton mass formula (A = 13/12, p = 5/6 from SU(2)meta Clebsch–Gordan structure, [T2]):

$$mn/me = \exp(A\ |mJ_R|^{p} / be). \qquad (12)$$

Lepton	|mJ_R|	m_pred/me	m_pred	Error
e^-	0	1	0.511 MeV (anchor)	0.0%

μ^-	3/2	218	111.4 MeV	5.4%
τ^-	5/2	3795	1939 MeV	9.1%

Table 5: Lepton masses (leading order, [T2]). Residual errors target P8b.

11 Neutrino Masses

SC input: $\Delta m^2_{21} = 7.530 \times 10^{-5}$ eV^2 — equivalently the adimensional ratio $\Delta m^2_{21}/m_e^2 = 2.908 \times 10^{-16}$, the framework's second (and only adimensional) empirical datum, serving purely as the neutrino-sector scale. Normal hierarchy and the mass ratios are the sector's predictive content. A closed-form derivation candidate, $\Delta m^2_{21} = (18/25)\, m(15)^2$ (-0.07%), is registered at T4 in the companion repository (C-Δm21; promotion path P26-C); it is not promoted here.

ν	**mν_pred**	**Δm^2_pred**	**Δm^2_obs**	**Error**
νe	7.50×10^{-9} eV	—	< KATRIN	—
$\nu\mu$	8.68×10^{-3} eV	7.530×10^{-5} eV^2	7.530×10^{-5}	0.0% (SC)
$\nu\tau$	4.73×10^{-2} eV	2.239×10^{-3} eV^2	2.453×10^{-3}	8.7%

Table 6: Neutrino mass predictions.

12 Cosmic Accountability

Observable	**Predicted**	**Observed**	**Error**	**Status**
$\rho\Lambda$ (GeV^4)	2.492×10^{-47}	2.515×10^{-47}	0.93%	[T2+]
Δm^2_{21} (eV^2)	7.530×10^{-5}	7.530×10^{-5}	0.0%	SC input
GN (GeV^{-2})	6.70883×10^{-39}	6.70883×10^{-39}	0.0%	[SC][1]
me	0.5089 MeV	0.511 MeV	0.42%	cascade output
GN (GeV^{-2})	6.653×10^{-39}	6.709×10^{-39}	0.83%	[T2][1]
$\rho\Lambda$GN^2	1.121×10^{-123}	1.132×10^{-123}	0.93%	[T2+]
ηB	withdrawn (§13.3)	6.13×10^{-10}	—	open; candidate §13.7
ΩDM/Ωb	5.43 / 5.64	5.364	1.2% / 5.1%	direct/inference; mutual 3.9% (§14.3)
mμ/me	218	206.8	5.4%	b-dep.
mτ/me	3795	3477	9.1%	b-dep.
Δm^2_{31} (eV^2)	2.239×10^{-3}	2.453×10^{-3}	8.7%	indep.
Nν	3	3	exact	theorem
Ng	8	8	exact	theorem
Neff	3.054	2.99 ± 0.17	$< 1\sigma$	indep.; SM: 3.044

Table 7: Cosmic accountability table. SC = self-consistency input. b-dep. = sensitive to cascade parameter b at $\lesssim$ 35% per 1% change in b.

Remark 12.1 (Cascade output vs. empirical anchor). The cascade output is unique: me = Λmeta exp($-$Ne/be) = $\sqrt{(K(\xi)/GN)}$ exp($-$S/2g^2 $-$ Ne/be) = 0.5089 MeV (0.42% against the observed 0.511 MeV), since K(ξ) sits inside Λmeta (eq. 6). A K(ξ)-free cascade would give 0.4714 MeV (-7.8%); the prefactor is part of Λmeta itself (eq. 6). In P8b, the observed 0.511 MeV serves only as a reference for computing lepton mass ratios mμ/me, mτ/me; it does not enter the GN $\to$ Λmeta $\to$ me derivation chain.

[1] This row is the single dimensional input (SC) itself, entered at its measured value by construction — 0.0% is not a prediction. The second GN row below is the independent, genuinely predicted quantity: GN recovered from the Λmeta$\to$me cascade run in reverse, at [T2], with a non-zero 0.83% residual because it is not fitted to the input.

13 Baryogenesis: Status after the P14 Audit

Summary. Earlier releases carried $\eta_B = 6.02 \times 10^{-10}$ as a derived output at [T2]. The P14 audit withdraws that claim. Equation (1) conserves psion number exactly, field-theory CP is exact, and the Standard-Model sphaleron conversion used in the former eq. (14) does not apply in the PH. Neither eq. (1) nor its minimal extension produces the asymmetry (13.3). One candidate mechanism remains, level bifurcation in the pair production of unconfined composites after inflationary reheating (13.7). Its structure follows from symmetry, but its magnitude is not yet explained. This section replaces the former 13.1–13.5.

13.1 What Equation (1) Conserves

Psion number. N_ψ is the Noether charge of the overall phase and is exactly conserved by every term of eq. (1). In the broken phase it is the only exact additive charge (exact charge audit over 20 triples and four condensate variants, zero further candidates).

Continuous internal symmetry. The full Lie algebra of internal transformations leaving $(\bar\psi J^a \psi)(\bar\psi \Sigma^a \psi)$ invariant is two-dimensional and abelian, $U(1)_N \times U(1)_M$, with $M = \sum m_k$; no continuous chiral extension survives the $G_{\rm UV}$ term. The winding condensate of App. J.6.2 breaks $U(1)_M$, leaving a residual discrete charge $D = (2M - 6Q + 3) \bmod 4$.

CP. The coupling tensor is real and ε_ψ is CP-even: field-theory CP is exact. The Z6 vacua break CP spontaneously, but conjugate vacua are degenerate ($\Delta C(s)$ symmetric).

Antimatter. Only one reading is consistent with observed reactions: antiparticles are composites of antipsions ($N_\psi = -3$, charge $-Q$). The index inversion $k \to (6-k) \bmod 6$ (App. A.4) describes the Z6 label of the antipsion, not a psion composite. The alternative reading (antiparticle = psion triple with inverted indices) fails psion-number conservation already in $\beta^\pm$ decay, fails charge conservation in $n \to p e^- \bar\nu$ and $\mu^- \to e^- \bar\nu_e \nu_\mu$, and gives $\bar\Sigma^+$ the charge $+1$.

13.2 The Sakharov Conditions [16], Revisited

(1) Baryon-number violation. Nothing built from psions violates N_ψ, including an emergent sphaleron. With $N_\psi/3 = B + L$ (13.4), $B + L$ is conserved and only $B - L$ can change: the reverse of the Standard Model. The sphaleron factor 8/23 and the coefficient $C_{\rm core}$ of the former eq. (14) therefore have no basis in the PH. In addition, $C_{\rm core} = 8.4434 \times 10^{-4}$ could not be reproduced from its stated inputs.

(2) C and CP violation. Eq. (1) supplies none (13.1). The former statement that the chirality-flip amplitude $\propto (m_e/\Lambda_{\rm meta})^2$ is a CP-odd phase is withdrawn: a real amplitude carries no phase.

(3) Departure from equilibrium. Available: the formation of composites is out of equilibrium when the contact rate is comparable to the expansion rate (13.7). Temperature histories alone redistribute abundances among cascade levels but never create an asymmetry.

13.3 Withdrawn Results and Closed Mechanisms

Withdrawn. (i) The former eqs. (14)–(15), $\eta_B = C_{\rm core}(g_{1/2}/g_S)\varepsilon_\psi = 6.02 \times 10^{-10}$: the sphaleron input has no PH basis, and $C_{\rm core}$ is not reproducible. (ii) Eqs. (37)–(38) of App. J: $\xi^* = 434.09$ was computed from the target η_B, so eq. (38) is an identity, not a check. Independent routes to ξ^* give $\xi^* \le 1.08$ (Potts Z3; second-order NJL transition with $T_c = 9.6$ MeV) and $\xi^* = 3.6$–39 (Coleman–Weinberg with thermal

inflation). (iii) The claim that the PH satisfies all three Sakharov conditions without additional ingredients. (iv) Prediction F12 is suspended (§9).

Reclassified. $\varepsilon_\psi = 25/169$ remains a result of $\mathrm{SU}(2)_{\mathrm{meta}}$ geometry (App. J, conditional on P23). Its link to η_B is not established.

Closed mechanisms. Each was tested under a pre-registered definition, and each fails for a stated reason:

- Emergent sphalerons: they conserve N_ψ.
- Z6 domains with spontaneous CP: they form at $T \approx 3$ MeV, when assembly is frozen ($\Gamma/H = 3\times10^{-37}$). Their maximal comoving size today is ~ 30 pc, against the $\gtrsim 1$ Gpc required of antimatter regions [19]. Degenerate conjugate vacua give a zero global average.
- Temperature histories: no asymmetry from symmetric initial data.
- CP terms built from PH operators: 16 commutator and rearrangement patterns, alone or in pairs, fit the observed composition no better than random directions.
- A phase inserted in $J\cdot\Sigma$: it generates an asymmetry, but exotic triples receive 2–2500 times the proton asymmetry. Without a conserved label, no CP source selects the observed composites.
- Species mixing in the minimal extension of 13.4: the proton lifetime bounds its coefficient to $c \leq 1.3\times10^{-8}$, which places the active window above 2.7×10^{15} GeV, outside the contact description.

13.4 Baryon Number and the Level Rule

What observation requires. With the cascade level treated as a quantum number, the observed reactions $\beta^\pm$ decay, e^- capture, $\pi^+ \to \mu^+\nu_\mu$, μ decay, $\nu_\mu n \to \mu^- p$, $\bar\nu_e p \to e^+ n$, $\Lambda \to p\pi^-, n\pi^0$, $\Sigma^0 \to \Lambda\gamma$ and neutrino oscillations require exactly two additive charges: B (states of the level-8 class) and L (level-10 and neutrino classes), with $N_\psi/3 = B + L$.

What eq. (1) allows. With only N_ψ and Q conserved, seven unobserved processes are allowed: $p \to \nu\pi^+$, $n \to \nu\pi^0$, $n \to 3\nu$, $pe^- \to \nu_\mu\nu_\tau$, and three level jumps within one triple ($n \to \nu_\tau\gamma$, $\Lambda^0 \to \nu_e\gamma$, $\Sigma^0 \to \nu_\mu\gamma$). All have $\Delta B = -\Delta L$. Against the measured limits ($p \to \bar\nu\pi^+ > 3.9\times10^{32}$ yr and $n \to \bar\nu\pi^0 > 1.1\times10^{33}$ yr [17]; $n \to$ invisible $> 5.8\times10^{29}$ yr [18]), the required amplitude suppression is $\lesssim 10^{-8}$ relative to a contact operator at Λ_{meta}. For same-triple jumps it is $\lesssim 5\times10^{-30}$ relative to an electromagnetic transition (reference lifetime 10^{29} yr).

What eq. (1) cannot supply. B is not a Noether charge: it lies outside the span of (M, Q, N_ψ). It is not topological: the vacuum manifold has $\pi_2 = \pi_3 = 0$, so no skyrmion-like baryon number exists [21]. And the level is not an internal degree of freedom, since states are labelled by the triple alone. Hence the dual-sector identification of §6 and L.4.4, in which the same triple is either a baryon or a neutrino ((0,1,5) $= n/\nu_\tau$, (0,2,4) $= \Sigma^0/\nu_\mu$, (1,2,3) $= \Lambda^0/\nu_e$), is incompatible with neutron stability as eq. (1) stands. Correction to §6: there are five neutral three-body triples, not three. They are (0,1,4), (0,1,5), (0,2,4), (0,2,5) and (1,2,3); the two additional ones carry $D = 2$.

Minimal extension (registered, not adopted). A two-valued species index, $\psi \to \psi_s$ with $s \in \{B, L\}$ and the currents of eq. (1) summed over species, preserves $U(1)_B \times U(1)_L$ if $SU(2)_{\mathrm{species}}$ is broken. All observed reactions are allowed, and all seven unobserved processes are forbidden exactly. Its costs:

- The L sector contains six positively charged matter leptons that must be heavy.

- The numerical predictions survive only if the species index does not enter $|\beta_0| = N_{\text{Dirac}} + 2/N_f$. Doubling N_f gives $m_e = 0.065$ MeV.
- A label carried by the constituents cannot be chosen at formation.

This is recorded as open problem **P14-B**: a conserved baryon (rung) label.

13.5 Link to N_{eff} (status note)

$$\Delta\text{Neff_PH} = 3 \cdot (15/7\pi^2) \cdot (\mu\nu/\text{T})^2 = 0.010, \text{ Neff_PH} = 3.054. \qquad (16)$$

Equation (16) is retained, but its shared-parameter status with η_B lapses with 13.3. If $\mu_\nu/T = 0.123$ is read as a lepton-number asymmetry, it conflicts with psion-number bookkeeping. With total $N_\psi = 0$, $B = -L$ and charge neutrality, the antineutrino excess is two per hydrogen atom, a degeneracy $\xi_\nu \approx 2.5 \times 10^{-9}$, some 5×10^7 times smaller than 0.123. Equation (16) must therefore rest on a different reading of μ_ν or be reclassified.

13.6 Structural Predictions

(a) No nucleon decay with $\Delta B = \Delta L$: $p \to e^+\pi^0$ and similar modes change N_ψ and are forbidden exactly [T1]. **(b)** Any nucleon decay, if observed, has $\Delta B = -\Delta L$ ($p \to \nu\pi^+$, $n \to \nu\pi^0$, $n \to 3\nu$). **(c)** No $0\nu\beta\beta$ and no n–$\bar{n}$ oscillation, since both change N_ψ. **(d)** If the asymmetry formed with total $N_\psi = 0$: a cosmic antineutrino excess of two per hydrogen atom. **(e)** Dark matter as a thermal relic of the cascade top would require level ≈ 4.8 (≈ 56 TeV); the canonical $m_{\text{DM}} = 7.2 \times 10^{10}$ GeV requires a non-thermal origin (§14.3).

13.7 A Candidate Mechanism: Level Bifurcation at Reheating

Unconfined spectrum. By the confinement rule of App. H/I (summed index I mod $6 \in \{0,3\}$), exactly eight triples are unconfined: (0,1,2), (0,1,5), (0,2,4), (0,4,5), (1,2,3), (1,3,5), (2,3,4) and (3,4,5). These are the triples with one constituent of each proto-colour, and they are the states identified with observed particles. All fractional-charge triples and both $D = 2$ neutrals are confined. Exotic triples therefore do not constrain any mechanism below Λ_{meta}.

Postulates. Below Λ_{meta} only these composites and radiation exist, so composites are pair-produced from radiation. Each triple has four channels, and $t_B\bar{t}_L$ and $t_L\bar{t}_B$ are CP conjugates from a self-conjugate initial state, the minimal setting for CP violation compatible with CPT and unitarity [20]. (P1) The rung (baryonic t_B or leptonic t_L) is fixed at production and conserved by every process except annihilation back into radiation. (P2) The conjugate channels are biased by ε_t, with source ε_t times the net production rate, so the source vanishes in equilibrium. (P3) Production occurs after reheating at $T_{\text{RH}} < \Lambda_{\text{meta}}$ with the natural coupling, so a single condensate vacuum covers the observable Universe.

Structural consequences ([T1] given P1–P2). Each production event changes B and L oppositely and creates a triple with its antitriple. Hence $B = -L$, total charge zero and $\Delta N_\psi = 0$, and the late-time state is hydrogen plus two net antineutrinos per baryon.

CP source. The vacuum phase of the winding condensate changes M by two units, and $U(1)_M$ is exact in eq. (1); since the condensate itself generates the bond of J.6.3, a $U(1)_M$ rotation removes the phase exactly in every vacuum. The phase is physical only relative to an explicit breaking of strength δ, with CP-odd effects

$\propto \delta^2$. The only explicit breaking in the PH, $\kappa_{\rm CP} = (m_e/\Lambda_{\rm meta})^2$ (K.10), gives a bias $\sim 10^{-62}$. The PH as written has no CP source of adequate size.

Efficiency and open problem. With the natural coupling, η_B/ε reaches ~ 0.1 for $T_{\rm RH} \sim 4$–7×10^9 GeV, and the asymmetry is washed out above $\sim 1.5 \times 10^{10}$ GeV. The result is insensitive to gradual reheating (peak temperature up to 60 $T_{\rm RH}$). The observed η_B needs a bias of 10^{-7}–10^{-6}, far above what the PH supplies, and the sign depends on the binding law and on $T_{\rm RH}$. The reheating temperature is not a prediction.

Status ([T3]) and conditions. (C1) Derive the rung label, for instance as a winding charge changed only through the deconfined core, where the action per mode S/N_e gives a suppression $e^{-21\Delta\ell}$, enough for 13.4. (C2) Identify a derived explicit breaking of $U(1)_M$ (or another CP-odd ingredient) of strength $\delta \sim 10^{-3}$. (C3) Derive the pair-production rate of composites.

14 Dark Matter: Identity, Mass, and Relic Density

14.1 Dark Matter as Stable Z6 Singlets

State	I	Stability
(0, 0, 1, 1, 2, 2, 3, 4, 5)	0	Ground configuration (unique s-wave state: 3 forced singlets + free-spin triplet)
(0, 0, 0, 0, 0, 0, 1, 1, 4)	0	Neutral singlet (census; high shell cost)
(0, 0, 0, 0, 0, 1, 1, 1, 3)	0	Neutral singlet (census; high shell cost)

Table 8: Representative neutral N = 9 singlets (generated census, `gen_composite_tables.py` [2]); stability filter per [2], pending the parity-correct re-enumeration (P-Pauli).

Theorem DM-S (statistics by parity, [T1]). All stable dark singlets carry Q = 0 (neutral by the Z6 charge formula), with statistics fixed by body-number parity: fermionic at N $\in$ {9, 15, 21, 27}, bosonic at N $\in$ {12, 18}; scalar states are possible only at even N. DM is dark by group-theoretic necessity (Q = 0), and its stability is protected independently of spin by psion-number conservation (gauge bosons are $\psi\bar{\psi}$ bilinears with net psion number zero), Theorem KS, and — for odd N — fermion-number superselection.

Absolute stability (Theorem KS + App. G, [T1]): By App. G (exhaustive enumeration), no Z6-allowed decay channel exists. By Theorem KS (§8), ALL fragmentation decays N $\to$ {Ni} are kinematically forbidden by the cascade mass function, independently of Z6 quantum numbers (Σm(Ni)/m(N) $\geq$ 69 for any 2-body decay). Together these establish full dynamical stability: Z6 algebraic stability is both necessary and sufficient.

DM stability: 39 stable neutral singlets at N $\in$ {9, 12, 15, 18, 21, 27} by exhaustive enumeration (dynamic programming + brute force, two independent algorithms; termination proved to N = 48). Mass scale: §14.2. Relic abundance: §14.3.

14.2 Dark Matter Mass from the Landau Pole

$$\text{mDM} = 2\Lambda\text{meta}/\text{NLandau} = 2 \times 1.2609 \times 10^{12}/35 = 7.205 \times 10^{10}\ \text{GeV}. \qquad (17)$$

NLandau = 35 counts cascade modes to the Landau pole (distinct from Ne = 10 binding levels, Theorem G.1).

14.3 Relic Density Derivation

Two partially independent routes to ΩDM/Ωb; both are quoted against the observed ratio 5.364 = Ωc h^2/Ωb h^2 (Planck 2018 base chain: Ωc h^2 = 0.1200, Ωb h^2 = 0.02237, h = 0.6736).

Inference route. ΩDM = 1 − ΩΛ(PH) − Ωb(PH), with both terms computed from PH predictions and no observed cosmological density imported: ΩΛ(PH) = $\rho\Lambda$,pred/ρ_crit(h) = 2.492 × 10^{-47}/3.673 × 10^{-47} = 0.6784, and Ωb(PH) = η10,pred/(273.9 h^2) = 0.0484. Result: ΩDM/Ωb = 0.2732/0.0484 = 5.64 (5.1% error) [T2]. Declared inputs: h = 0.6736, TCMB = 2.7255 K, flatness. This route makes no assumption about the DM production mechanism.

Status (v86). The direct route below uses ξ^* = 434.09, obtained from the predicted ηB through eq. (33); both are withdrawn (§13.3). The direct route is therefore carried as unsupported until ξ^* is derived independently, and the mutual-consistency and sensitivity statements below are conditional on it. The inference route is unaffected if the measured Ωb is used.

Direct route. ΩDM/Ωb = (4π/3) × (mDM/mp) × $e^{-\xi DM}$ = 5.43 (1.2% error) [T2+], with ξDM = (mDM/Λmeta) × ξ^* = (2/35) × 434.09 = 24.805 and ξ^* = 434.09 obtained self-consistently from the predicted ηB = 6.02 × 10^{-10} via §13.3. The exponential factor $e^{-\xi DM}$ is derived from the baryogenesis sector and is proved independent of TCMB: comoving yields Y = n/s eliminate s(T) in the ratio, and Tnuc cancels in the nucleation-epoch density ratio. The prefactor 4π/3 has geometric motivation (SO(3) isotropy) and is under active derivation (P-4π/3, [T2+]).

Mutual consistency. The two routes share the baryogenesis output ηB (which sets Ωb in the inference route and ξ^* in the direct route); their discriminating second ingredients are $\rho\Lambda$ (inference) and the Landau-pole mass mDM (direct). Mutual agreement: 3.9% — fully accounted by propagation of the upstream Class-A deficits through the route elasticities (see Sensitivity below): the inference route amplifies the $\rho\Lambda$ deficit (−0.92% in ΩΛ, elasticity −2.5) and the ηB deficit (−1.75% in Ωb, elasticity −1.2), while the direct route carries only the mild $GN^{-1/2}$ dependence of mDM. The spread is error propagation, not an independent tension; no new problem is registered. Both routes are independent of be (Class A).

Sensitivity. The direct route is exponentially sensitive to the ηB convention: d ln(ΩDM/Ωb)/d ln ηB = +ξDM/3 = +8.27, so switching ξ^* from the predicted ηB (434.09) to a measured one (434.6) moves the direct value from 5.43 to 5.27 — a 2.9% swing, larger than the quoted agreement itself. The predicted-ηB convention is canonical (zero-parameter purity; §J.4). The inference route elasticities are −2.5 (wrt ΩΛ) and −1.2 (wrt Ωb). Headline percentages in this section are convention-conditional at these levels.

14.4 Experimental Signatures

Direct detection: no signal at any mass other than m_{DM} (F5a), and a rate ∼ 10^{-54} cm^2 at m_{DM} conditional on zero proto-colour coupling (F5b, [T4]); see §9.1 for the kill criterion. Monochromatic γ-line DM2 + DM2 → $\gamma\gamma$ at 72 EeV (F13; AugerPrime). UHECR excess above 10^{22} eV (F8; Auger, GRAND). XRISM (2025) null keV result consistent with PH; falsifies keV sterile-ν DM.

15 Open Problems

ID	Problem	Status	Blocks
P23	κ = 1 normalisation identity (Lemma I.1')	[T2+]	$\varepsilon\psi \to$ T1 (ηB link withdrawn, §13.3)

P14-B	Conserved baryon (rung) label; sign and size of the bifurcation bias (§13.4, §13.7)	open	ηB, nucleon stability, Theorem KS
P14-C	Baryon binding scale ($(\hbar c)/a \approx 163$ MeV) and internal structure consistent with $J{\cdot}\Sigma$ mixing and with the Gottfried sum (valence $-1/9$; sea term $+0.346$ needed, not derived); hexagonal radii are phenomenological (App. L.4.7); dynamic-ring series closed: negative on radius–mass, hybrid description demoted, per-composite geometry unsupported, minimal spin extension fails on magnetic moments (App. L.4.8)	open	hadron radii, $m_n - m_p$, sea model
P-PMNS	Sector-bridging PMNS texture (e^-: I = 3; ν: I = 0); the CC bridge is the proto-W (I = 3) — texture open; flavour basis fixed, eq. (1) alone gives $\sin^2\theta_{23} \approx 0.21$ (§17.3)	[T3]	mixing sector; F3
P-|O|	Overlap sector for absolute rates; first datum: $\|O\|^2(CF) \leq 0.370$ (2σ)	[T2]	absolute rates → T1
P-εconf	Formation-overlap democracy behind εconf = 4/9; bounded empirically at 1–2% via the +8.27 lever	[T2]	withdrawn with eq. (33) (§13.3)
P-Pauli	Pauli structure restated: auto-singlets from N $\geq$ 7; spatial shells from N $\geq$ 13 (nuclei at N = 15+); per-level structure open; 39-count re-audit (floor)	[T2]	DM spin sector
C-Q-rat	Charge rationality / integer quantization as Z6 selection principle	[T3]	why Nc = 6
F-fork	RESOLVED → twelve-state bookkeeping: the U3 decider fired ((2, $2\pi/3$)); residuals R-pair (auto-singlet + uniqueness + lepton anchor) and R-norm (bias/formation split, 0.9% seal) reviewed and held	[T2]	closed
C-web	Closed-form web RESOLVED: S definitional; $\rho\Lambda$ = C·me^6/Λmeta^2 exact rearrangement of Theorem 3.1; O(1/S) = two-loop registration	[T2+]	master invariant; F17
P26	ν-quantum number locking (ce = 11/6 vs 2)	[T2]	P23 (locking mechanism)
P-4π/3	Prefactor 4π/3 in ΩDM/Ωb direct	[T2+]	ΩDM → T1
P1	be = 325/1152 from 2PI kernel	[T2]	mass ratios Class B
P2/P12	$\Phi(\Lambda) = \Lambda$ without me; closes all adimensional ratios; 1 dimensional datum remains (irreducibility)	[T2]	me → prediction

Table 9: Twelve critical open problems. Complete registry: Appendix K.7 and psion-proofs.

16 The Psion in Hawking Radiation

The Z6 condensate modifies black hole thermodynamics when the Hawking temperature approaches Λmeta. The Planck mass MPl = ($\hbar$c/GN)^{1/2} = 1.22×10^{19} GeV is the scale at which quantum gravitational effects become order-unity.

• **Critical mass:** M* = MPl^2/(8$\pi\Lambda$meta) = 8.38 g — the black hole mass below which the horizon temperature reaches Λmeta.

• **Deconfinement radius:** rdc = (GN M/Λmeta^4)^{1/3} for M > M*.

• **Modified evaporation rate:** (dM/dt)PH/(dM/dt)SM = 127.75/106.75 = 1.197 (19.7% faster above M*).

Full derivation of M* = 8.38 g, deconfinement radius, and the 19.7% modified Hawking rate: companion PH P19 Black Holes (psion-proofs [2]).

17 Exotic Signatures and Experimental Context

17.1 ANITA: Z6 Exotic Carrier

The ANITA balloon experiment observed anomalous upward-going radio pulses inconsistent with standard neutrino interactions. The PH identifies the carrier as νS = (1, 4, 4): a colour-singlet (I = 3) composite with Q = +1/3, M $\approx$ 7.5 GeV, via:

$$\nu\mathrm{S}(1,4,4) + \mathrm{n}(0,1,5) \to \tau^{-}(0,1,5)\mathrm{DI} + \mathrm{X}(0,0,3). \qquad (18)$$

Composite identification: Appendix H (PD protocol). Mass M $\approx$ 7.5 GeV: §6 four-body mass formula. The carrier is charged (Q = +1/3) and hence not sterile; the identification is carried under E4 quarantine (outside the Table 4 census) pending confrontation with fractional-charge limits (ArgoNeuT, milliQan).

17.2 CF-CC Channel: Structural Prediction (see Appendix K)

The Chiral-Flip Charged Current (CF-CC) is a channel predicted by the Z6 composite structure:

$$\nu\mu(0,4,2) + \mathrm{n}(0,1,5) \to \mathrm{e}^{-}(0,1,2) + \mathrm{p}(0,4,5). \qquad (19)$$

Structure ([T1]): selection rules and existence. The absolute rate factorizes as RCF = NDirac g^2meta $(\mathrm{me}/\mathrm{m}\mu)^2 \times |\mathrm{O}|^2 = 6.16 \times 10^{-4}\ |\mathrm{O}|^2$, with $|\mathrm{O}|^2$ the BSE formation overlap (P-|O|); the rate is energy- and baseline-independent at every accessible order (Theorem K.8.1). MicroBooNE bounds $|\mathrm{O}|^2 \leq 0.370$ (2σ). Full treatment and the e^+e^- uniqueness theorem: Appendix K §K.8.

17.3 IceCube Tau Flavor Ratio (F3, conditional)

F3 concerns the τ–e asymmetry of the astrophysical flux at Earth, $\mathrm{A_{te}} = (\mathrm{f}_\tau - \mathrm{f_e})/(\mathrm{f}_\tau + \mathrm{f_e})$ for a pion-decay source. This quantity is fixed by the PMNS matrix alone. With standard oscillations it spans roughly -10% to +9% over the NuFIT 6.0 ranges; +7% corresponds to $\sin^2\theta_{23} \approx 0.574$ with $\delta_{CP} \approx \pi$, and with real mixing ($\delta \in \{0, \pi\}$) $\delta = 0$ gives $\mathrm{A_{te}} < 0$ for every allowed θ_{23}. The PH does not derive the PMNS matrix (open problem P-PMNS), and no derivation of the 7% figure is contained in this release; earlier versions of this section cited one in [2]. F3 is therefore carried at [T4], conditional on P-PMNS.

Status of the PMNS programme ([2], `proofs/docs/PMNS_PROGRAMME.md`). The flavour basis is fixed exactly: the generation of a neutral triple is the proto-colour of its single right-handed constituent, and the charged-current flip is colour-diagonal. With the charged leptons taken diagonal in that basis, both terms of eq. (1) together with the locking gap give $\sin^2\theta_{12}$ close to the measured value, but $\sin^2\theta_{23} \approx 0.21$ and $\mathrm{A_{te}} \approx -34\%$, the opposite sign to +7%. The charged-lepton rotation, which could change θ_{23}, requires the full P8b transfer matrix and has not been tested.

Experimental context. IceCube's flavour measurement ($\mathrm{f_e} : \mathrm{f}_\mu : \mathrm{f}_\tau = 0.30 : 0.37 : 0.33$, 5 TeV–10 PeV) does not constrain $\mathrm{A_{te}}$ at the required level. IceCube-Gen2 and KM3NeT target $\sim$5% precision on the flavour ratio, while DUNE and Hyper-K will measure θ_{23} and δ_{CP} directly, which determine $\mathrm{A_{te}}$ in any case.

17.4 Sterile Neutrino DM vs. PH

Nν = 3 (T1 theorem) forbids a fourth neutral fermionic stable composite. XRISM (2025) found no keV X-ray lines. PH and sterile-neutrino DM are mutually exclusive. Nν = 3 T1 proof: Theorem 6.1 (§6); Appendix G §G.5.2; prediction F6 (§9).

17.5 Neff and Baryogenesis

$\varepsilon\psi$ = 0.1479 generates δL = 0.370, giving $\mu\nu$/T = 0.123 and Neff = 3.054 (F12). Structurally distinct from leptogenesis: no right-handed neutrinos. The ηB link of this chain is withdrawn in v86 (§13.3); Neff is retained with the status note of §13.5. Predictions F7 and F12 (§9). Structural derivation of $\varepsilon\psi$: Appendix J §J.6.

18 Discussion

18.1 Class A vs. Class B Results

Class A (independent of b): ΩDM/Ωb, Nν = 3, Ng = 8, normal hierarchy, DM darkness. Class B (b-dependent, $\lesssim$ 35% per 1% change): mμ/me, mτ/me, Δm^2_{31}. Registered observation: Koide's empirical relation Q $\equiv$ (me + mμ + mτ)/($\sqrt{}$me + $\sqrt{}$mμ + $\sqrt{}$mτ)^2 = 2/3, satisfied at the ~10^{-5} level with no SM explanation, is listed among the b-dependent targets of the lepton sector; no PH derivation is registered.

18.2 SM Gauge Group from Z6

Z6 = Z3 × Z2 = Center(SU(3) × SU(2)). The SM gauge symmetry SU(3) × SU(2) × U(1) is derived, not postulated.

18.3 Framework Comparison: PH vs. SM and GUTs

Framework	**Free inputs**
Standard Model	∼19
Minimal GUT (SU(5))	5–8
PH (this work)	2 (GN, Newton's constant; Δm^2_{21}, neutrino-sector scale, §11)
PH (P2/P12 closed)	1 (Δm^2_{21} remains; a separate candidate closure, P26-C, is registered at T4 and not yet promoted, §11)

Table 10: Input count comparison.

18.4 What the PH Derives That the SM Assumes

Five structural facts that enter the SM as inputs or empirical accidents are theorems here: (i) electron uniqueness (Theorem 6.2) — the SM offers no reason why exactly one stable charged lepton exists; (ii) the charge–chirality identity (Theorem 6.3) — where Gell-Mann–Nishijima expresses Q through two further inputs, the PH reads Q as chirality content; (iii) Pauli-triviality of the observable sector (§6.1) — the inverse of the quark-model Δ++ story; (iv) the statistics of dark matter (Theorem DM-S) — fixed by parity, not chosen; (v) the Dirac large number — the electric-to-gravitational force ratio between two electrons, RFG = αem × ξcond^{-1} × (Λmeta/me)^2 = 4.17 × 10^{42}, and the closed relation ke/GN = Λmeta^2/K(ξ) (A.2.4): where the SM holds two independent couplings and a 42-order-of-magnitude mystery, both inverse-square laws are bookkeepings of one field — charge as chirality content (Theorem 6.3), mass as cascade binding — sharing one condensate scale. Each carries an internal falsification handle through the Z_N anchor suite ([2]: $\varepsilon\psi$(Z4) = 9/25 in the chirality sector; five Q = −1 states in the exact Z10 charge census).

19 Conclusions

The Psion Hypothesis derives the observable universe from a single fermionic field ψ with Z6 = Z3 $\times$ Z2 and spin j = 5/2, with one empirical anchor: Newton's constant GN.

• **Exact theorems ([T1]):** Nν = 3, Ng = 8, Ngen = 3; 39 stable DM states; det′/det = 1/48; σdw/N0 = 1.

• **Class A results ([T2]):** Neff = 3.054 (status note, §13.5); ΩDM/Ωb = 5.64 (inference, 5.1%); the direct route (5.43) is unsupported in v86 (§14.3). ηB: derivation withdrawn; candidate mechanism in §13.7.

• **Master invariant ([T2+]):** $\rho\Lambda GN^2$ = 1.121 $\times$ 10^{-123} predicted vs 1.132 $\times$ 10^{-123} observed (0.93%; Λmeta cancels and the prediction is GN-independent). Closed form: C·K^2·e^($-$S(1+3/π^2)) with S = Nc·Ne/be; $\hbar$-linear — the accelerated expansion is a first-order quantum effect, with co-drift falsifier $\delta\rho\Lambda/\rho\Lambda = -2\,\delta$GN/GN (F17).

• **SC input:** 1 dimensional anchor (GN) + 1 adimensional datum ($\Delta m^2_{21}/m_e^2$) $\rightarrow$ 15+ predictions. P2/P12 ([T2]): closure $\rightarrow$ all adimensional ratios internal; 1 dimensional datum remains (Remark 5.1).

• **Hard T1 falsifiers:** any fourth charged lepton, fourth active neutrino, or ninth gluon refutes the framework.

• **Structural theorems ([T1]):** unique electron; charge–chirality identity Q = nR $-$ 1; DM statistics by parity (DM-S); Pauli-trivial observable sector; CF-CC flatness with $|O|^2 \leq 0.370$ (2σ).

• **Gravity-sector consistency:** the condensation channel is a Lorentz scalar (K.9; c_gw = c_γ by structure) and the Z6 relic-wall problem resolves by infrared tension and exact chiral bias (K.10).

• **Critical open step (P23):** the normalisation identity $\delta\chi$ = Ind(Dχ)/$(2j)^2$ — one equation from T1 closure of $\varepsilon\psi$ (its link to ηB is withdrawn, §13.3). **New open step (P14-B):** a conserved baryon (rung) label, required by nucleon stability and by the candidate baryogenesis mechanism (§13.4, §13.7).

A Complete Inventory of PH Results

This appendix catalogues all results by category with T-level classification. The framework derives everything from ψ + Z6 + NJL fixed point + GN ([SC]); nothing beyond GN is a free parameter.

A.1 Derived Fundamental Parameters

be = 325/1152 $\approx$ 0.2821 (cascade parameter, T1 given the 2PI kernel). Ne = 10 (Z6 theorem, T1). S = 6Ne/be = 212.677 (condensate action, T1). K(ξ) = 1.16335 (Gel'fand–Yaglom coefficient, T1). fπ = $\sqrt{(3/\pi)}$ me = 0.4994 MeV (condensate decay constant; confirmed by three independent checks: Yukawa coupling, vortex-action consistency, GY $\sqrt{\xi}$). Λmeta = $\sqrt{(K(\xi)/GN)}$ exp($-S/2g^2$) ([SC] $\rightarrow$ [T2], eq. 6).

A.2 Exact Theorems (T1)

A.2.1 Counting Theorems

Nν = 3 active neutrino species (Z6 singlet counting, [T1]). Ng = 8 gluons (adjoint of the emergent colour group, [T1]). Ne = 10 single-particle modes (Z6 combinatorial theorem, [T1]). Ngen = 3 charged-lepton generations: the fourth would require JR = 7/2 > jmax = 5/2 and is kinematically forbidden within Z6 ([T1]). This is a kinematic upper bound, not a constructive identification of the second and third generations; the latter is addressed separately via the Leptonic Transfer Matrix (P8b, App. A.2.5; Appendix L.4.4). SM gauge group SU(3) $\times$ SU(2) $\times$ U(1) is the unique continuous gauge group consistent with the Z6 centre ([T1]).

A.2.2 Algebraic Identities

σdw/N0 = 1 (domain-wall tension over zero-mode normalisation; one-line algebra, [T1]). GY ratio = 1/48 (Gel'fand–Yaglom domain-wall ratio; three independent verifications, [T1]). ζ'4D(0) = $-3/(2\pi)$ (Kramers–Minakshisundaram spectral zeta, [T1]). Z3 overcounting = 27 (fixed-point Z3 degeneracy, [T1]). λ = 9/2 in V(σ) = $\lambda(\sigma^2 - f\pi^2)^2$: forced by mZ6 = 6f$\pi$ (Theorem W.1: $m^2 = 8\lambda f\pi^2$, imposing mZ6 = 6fπ forces λ = 9/2 exactly, [T1]). k $-$ 2j = (2j + 1) $-$ j for j = 5/2, k = 3: unique self-consistency condition of the leptonic sector ([T1]). $1/r^2$ for both Coulomb and Newton (Gauss's theorem in 3D from masslessness, [T1]).

A.2.3 Z6 Gauge Structure Theorems

V.1: Qk+3 = Qk + 1; unit Z3 triality; ΣQk = 1 structural ([T1]). V.2: the flip 2 $\rightarrow$ 3 conserves triality; intermediaries are colour singlets ([T1]). V.3: Eflip must be rational, killing one-meta-boson exchange ([T1]). V.4: the charge interlock is a flux-quantum count, 2j quanta of size 1/N ([T1]). V.5: Z6 = Z3 $\times$ Z2 factorisation removes the BKT caveat, restoring sharp self-duality ([T1]). V.7: the equivalence web — five reformulations of Lemma I.1' are exactly equivalent; no further reformulation can yield T1 ([T1]).

A.2.4 The ν Quantum Number

$\nu \equiv \lfloor k/3 \rfloor \oplus \chi$ is a hidden good quantum number of the Z6 structure. Theorem ν.1: $\nu = 0 \Leftrightarrow$ locked subspace, verified exactly on all 12 single-particle states ([T1]). Theorem ν.2: G $\cdot$ Punlocked = G $\cdot$ Nν with Nν = $-(1 - \hat{Z}2 \cdot \chi)$ — the deep-gap projection operator of the I.6 algebraic model is the ν charge, exact ([T1]).

A.2.5 Leptonic Transfer Matrix (P8b)

t12 = -8 (J^+ element, exact, [T1]). $\langle 3|Hdd|1\rangle = t12^2 = 8$ (three-body symmetry cancellation, exact, [T1]). 15 active pairs (direct count). $k - 2j = (2j + 1) - j$ identity. Algebraic cancellation of be in t13 ([T1]). C2 diagonals = {10.75, 8.75, 6.75} (exhaustive Q = -1 enumeration, [T1]). Mass ratios $m\mu$/me and $m\tau$/me: $\lambda i/\lambda j$ vary < 0.01% for $\pm 28\%$ change in be (insensitivity theorem, repository [2] §I.9.3, [T1]). CF-CC structure (selection rules, existence): [T1]; absolute rate $6.16 \times 10^{-4} \times |O|^2$, ceiling $|O|^2 \leq 0.370$ (2σ) ([T2], P-|O|). Absolute DM stability: four-body Z6 singlets have no kinematically accessible decay channel (exhaustive enumeration, termination proved to N = 48, [T1]).

A.2.6 Gravitational Structure

GN = $K(\xi)\ \Lambda meta^{-2} \exp(-S/g^2)$ (T1 at leading order). PC1 ([T2], resolved): hAB is a massive spin-2 composite ($\sim$0.6 GeV), satisfying the Weinberg–Witten theorem by being massive; GN emerges via the Sakharov mechanism [3] as a collective condensate property; there is no massless composite graviton in the PH. The $1/r^2$ law for gravity follows from masslessness of the effective long-range coupling and 3D isotropy ([T1]).

A.3 Structural Results (T2)

A.3.1 Mass Spectrum (Class B — b-dependent)

me (cascade output) = $\Lambda meta \exp(-Ne/be)$ = 0.5089 MeV (0.42% error; the observed 0.511 MeV is the matrix anchor in P8b). $m\mu$/me: +0.46%. $m\tau$/me: +0.56%. Zero free parameters. mp/$m\tau$ = 0.52805 at 0.000% error (structural, b-independent). GN = 6.653×10^{-39} GeV^{-2} at 0.83% error against CODATA 2018 in natural units (6.70883×10^{-39} [11]).

A.3.2 Cosmological Results (Class A — b-independent)

ηB: derivation withdrawn in v86 (§13.3); level bifurcation is a candidate mechanism conditional on P14-B (§13.7, [T3]). Neff = 3.054 ($< 1\sigma$ from CMB; CMB-S4 target). $\Omega DM/\Omega b$ = 5.43 (1.2% error, direct, [T2+]) / 5.64 (5.1% error, inference, [T2]); mutual 3.9% (error propagation, §14.3). $\rho\Lambda = 2.492 \times 10^{-47}$ GeV^4 (0.93% error). $\rho\Lambda GN^2 = 1.121 \times 10^{-123}$ predicted vs 1.132×10^{-123} observed (0.93%; the prediction is GN-independent).

A.3.3 Force and Coupling Results

ke/GN = $\Lambda meta^2/K(\xi)$ at 0.000% error (P22, [T2]). CZ6 = $(2j + 1)^2 \times 2 = 72$ (Z6-exact integer; observed ratio $\alpha meta/\alpha em$ = 71.75 differs by 0.35% = predicted running of αem). RFG = Fem/Fgrav = $\alpha em \times \xi cond^{-1} \times (\Lambda meta/me)^2 = 4.17 \times 10^{42}$ (three PH factors, no free parameters). The Coulomb and Newton forces share common Z6 origin: both electric charge and gravitational mass are quantum numbers of psion composites built from the same field ψ.

A.3.4 Condensate Structure

C = Nc/(27π^4) (dark-energy coefficient, [T2+]). fπ = $\sqrt{}$(3/π) me = 0.4994 MeV (Branch B; confirmed by three independent internal checks: Yukawa coupling, vortex-action consistency, Gel'fand–Yaglom ξ). ucond = fπ^4 = (3/π)^2 me^4 $\approx$ 0.912 me^4 (Branch B; Conjecture CEC). Self-consistency triangle: GN, $\rho\Lambda$, and me are three projections of the single action S = 6Ne/be through the coupling g^2 = 2π^2/3.

A.3.5 Locking Mechanism (ν-channel, P26)

The gap operator G · Punlocked acquires a derived dynamical origin: f1 = 0 (the direct J$\Sigma \to \nu$ route is killed by kinematic identity, exact). f2 = 1 (universal scalar piece). f3 = 3/4 (tensor-induced ν self-energy, exact rational). f4 = 1/12 (bubble). Mechanism: tensor-induced ν condensation in two stages — ΔT (tensor condensate) $\to \Delta\nu$ (induced ν gap) through the 3/4 coefficient. Quantitative open step: ceff = f2 + f3 · w3 + f4 = 11/6 with independently registered multiplicities, vs target 2 required by the operational definition; fork R1/R2 registered (P26, [T2]).

A.4 Ontological Derivations

Particles are Z6 tuples (k1, …, kN); antiparticles are the same tuples made of antipsions ($N_\psi = -3$, charge $-Q$), whose Z6 labels are the inversions k $\to$ (6 − k) mod 6 (v86 correction; §13.1). Dark matter: absolutely stable Z6-singlet composites (Q = 0, I $\in \{0, 3\}$; statistics by body-number parity, Theorem DM-S); stability proved by exhaustive enumeration (39 states at N $\in \{9, \ldots, 27\}$, termination theorem, [T1]). Mass scale mDM = 7.205 $\times$ 10^{10} GeV ([T2]). The full composite spectrum contains 76,761 states; the SM is the 8-state shadow under the projections Q $\in \mathbb{Z}$, N $\leq$ 3, stable ground states.

A.5 New Forces Predicted

Force 1 — Proto-colour force ([T3]): mediated by the tower of Z6 proto-colour bilinears. The observable member is the proto-colour singlet of the four neutral bilinears ((1,2)$\otimes$(5,4): I = 0, Q = 0, net psion number zero — the exact analogue of hadrons as colour singlets); individual bilinears carry I $\in \{1, 5\}$ and are confined. The lightest observable member is identified with the X17 singlet (F15): mX17 = Λmeta exp(−9/be) = 17.62 MeV (observed 17.01 $\pm$ 0.16 MeV, 3.59% error, Atomki [8]). Universal coupling across three nuclei follows from Z6 group structure. The tower predicts heavier members at each cascade level N < 9.

Force 2 — Chiral-Flip force ([T1]): a new interaction exchanging chirality within the Z6 internal space ($\Delta\chi$ = 1, ΔQ = 0, ΔI = 0), absent from SU(3) $\times$ SU(2) $\times$ U(1). Structure [T1]; absolute rate 6.16 $\times$ 10^{-4} $\times$ |O|^2 with |O|^2 $\leq$ 0.370 (2σ, MicroBooNE exclusive 1e1p [14]) — see App. K §K.8.

Force 3 candidate — resonant gravitational coupling ([T4], P18): oscillatory δGN/GN $\sim$ 10^{-10} at fdrive $\in$ [0.38, 0.43] GHz (UHF). Mechanism conjectural ([T4]): requires Z6 composite gauge structure (PC1-sub, open) and resolution of the static-vs-dynamic screening inconsistency (PC2, frozen). Not falsifiable until both close.

A.6 Experimental Anomalies Illuminated

CF-CC channel (structural [T1] prediction): RCF = 6.16 $\times$ 10^{-4}; MiniBooNE 4.8σ excess was consistent, but MicroBooNE (2022) found no excess — experimental status disputed (see App. K §K.8). X17 anomalous

boson (Atomki, three nuclei): mX17 = 17.62 MeV (F15, [T3], disputed†). ANITA anomalous upward-going pulses: candidate carrier νS = (1, 4, 4), a Z6 composite with I = 3, Q = +1/3, M $\approx$ 7.5 GeV, derived from Z6 selection rules independently of the experimental result.

A.7 Falsifiable Predictions (16 in Table 4)

Hard T1 falsifiers (immediate): fourth charged lepton, fourth active neutrino, ninth gluon. Near-term: F2 (δme/me = $-\frac{1}{2}$ δGN/GN); F12 (Neff = 3.054, CMB-S4). Existing data: F15 (X17, Atomki, disputed); F5a/F5b (direct detection: exclusion outside m_{DM}, E2; rate at m_{DM}, [T4]; XENONnT/LZ). See Table 4 for the complete registry.

A.8 Principal Open Problems

P23 ([T2+], critical path): Lemma I.1', $\delta\chi$ = Ind(Dχ)/(2j)^2; closure cascades to $\varepsilon\psi$ = 25/169 (T1); the former $\varepsilon\psi \to \eta$B link is withdrawn (§13.3). P14-B (open; critical for baryogenesis and nucleon stability): a conserved baryon (rung) label (§13.4, §13.7). Two open components: the unlocked-gap mechanism (P26; mechanism at T2, quantitative closure open) and the winding placement with Ind = 1 in the §2.3 construction. P2/P12 ([T2]): self-consistent fixed-point $\Phi(\Lambda) = \Lambda$, determining Λmeta from internal consistency; closure makes all adimensional ratios internal; exactly one dimensional datum remains, by the irreducibility established in §5 (a scale-free kernel cannot fix a scale). P1 ([T2]): formal derivation of be = 325/1152 from the 2PI kernel (28% theoretical gap; proven not to propagate to mass ratios; repository [2] §I.9.3). P8b Passo A.3 ([T2]): formal integration of Σ2 = be $\times$ 15/8. PC1-sub ([T3]): minimum energy scale to perturb GN via collective modes.

B Derivation of the Kink Fluctuation Operator

The second derivative of the kink potential V(σ) = $\lambda(\sigma^2 - f\pi^2)^2$ evaluated on the kink solution σk(x) = fπ tanh(mZ6 x/2) gives:

$$V''(\sigma k) = mZ6^2 - (3/2)\, mZ6^2\, \mathrm{sech}^2(mZ6\, x/2). \qquad (20)$$

Substituting y = mZ6 x/2 yields the fluctuation operator:

$$M = (mZ6^2/4)\, [\, -d^2/dy^2 + 4 - 6\, \mathrm{sech}^2(y)\,],\ n(n+1) = 6,\ n = 2. \qquad (21)$$

This is the n = 2 Pöschl–Teller Hamiltonian — a quantum-mechanical potential well exactly solvable in closed form, with two bound states and a reflectionless continuum.

The two bound states have eigenvalues ε0 = 0 (zero mode / Goldstone) and ε1 = 3mZ6^2/4 (shape mode). The zero mode corresponds to translation of the kink along x and must be excluded from the functional determinant; the shape mode contributes the lowest eigenvalue of the fluctuation operator, entering the Gel'fand–Yaglom calculation of Appendix D.

C Exact Identities Used in Theorem 3.1

The coefficient C = Nc/(27π^4) rests on three exact identities.

Domain-wall tension. σdw = 4fπ^3. This follows from integrating the kink solution σk(x) = fπ tanh(mZ6 x/2) through the potential: σdw = $\int$[V(σk)]^{1/2} dσk = 4fπ^3.

Zero-mode normalisation. N0 = (2/3) mZ6 fπ^2. The zero-mode wave function ψ0(x) $\propto$ sech^2(mZ6 x/2) integrates to N0 = (2/3) mZ6 fπ^2.

Ratio (exact, T1).

$$\sigma\text{dw/N0} = 4f\pi^3 \,/\, [(2/3) \times 6f\pi \times f\pi^2] = 4f\pi^3/4f\pi^3 = 1. \checkmark \text{ [T1]} \qquad (22)$$

The Z3 Faddeev–Popov factor. The Z3 fixed-point degeneracy contributes an overcounting factor of 27, giving C = Nc/(27π^4) with Nc = 6.

D Theorem P16: Renormalisation of the Z6 Wall Determinant

This appendix proves Theorem P16 (T1): the explicit renormalisation equation σwall_ren = σwall_bare + $\delta\sigma$bulk for the Z6 domain wall. The proof uses the exact heat-kernel trace ΔK(t) = erf(m$\sqrt{}$t) + e$^{-3m^2t/4}$ erf(m$\sqrt{}$t/2) (Lemma I), Seeley–DeWitt locality, analytic pole cancellation in Mellin space, and the exact finite part Γren/A = $-\sqrt{}$3 mZ6^3/(96π).

D.1 The Kirsten–McKane Spectral Zeta Result

The Kirsten–McKane formula [6] for the operator L = $-\Delta$3 + A(x) in d = 3 transverse dimensions:

$$\zeta'\text{l}(0) = (4\pi)^{-3/2} \int_0\text{\^}\infty \text{ dt } t^{-5/2} \text{ Kdiff(t)}, \qquad (23)$$

where Kdiff(t) = e$^{-\varepsilon 1\, t}$ and ε1 = 3mZ6^2/4. Evaluating the Laplace transform:

$$\zeta'\text{4D,diff}(0) = -3/(2\pi). \text{ [exact, [T1]]} \qquad (24)$$

D.2 The Gel'fand–Yaglom Result

The GY method [6] gives the fully renormalised determinant ratio (including bulk subtraction):

$$\det'(\text{Mk})/\det(\text{Mvac}) = 1/48. \text{ [T1], exact.} \qquad (25)$$

Three independent verifications: (i) analytic zeta function of exact ΔK(t) — $\zeta'(0)$ = ln 12; (ii) lattice spectral pairing (L = 120, N = 32000): 0.08333461 vs 1/12 = 0.08333333; (iii) Jost threshold limit: [Π(n $-$ j)/(n + j)]/(2n)^2 = 1/48.

$$\Gamma\text{GY} = (1/2)\log(1/48) = -(3/2)\log 2 - (1/2)\log 4 = -1.93574\ldots \text{ [T1]} \qquad (26)$$

D.3 Theorem P16: Renormalisation Equation ([T2+])

Theorem D.1 (Scheme-equivalence of domain-wall determinant, [T2+]). *The fluctuation determinant of the kink background admits two renormalisation-scheme representatives:*

- **GY scheme:** $\det'$(Mk)/det(Mvac) = 1/48 (y-units; 1/12 unit-gap).
- **KM scheme:** ζ'4D,diff(0) = $-3/(2\pi)$.

The mismatch defines the bulk-subtraction cocycle: Γbulk−ct = ΓGY − ΓKM = −(3/2) log 2 + 3/(4π) = −0.901889… This cocycle is a scheme-transformation artefact absorbed by the domain-wall renormalisation condition. By the standard renormalisation principle, physical observables are scheme-independent: det_ren^ζ = det_ren^GY. Therefore C = Nc/(27π^4) is scheme-independent.

Consequence ([T2+]): Theorem P16 closes the renormalisation equation at T1. The prediction ρΛ = C Λmeta^4 e^{-S} = 2.492 × 10^{-47} GeV^4 (0.93% error) is [T2+]: the renormalisation equation is proved (T1), and the C = Nc/(27π^4) assembly retains [T2+] status pending the identification μ4D = 1/π^4.

The physical coefficient C is constructed from the GY-renormalised wall partition function (Steps 1–4 of Theorem 3.1). The KM result ζ′4D,diff(0) = −3/(2π) is not an additional multiplicative factor in C — it is the heat-kernel representative of the same determinant. The cocycle Γbulk−ct is absorbed by the domain-wall renormalisation condition and does not appear in physical observables.

Factor decomposition: C = Nc × |Z3|$^{-3}$ × μ4D with Nc = 6, |Z3|$^{-3}$ = 1/27, μ4D = 1/π^4 holds independently of regularisation scheme. The 1/π^4 origin: the Faddeev–Popov path-integral measure over S^1-valued transverse fluctuations contributes a factor π^{-4} in the 4D functional integral (see P16, [T2]).

E Unit Conversion: ρΛ in GeV^4

$$\rho\Lambda_obs = \Omega\Lambda \cdot (3H0^2c^2)/(8\pi GN) \cdot (\hbar c)^3/(1\ \text{GeV}) = 2.515 \times 10^{-47}\ \text{GeV}^4. \qquad (27)$$

Using ħc = 0.197327 GeV·fm, H0 = 67.36 km/s/Mpc, ΩΛ = 0.6847 (Planck 2018 TT,TE,EE+lowE+lensing [1]); equivalently ρ_crit = 3.673 × 10^{-47} GeV^4.

Convention note: every observational comparator in this paper (ρΛ,obs; ρΛGN^2; ΩDM/Ωb = 5.364) is drawn from this single chain — do not mix Planck chains.

F Principal Conclusions and Spectrum Analysis

F.1 αmeta = π/6: Formal Proof

Theorem F.1 (αmeta = π/Nf, [T1] conditional). *The Molien series for Z3 acting on $\mathbb{C}^6$: F(x) = (1/3)[1/(1 − x)^6 + 2/(1 − x^3)^2]. The identity gmeta = 2π/√Nf gives αmeta = g^2/(4π) = π/Nf = π/6.* □

F.2 PH Composite Spectrum: Molien Series

The generating function F(x) counts Z6-invariant composites at each body number N. The coefficient of x^N gives the total number of colour-singlet states.

N	C(N) total	Cstable	Cunstable	% stable	Cum. stable
1	2	2	0	100%	2
2	7	6	1	86%	8
3	20	14	6	70%	22
4	42	20	22	48%	42
5	84	28	56	33%	70
6	156	29	127	19%	99
7	252	30	222	12%	129
8	462	30	432	6%	159
9	670	35	635	5%	194

10	1001	30	971	3%	224
12	2066	30	2036	1%	284
Σ (1–20)	76761	534	76227	0.7%	534

Table 11: PH composite spectrum. Cstable(N) ∈ {29, 30, 31} for all N ≥ 10 (Molien saturation, falsifiable prediction F14, [T3]).

N = 9 (Cstable = 35) is the final pre-saturation transition level; Cstable(N) ∈ {29, 30, 31} applies strictly for N ≥ 10. Any value Cstable(N) ≥ 32 falsifies the saturation mechanism (prediction F14, [T3]; testable by running `enumerate_z6.py`).

F.3 What the Standard Model Represents

The SM is the triple projection of the PH spectrum onto: (1) Q ∈ ℤ; (2) N ≤ 3; (3) stable ground states. This selects 8 states from 76,761.

F.4 PH Boson Spectrum: 36 Bilinears

All spin-1 force carriers are bilinears B^μ_kk′ = $\bar{\psi}$k γ^μ ψk′ with Z6 gauge charge Ibln = (k′ − k) mod 6.

Family	**Count**	**Q**	**Mass**	**Force**
Photon (diagonal)	1	0	massless	EM (✓)
Proto-W±	6	±1	MW	Weak (✓)
Z^0 (mixing)	—	0	MZ	Weak (✓)
Gluons (Z3)	8	0	massless	Strong (✓)
X17 family	4	0	17.6 MeV	Proto-colour
Exotic bilinears	20	±n/9	cascade	Z6 sector

Table 12: PH boson spectrum. Two forces beyond the SM are predicted: the proto-colour force (X17 family, scale Λmeta) and Z6 Sector Isolation (topological).

G PH Proofs Repository

All numerical verifications, enumeration algorithms, and derivation scripts are maintained in psion-proofs (GitHub: `https://github.com/karlosmardenmaia/psion-proofs` [2]).

Module	**Section**	**Content**
`proofs/P_dark_matter.py`	§14.1	39 stable Z6 dark singlets ([T1])
`proofs/P_dark_energy.py`	§5.2	C = Nc/(27π^4) functional det ([T1])
`proofs/P_eps_psi.py`	§13.2, App. J	$\varepsilon\psi$ = 25/169 chain ([T2])
`proofs/P_counting_theorems.py`	§7–§9	Nν = 3, Ng = 8, Ngen = 3 ([T1])
`proofs/P_wall_tensions.py`	§5.2, App. B–C	σdw/N0 = 1, det ratio 1/48 ([T1])
`proofs/P_yield_ratio.py`	former §13.3–13.4	ηB derivation (withdrawn in v86)
`p14d_psion_number_audit/`	§13	P14 audit; level bifurcation ([T3])
`scripts/cascade_anchor.py`	§3	GN SC input; me cascade output
`scripts/dm_prova_real.py`	§14.2	DM mass verification ([T2])
`scripts/proof_T_CMB_cancels.py`	§14.3	TCMB-independence ([T1])
`scripts/cascade_level_anchors.py`	§14.2	Cascade level → body number ([T1])
`docs/DARK_MATTER_PROOFS.md`	§14	Full DM proof inventory
`docs/PROGRESS_AND_ROADMAP.md`	App. J	P23 derivation log

Table 13: Core modules of the psion-proofs repository.

All proofs run against a pytest suite ($\geq$ 25 tests, all passing): pip install -r `requirements.txt` && pytest tests/. Reproducible on any machine with Python 3.10+.

Theorem G.1 (Ne = 10, [T1]). *Ne = Nγ + Ng + Nφ = 1 + 8 + 1 = 10.*

Proof. Nγ = 1: unique Z6-singlet spin-1 massless mode (proved in §7). Ng = 8: $|Z3|^2 - 1 = 8$ coloured gluonic bilinears (proved T1 in §7). Nφ = 1: unique Z6-singlet scalar mode. Total: 1 + 8 + 1 = 10. This is a counting theorem of Z6 = Z3 × Z2; no free parameter. □

Falsifiable prediction F14 ([T3]). For all N $\geq$ 10, Cstable(N) $\in$ {29, 30, 31}. Any value Cstable(N) $\geq$ 32 falsifies the saturation mechanism. Testable by running `enumerate_z6.py` for N = 7, 8, 11, 12.

DM stability theorem ([T1]). 39 stable neutral singlets at N $\in$ {9, 12, 15, 18, 21, 27} by exhaustive enumeration using two independent algorithms (dynamic programming + brute force); termination proved to N = 48. Code: `proofs/P_dark_matter.py`.

H The Psion Diagram: Complete Calculus for Composite Classification

This appendix makes the PH composite spectrum self-contained. It provides: (H.1) the six psion states and their quantum numbers; (H.2) composite formation rules; (H.3) the five-step Psion Diagram (PD) protocol; (H.4) the complete N = 1–4 spectrum; (H.5) the Molien series and proof of Ne = 10; (H.6) exactly forbidden channels; (H.7) the proto-quark sector; (H.8) the αmeta = π/6 formal proof; (H.9) worked examples. No external reference is required to verify these results.

H.1 Composite Formation Rules

Rule 1 (Z6 colour singlet, [T1]): a composite is colour-singlet (observable) if and only if I = (Σki) mod 6 $\in$ {0, 3}.

Rule 2 (Electric charge): Q = ΣQki. Observable particles have Q $\in \mathbb{Z}$ or Q $\in$ {±1/3, ±2/3} (fractional, confined below Λmeta).

Rule 3 (Spin): total spin J follows from coupling individual mJ(ki) values under SU(2)meta. Jmax = N × 5/2.

Stability criterion: a composite is stable if it is the lightest state in its (I, Q, J) sector.

H.2 The Five-Step Psion Diagram (PD) Protocol

Step 1: Assign Z6 triples. Compute Q = ΣQki, I = (Σki) mod 6, chirality. For unknown particles: enumerate all Z6-allowed triples consistent with observed Q and J.

Step 2: Apply conservation laws at every vertex. (a) Qin = Qout. (b) Iin = Iout mod 6. Sector Isolation Theorem: I = 0 and I = 3 sectors never mix perturbatively.

Step 3: Classify diagram topology. L = 0 (tree): rate $\sim g^2$meta. L = 1 (one loop): rate $\sim g^4$meta. CF-CC is the unique inter-composite channel exchanging a single constituent between a three-body lepton and a three-body baryon.

Step 4: Determine amplitude type. O1 (scalar): |ΔmJ| > 1. O2 (spin): |ΔmJ| = 1. Tree-level chiral-flip rate: R $\sim$ NDirac × g^2meta × (mass ratio)^2.

Step 5: Predict observables. Conservation laws give selection rules. Rate (leading order): BSE overlap integral required for precision. Mass: m $\sim$ Λmeta $\times$ exp(−Ncascade/b).

H.3 Complete Composite Spectrum: N = 1 to N = 4

N = 1. Only k = 0 (I = 0) and k = 3 (I = 3) are colour-singlets, but both are unstable. No observable N = 1 particles.

N = 2 (bilinears).

Representative state	I	Q	J	SM identity	Status
(k, k′), I = 0, Q = 0, J = 1	0	0	1	Photon γ	[T1]
$\|Z3\|^2 - 1 = 8$ coloured states	0	±frac	0/1	Gluons (8)	[T1], Ng = 8
(k, k′), J = 2, I = 0, Q = 0	0	0	2	hAB $\sim$ 0.6 GeV	[T2], not graviton
Others $\notin \{0, 3\}$	—	—	—	Confined	—

N = 3 (leptons and nucleons).

Configuration	I	Q	ΣmJ	Identity
LLL (0, 1, 2)	3	-1	-9/2	e^- (unique Q = −1, Theorem 6.2)
LLR (0, 1, 5)	0	0	-3/2	n / $\nu\tau$ (dual-sector)
LLR (0, 2, 4)	0	0	-3/2	$\nu\mu$ / Σ^0-like (dual-sector)
LRR (0, 4, 5)	3	+1	3/2	Proton p (charge check: 1/9 + 4/9 + 4/9 = +1)
LLR (1, 2, 3)	0	0	-3/2	νe / Λ^0-like (dual-sector; generation ordering open)
LRR (1, 3, 5)	3	+1	3/2	Σ^+-like
LRR (2, 3, 4)	3	+1	3/2	Σ^+-like (p-alternative configuration)
RRR (3, 4, 5)	0	+2	9/2	Q = +2 exotic (BSM)

N $\geq$ 4 (dark matter: neutral singlets).

Configuration	I	Status
(0, 0, 1, 1, 2, 2, 3, 4, 5)	0	Ground configuration (closed shell + valence) — stable, neutral
(0, 0, 0, 0, 0, 0, 1, 1, 4)	0	Neutral singlet (census; high shell cost)
(0, 0, 0, 0, 0, 1, 1, 1, 3)	0	Neutral singlet (census; high shell cost)
All others	—	Confined

H.4 Exactly Forbidden Channels (T1)

Proposed transition	Reason forbidden	Experimental test
Flavour-changing NC vertex ($\nu\mu \to \nu$e at one vertex)	I = 0 and I = 3 sectors never mix at a perturbative vertex; mass-eigenstate oscillation is outside the theorem's scope (P-PMNS)	FCNC suppression
$\nu\mu \to \nu$S (exotic)	Single-carrier vertex needs ΔI = 3 with a neutral mediator; the only I = 3 bilinears (proto-W) carry Q = ±1	No mixing; XRISM null (2025)
4th neutrino	Nν = 3 is T1 theorem; no 4th Q = 0, I $\in \{0, 3\}$ three-body singlet	Neff < 4; LEP/FCC null
DM $\to \gamma$ direct decay	Net psion number (DM: N $\geq$ 9) vs γ (0); no lighter same-N state (ground-state definition); odd N additionally fermion-protected (DM-S)	No keV γ line; XRISM 2025
FCNC (u $\to$ c, d $\to$ s)	Each transition changes I; Sector Isolation forbids single-vertex FCNC	GIM mechanism; FCNC suppressed
Any Qin $\neq$ Qout	Electric charge exactly conserved at every vertex	Universal

Table 14: Exactly forbidden transitions and their group-theoretic reasons.

H.5 Worked Examples: PD Protocol Applied

Example 1 — CF-CC reaction (structure, [T1]; experimental status disputed).

$$\nu\mu(0, 4, 2) + \mathrm{n}(0, 1, 5): \mathrm{Iin} = 12 \bmod 6 = 0, \mathrm{Qin} = 0. \qquad (28)$$

$$\mathrm{e}^{-}(0, 1, 2) + \mathrm{p}(0, 4, 5): \mathrm{Iout} = 12 \bmod 6 = 0 \checkmark, \mathrm{Qout} = 0 \checkmark. \qquad (29)$$

RCF = NDirac g^2meta $(\mathrm{me}/\mathrm{m}\mu)^2 = 4 \times (2\pi^2/3) \times (0.511/105.66)^2 = 6.16 \times 10^{-4}$. Structural prediction: RCF $= 6.16 \times 10^{-4} \times |O|^2$ (structure T1; overlap open, P-|O|). MicroBooNE bounds $|O|^2 \leq 0.370$ at 2σ (see App. K §K.8).

Example 2 — Photon identification ([T1]). Two-body (k, k′) with I = 0, Q = 0, J = 1: unique solution is diagonal bilinear B^μ_kk = $\bar{\psi}$k γ^μ ψk, I = 0, Q = 0, J = 1, massless. ⇒ Photon. ✓

Example 3 — Dark matter stability ([T2]). Composite (0, 1, 2, 3): I = 0, Q = 0, J = 0. Decay DM → νe + X? X must have J = 1/2, Q = 0, I ∈ {0, 3}. No such lighter composite exists ⇒ DM absolutely stable. ✓

I The Proto-Quark: PH Hadronic Constituents and Their Relation to SM Quarks

This appendix introduces the term proto-quark for the PH hadronic constituent and develops the formal argument distinguishing it from the SM quark: (I.1) why a new name is warranted; (I.2) three exact structural distinctions; (I.3) what the two frameworks agree on; (I.4) falsifiable predictions; (I.5) responses to expected objections; (I.6) summary table.

I.1 Why a New Name is Warranted

The SM quark is defined operationally: charge Q ∈ {+2/3, −1/3}, colour charge under SU(3), spin 1/2, and confinement. The PH does not dispute these observations. What it disputes is the ontological status: the SM treats the quark as fundamental. The PH derives the same observable object as a composite of psions.

Definition I.1 (Proto-quark). *A proto-quark is an N = 3 psion composite in a colour-charged Z6 sector (I ∉ {0, 3}), permanently confined below Λmeta. Observable SM charges ±2/3, ±1/3 emerge as the low-Q^2 effective description at $Q^2 \ll \Lambda\mathrm{meta}^2$.*

I.2 Three Exact Structural Distinctions

Confinement: arithmetic vs. dynamical. In QCD, confinement is dynamical (non-Abelian SU(3) running coupling). In the PH, confinement is arithmetic: I ∉ {0, 3} ([T1]). No confinement problem exists in the PH.

Fractional charge: axiomatic vs. derived. In the SM, Q = +2/3 and Q = −1/3 are inputs. In the PH, all charges are derived from the Qk formula (§1).

Asymptotic freedom: present vs. absent. QCD: αs(Q^2) decreases logarithmically. The PH: no running coupling between proto-quarks below Λmeta — proto-quarks are permanently confined for all $Q^2 < \Lambda\mathrm{meta}^2$.

I.3 What the Two Frameworks Agree On

Observable	**SM QCD**	**PH Proto-Quark**	**Agreement**
Proton charge Q(p) = +1	+2/3 + 2/3 − 1/3	(0, 4, 5): +1/9 + 4/9 + 4/9 = +1	Exact
Neutron charge Q(n) = 0	+2/3 − 1/3 − 1/3	(0, 1, 5): +1/9 − 5/9 + 4/9 = 0	Exact
Ng gluons	8 (SU(3) adjoint)	$\|Z3\|^2 - 1 = 8$ ([T1])	Exact
Quark confinement	Always confined	I $\notin \{0, 3\}$: always confined	Exact
mp/mτ ratio	0.52805	exp(−ΔXD/b) = 0.52805	0.000% ([T2])
Integer hadron charges	By construction	By Z6 singlet rule	Exact

Every measurement on hadrons to date is consistent with both frameworks. The PH reduces to the SM at $Q^2 \ll \Lambda meta^2$.

I.4 Falsifiable Predictions

CF-CC inter-composite scattering (structure [T1]; App. K §K.8). RCF = NDirac g^2meta $(me/m\mu)^2 \times |O|^2 = 6.16 \times 10^{-4}\ |O|^2$, flat in energy and baseline (Theorem K.8.1); MicroBooNE bounds $|O|^2 \leq 0.370$ at 2σ [14], and SBND's percent-level νe-like rate measurement tightens the ceiling directly.

Deep inelastic scattering at $Q^2 \sim \Lambda meta^2$. At $Q^2 \ll \Lambda meta^2$, PH and QCD predictions are identical. At $Q^2 \sim \Lambda meta^2 \sim 10^{24}$ GeV^2, individual proto-quark charges become resolvable and F2(x, Q^2) deviates from QCD. Prediction F4 ([T3]).

I.5 Addressing Expected Objections

Objection 1: "DIS has directly confirmed quark charges ±2/3, −1/3." DIS measures $\Sigma q\ e^2 q\ q(x)$ — not individual charges directly. PH proto-quark charges give the same result at $Q^2 \ll \Lambda meta^2$.

Objection 2: "Asymptotic freedom is confirmed." Correct — and this is a PH prediction for $Q^2 \ll \Lambda meta^2$. The PH predicts departure from asymptotic freedom at $Q^2 \sim 10^{24}$ GeV^2.

Objection 3: "The constituent quark model predicts hadron masses." The PH mass formula gives the proton/tau ratio at 0.000% error with zero free parameters.

Objection 4: "Why introduce new terminology?" The SM quark is defined as fundamental. Calling the PH constituent a 'quark' would contradict the framework. 'Proto-quark' signals the correct relationship: the quark is the effective description; the proto-quark is its sub-structure.

I.6 Summary Table

Property	**SM Quark**	**PH Proto-Quark**	**Observationally distinct?**
Fractional charge	Input axiom	Derived from Qk formula	At $Q^2 \sim \Lambda meta^2$ only
Individual charges	+2/3, −1/3	+1/9, +4/9, +5/9	At $Q^2 \sim \Lambda meta^2$ only
Charge sums	Q(p) = +1, Q(n) = 0	Q(p) = +1, Q(n) = 0	Identical
Asymptotic freedom	Yes (running αs)	No (condensate at Λmeta)	At $Q^2 \sim 10^{24}$ GeV^2
CF-CC scattering	Not in QCD	RCF = 6.16×10^{-4}	Testable now
Mass formula	Fitted parameters	exp(A|mJ_R|^p/b); p, A derived	Testable (mass ratios)

Epistemic summary: The proto-quark is not a falsification of the quark — it is its sub-structure. The SM quark is the correct effective description at $Q^2 \ll \Lambda meta^2$. Two predictions are testable at current energies: the CF-CC rate (MicroBooNE) and the zero-parameter mass formula.

J The Chirality Asymmetry $\varepsilon\psi$ = 25/169: Results

This appendix contains the complete result set behind the chirality asymmetry parameter. Full derivations, verification scripts, and the problem registry are maintained in the companion repository [2].

J.1 Physical Mechanism

The contact interaction $-(\bar{\psi}J^a\psi)(\bar{\psi}\Sigma^a\psi)$ is odd under Z2 $\subset$ Z6: three-body LLL composites bind more efficiently than RRR composites at the Z6 phase transition. **v86:** LLL and RRR are not matter and antimatter (antimatter is made of antipsions, §13.1), and $\varepsilon\psi$ is CP-even; it does not by itself produce a baryon excess (§13.3).

J.2 The Chiral Asymmetry Formula

$$\varepsilon\psi = 4\delta\chi/(1 + \delta\chi)^2. \qquad (30)$$

For $\delta\chi$ = 1/25: $\varepsilon\psi = (4/25)/(26/25)^2 = 100/676 = 25/169 \approx 0.1479$.

J.3 Structural Origin: $\delta\chi = 1/(2j)^2$ ([T2])

The binding asymmetry between the L-sector (k = 0, 1, 2) and R-sector (k = 3, 4, 5) is a second-order effect of the JaΣa coupling; second-order corrections in a representation of spectral width 2j scale as $1/(2j)^2$. Hence:

$$\delta\chi = 1/(2j)^2 = 1/25, \qquad (31)$$

$$\delta\chi = 1/25 \Rightarrow \varepsilon\psi = 25/169 = 0.14793\ldots. \qquad (32)$$

The value $(2j)^2 = 25$ is the squared spectral width that fixes the six-state Z6 assignment (j = 5/2 is the minimal half-integer spin with exactly six states). $\varepsilon\psi$ is an exact rational of the spin-5/2 structure, not a fitted number; its first-principles derivation is the open problem P23 ([T2+]).

J.4 Former Baryogenesis Chain (withdrawn in v86)

Status (v86). The chain below is withdrawn (§13.3): ξ^* in eq. (37) was computed from the target η_B, so eq. (38) is an identity, not a verification. It is retained for the record. Former chain:

$$\eta B = \varepsilon\psi \times \varepsilon conf \times (n/n\gamma) \times (\xi^*)^{-3}, \qquad (33)$$

$$\varepsilon\psi = 25/169 = 0.14793 \text{ (exact rational fraction)}, \qquad (34)$$

$$\varepsilon conf = N3/(N2 + N3 + N4) = 4/9, \qquad (35)$$

$$n/n\gamma = 0.749, \qquad (36)$$

$$\xi^* = 434.09 \text{ (computed from the target } \eta B\text{; see status note)}, \qquad (37)$$

$$\eta B = 0.147929 \times 0.444444 \times 0.749/434.09^3 = 6.02 \times 10^{-10}. \qquad (38)$$

Note on εconf. $\varepsilon conf = 4/9$ counts four lepton-number three-body composites (e^-, μ^-, τ^-, νe) out of nine three-body families. Verification: $4/9 \Rightarrow \xi^* = 434.09$ ✓; $4/13 \Rightarrow \xi^* = 384$ ✗. Declared assumptions ([T2]): $\varepsilon conf = 4/9$ and $n/n\gamma = 0.749$ are thermal-democracy counts — equal formation weight across the three-body families at the T $\sim$ Λmeta transition — and the factorization of eq. (33) (asymmetry $\times$ counting $\times$ density, no

cross terms) is likewise declared. The direct ΩDM/Ωb route (elasticity +8.27, §14.3) doubles as an empirical bound on democracy violations at the 1–2% level. Full derivation: repository [2], P14.

J.5 Index-Normalisation Conjecture

Conjecture J.1 (P23 reformulated, [T2] candidate).

$$\delta\chi = \mathrm{Ind}(D\chi)/(2j)^2 = 1/25. \qquad (39)$$

The Dirac operator in the Z6 kink background,

$$D\chi = i\gamma\mu\partial\mu + g\,\Phi(x)\,Ja\Sigma a,\ \Phi(x) = f\pi\tanh(mZ6\,x/2), \qquad (40)$$

carries one chiral zero mode by the Jackiw–Rebbi mechanism [7]:

$$\mathrm{Ind}(D\chi) = nL - nR = 1, \text{ [T1]} \qquad (41)$$

$$W = \Delta mJ = 5/2 - (-5/2) = 5 = 2j. \text{ [T1]} \qquad (42)$$

$$\delta\chi = \mathrm{Ind}(D\chi)/(2j)^2 = 1/25 \text{ [conditional on Conjecture J.1].} \qquad (43)$$

Ind = 1 counts the topological unit of chirality; $(2j)^2$ is the spectral volume. Once the identity is proved, $\varepsilon\psi$ = 25/169 becomes a T1 theorem (its link to ηB is withdrawn, §13.3).

J.6 The Chiral Mixing Angle: Exact Theorem Suite

J.6.1 Theorem I.1 ([T1]) — Bare J·Σ diagonal in the locked subspace

In the chirality-locked subspace ($|-j\ldots-1/2, L\rangle$, $|+1/2\ldots+j, R\rangle$, σ = sign m), the bare coupling $Jz\Sigma z + \lambda(J^+\Sigma^- + J^-\Sigma^+)$ is purely diagonal, equal to |m|: $J^+\Sigma^-$ and $J^-\Sigma^+$ cannot connect two locked states. The coupling producing the asymmetry must therefore come from the condensate.

J.6.2 Theorem I.2 ([T1]) — Winding condensate activates a single bond

The Z6 winding condensate (winding fraction $\varepsilon = 1/6$ per kink) generates $J^+\Sigma^+ + J^-\Sigma^-$, activating exactly one bond $|-1/2, L\rangle \leftrightarrow |+1/2, R\rangle$ with the exact SU(2) element:

$$\langle +1/2|J^+|-1/2\rangle = j + 1/2 = 3,\ t = \varepsilon(j + 1/2) = 1/2. \qquad (44)$$

J.6.3 Theorem I.3 ([T1]) — Universality of t = 1/2 and the zero mode

$t = \varepsilon(j + 1/2) = (j + 1/2)/(2j + 1) = 1/2\ \forall j$; $H = \mathrm{diag}(|m|) + t(|L\rangle\langle R| + |R\rangle\langle L|)$ has an exact E = 0 mode. (45)

Verified for j = 1/2 through 9/2: the winding normalisation $\varepsilon = 1/Nc$ and the Jackiw–Rebbi index theorem are equivalent statements in this algebraic model.

J.6.4 The Z6 Charge Interlock ([T1])

The zero mode is $|0\rangle = (|k = 2, L\rangle - |k = 3, R\rangle)/\sqrt{2}$. With the Z6 charges of §1:

$$Q2 = -5/9,\ Q3 = +10/9,\ \Delta Q = 5/3, \qquad (46)$$

$$\langle 0|Q\Sigma^z|0\rangle = \Delta Q/2 = 5/6 \text{ [offset-independent],} \qquad (47)$$

$$\mathrm{Nc}\langle 0|\mathrm{Q}\Sigma^z|0\rangle = 6 \times 5/6 = 5 = 2\mathrm{j} \cdot \mathrm{Ind}(\mathrm{D}\chi).\ \star\ \text{EXACT INTERLOCK}\ \star \qquad (48)$$

Universality ([T1]). For every ZN, the proto-colour charge coefficient obeys $aN = [(N-1)/N]/(1-\cos(2\pi/N))$, giving $\Delta Q = (2N-1)/N$ and hence $\mathrm{Nc}\langle \mathrm{Q}\Sigma^z\rangle 0 = 2N - 1 = 2j$ identically. The Z3 coefficient 4/9 is derived, not input. Corollary ($\Sigma Qk = 1$, [T1]): $Qk = q(k \bmod 3) + \lfloor k/3 \rfloor$ with $q = (1/9, -5/9, -5/9)$, so $\Sigma Qk = 2\Sigma q + 3 = 1$ — the unit sum is the integer Z2 chirality shift, a structural corollary of the Z3 $\times$ Z2 factorisation; the §J.6.4 charge difference is offset-independent.

J.6.5 Chiral Mixing Angle ([T1])

The formula $\varepsilon\psi = 4\delta\chi/(1+\delta\chi)^2$ is the double-angle identity under $\delta\chi = \tan^2\theta\chi$:

$$\varepsilon\psi = \sin^2(2\theta\chi);\ \tan\theta\chi = 1/5 \Rightarrow \sin 2\theta\chi = 5/13,\ \cos 2\theta\chi = 12/13, \qquad (49)$$

$$\varepsilon\psi = (5/13)^2 = 25/169.\ [\text{Pythagorean triple } 5^2 + 12^2 = 13^2] \qquad (50)$$

J.6.6 Lemma I.1′ — the single open step ([T2+])

$$\tan\theta\chi = \mathrm{Nc}\langle 0|\mathrm{Q}\Sigma^z|0\rangle/(2\mathrm{j})^2 = 5/25 = 1/5.\ [[\mathrm{T2+}], \text{open}] \qquad (51)$$

By spectral reduction (using only the interlock and t = 1/2, both T1), Lemma I.1′ is algebraically equivalent to $\tan\theta\chi = t/j$: the charge structure drops out; what remains is a pure spectral statement.

J.6.7 Jackiw–Rebbi Anchor and Structural Protection ([T1])

At j = 1/2, Lemma I.1′ gives $\tan\theta\chi = 1 \Rightarrow \varepsilon\psi = 1$ — exactly the Jackiw–Rebbi result (the kink zero mode is 100% chirally polarised) [7]. PASS. Edge protection: at j = 1/2 there is no leakage channel, so $\varepsilon\psi(\mathrm{Z2}) = 1$ for all condensate gaps G and all λ. Z4 invariant-subspace protection: the sector $S = \{(-1/2, \mathrm{L}), (+1/2, \mathrm{R}), (-3/2, \mathrm{R}), (+3/2, \mathrm{L})\}$ satisfies $H(\lambda, G)\cdot S \subseteq S$ for all λ, G, forcing $\varepsilon\psi(\mathrm{Z4}) = 9/25$ exactly for any eigenstate in S. Finite-gap correction law: $\varepsilon\psi(G) = 25/169 - \kappa 6/G^2 + O(1/G^3)$ for Z6, with exact leading term $-480/2197$ from first-order perturbation theory.

J.6.8 General Formula

j	**Group**	**tan $\theta\chi$**	**$\varepsilon\psi$**	**Pythagorean triple**	**sec($2\theta\chi$)**
1/2	Z2 (JR)	1	1	—	—
3/2	Z4	1/3	9/25	(3, 4, 5)	5/4
5/2	Z6	1/5	25/169	(5, 12, 13)	13/12
7/2	Z8	1/7	49/625	(7, 24, 25)	25/24

Pattern: $\tan\theta\chi = 1/(2j)$; triple $(2j, 4j^2 - 1, 4j^2 + 1)$. Internal falsification target: a full Z4 dynamical calculation must give $\varepsilon\psi = 9/25$ (protected exactly by the invariant-subspace theorem).

J.7 Z6 Gauge Structure Theorems ([T1])

V.1 (Z3 $\times$ Z2 charge decomposition). $Qk = q(k \bmod 3) + \lfloor k/3 \rfloor$; the Z2 factor acts as $Q \to Q + 1$; every psion carries unit Z3 triality, $9Qk \equiv 1 \pmod 3$.

V.2 (Selection rules). All $Q = -1$ lepton composites and their flip intermediates are colour singlets; the flip $2 \to 3$ conserves triality. The flip cost comes from the Z6 flux/contact sector, not from confinement.

V.3 (Rationality). Eflip = 2j·t = 5/2 must be rational; one-meta-boson exchange (carrying αmeta = π/6) is excluded as the mechanism.

V.4 (Flux-quantum count). The flip transports $\Delta Q = 5/3 = 2j$ quanta of size 1/3; for every Z2N, quanta = $Nc\Delta Q/2 = 2N - 1 = 2j$ identically — the charge interlock is a flux count.

V.5 (Factorisation). $Z6 \cong Z3 \times Z2$ (CRT) and the charge table factorises exactly; the relevant statistical system is a product of a Z2 and a Z3 system, each with a unique Kramers–Wannier self-dual transition (no BKT window).

Theorem J.3 (V.7: equivalence web, [T1]). *The following are exactly equivalent, verified on all four anchor groups: (1) Lemma I.1′; (2) tan $\theta\chi$ = t/j; (3) Eflip = 2j·t = j; (4) the flip intermediate sits at the locked spectral edge; (5) flux costs equal (t, N − 1) = (half bond splitting, bond→edge distance). Conditionally on L1 ([T2]), all are further equivalent to (6): the colour factor sits at its Wegner self-dual point. No algebraic reformulation can close the set to T1; a dynamical input is required.*

J.8 Self-Duality Results ([T1])

SD-1. The winding coupling $\varepsilon = 1/Nc$ equals the Villain 4D self-dual coupling $\alpha sd(ZN) = 1/(2N)$ exactly, on all four anchor groups: the Z3 colour factor of the condensate sits exactly at the Villain self-dual point.

SD-2. ZN is Pontryagin self-dual; invariance under $q \mapsto N - q$ forces E(flux q) = f(min(q, N − q)) — the conjugacy min-rule is derived by group theory, not chosen.

SD-3. With $\Phi(\Lambda) = \Lambda$ (P2/P12), the Z6-uniform condensate $\langle\bar{\psi}k\psi k\rangle = \sigma/Nc$ factorises the effective action exactly: Seff = Seff(Z3) × Seff(Z2).

SD-4. The single matter coupling t = 1/2 places both factors simultaneously at their self-dual couplings: $\alpha Z3 = t/3 = 1/6 = \alpha sd(Z3)$, $\alpha Z2 = t/2 = 1/4 = \alpha sd(Z2)$, with no free parameter. SD-4 fixes the matter coupling; Eflip runs through the gauge stiffness κ — a distinct coupling — so self-duality alone does not close P23.

J.9 The ν Quantum Number and the Locking Mechanism

Theorem J.4 (ν.1, [T1]). *$\nu \equiv \lfloor k/3 \rfloor \oplus \chi$ is a hidden good quantum number: $\nu = 0$ is exactly the locked subspace, verified state by state on all twelve.*

Theorem J.5 (ν.2, [T1]). *Punlocked = $-(1 - \hat{Z}2\cdot\chi)$: the deep-gap projector of the algebraic model is the ν charge, G·Punlocked = G·Nν, exactly.*

Locking mechanism ([T2], exact coefficients). Projections of the §2 interactions onto the ν-channel: f1 = 0 (direct route killed by kinematic identity), f2 = 1 (scalar), f3 = 3/4 (tensor-induced, exact), f4 = 1/12 (bubble). Locking is tensor-induced ν condensation: the tensor condensate ΔT forms in the $J\Sigma$ channel, then breeds the ν gap through the exact 3/4 coefficient — a self-reinforcing bootstrap in which ν-conservation is emergent in the deep-gap limit. Quantitative closure is open: the registered multiplicities give ceff = f2 + 2f3 + f4 = 11/6 against the 2G convention of the operational definition; resolving the discrepancy requires the self-consistent non-commuting two-stage solve (P26). Falsifiable corollary: if ν condenses, the $\nu = 1$ sector is a physical multiplet at mass ∼2G, to be confronted with the hAB resonance (∼0.6 GeV).

J.10 The κ/G Barrier: Exact Form and Interpretation

Every analytic route from the §2 action to the k-ladder gauge stiffness yields the same exact ratio ([T2]): electromagnetic duality with βmag = Nc (which follows from $g^2 = 2\pi^2/3$ via βelec $\times$ βmag = $Nc^2/4\pi^2$, itself a new T2 result) gives κ/G = $Nc^2/(4\pi^2) = 9/\pi^2 \approx 0.912$ — an 8.8% deficit against the pass criterion κ/G = 1, fixed by Nc = 6 versus 2π. Ward identities do not determine κ (charge is incompressible in the deep gap); κ is a construction datum of the emergent-gauge map, not a symmetry consequence.

J.10.1 The sinc^2 Identity ([T2])

$$\kappa/G = \mathrm{sinc}^2(\pi/Nc),\ \mathrm{sinc}(x) = \sin(x)/x;\ \mathrm{sinc}^2(\pi/6) = (1/2)^2/(\pi/6)^2 = 9/\pi^2. \qquad (52)$$

sinc is the lattice-to-continuum energy ratio of the fundamental ZNc mode: the deficit is the Nyquist sampling error of the massless psion on a six-point lattice; $\kappa/G \to 1$ as Nc $\to \infty$. Geometric form: $\sqrt{(\kappa/G)}$ = chord/arc of a Z6 circular sector (half-angle 30°): the discrete jump between Z6 sectors versus continuous propagation on the phase circle. The gap G is itself lattice-generated (the massless psion has Gcont = 0), so both κ and G are intrinsically lattice objects and the sinc^2 is a genuine physical ratio, not a removable artefact.

J.10.2 The Psionic Fine-Structure Constant ([T2])

$$\alpha\psi = 1 - \mathrm{sinc}^2(\pi/Nc) = 1 - 9/\pi^2 \approx 0.0881,\ 1/\alpha\psi \approx 11.35. \qquad (53)$$

The massless psion's fine structure separates into three constants: αmeta = $\pi/6$ (dynamical coupling), ε = 1/6 (winding unit), $\alpha\psi \approx 0.0881$ (geometric sampling deficit) — a separation that is itself the signature of masslessness (a massive field collapses all readings to one constant, as the electron does to 1/137). Numerical observation, not a derivation: $\alpha\psi \approx 12\alpha$ to 0.6%.

J.10.3 The Fundamental Identity ([T2])

$$\mathrm{sinc}^2(\pi/Nc) \times g^2 \times gL \times t/(2j+1) = 1. \qquad (54)$$

Exact: $[Nc^2/4\pi^2] \times [2\pi^2/3] \times [2] \times [(1/2)/6] = 1$, every factor independently [T1]. The identity states that the compensating factor $1/\mathrm{sinc}^2 = \pi^2/9$ equals $g^2 \times gL \times t/(2j+1)$; it establishes consistency of κ/G = 1 with the derived parameter set, but not that the effective stiffness κeff takes this form — that derivation is the §2.3 $\to$ k-ladder map, the open content of P23.

J.10.4 Scale-Hierarchy Link ([T2])

$Nc/(2\pi) = \mathrm{sinc}(\pi/Nc)$ exactly. The observed hierarchy $\ln(\Lambda\mathrm{meta}/mZ6) = 33.68$ lies between $S/(2\pi) = 33.85$ (continuum, 0.5%) and S/Nc = 35.45 (discrete, 5.2%); their ratio is exactly the P23 gap factor $\mathrm{sinc}(\pi/Nc) = 3/\pi$. The physical scale separation is set by the continuum psion dispersion while the κ/G deficit reflects the discrete Z6 lattice — two faces of one tension. Additional exact identity: S $\times$ be = 6Ne = 60 = 10Nc.

J.11 The fπ Branch Question

Two exact, mutually exclusive forms for the condensate decay constant coexist in the framework: Branch A, fπ = me (from the App. B identity σdw/N0 = 1 with mZ6 = 6me, P24); Branch B, f$\pi = \sqrt{(3/\pi)}$ me (explicit

in §3.2, where the normalisation $\lambda = 9/2$ is forced by mZ6 = 6fπ — Theorem W.1, [T1] — and $3/\pi = Nc/(2\pi)$ gives Branch B the shape of a derived relation). The branches differ by 2.28%; existing numerical routes sit between them. Decisive question: the provenance of the §3.2 matching. Quarantine ([T1] in scope): the baryogenesis pipeline (eq. 33) contains no fπ or σdw dependence, so the fork cannot propagate to ηB, Neff, or the dark-matter chain; it is contained to {P24, Conjecture CEC, §3.2}.

J.12 Axiom I.1 — The κ = G Postulate

Postulate P-κ (Axiom I.1). κ = G: in the deep-gap limit of the k-ladder construction, the electric stiffness of the emergent Z6 gauge sector equals the matter gap.

Status: postulate, not theorem. Every algebraic route is blocked by Theorem V.7; every analytic route from §2 yields $\kappa/G = \mathrm{sinc}^2(\pi/Nc)$. The axiom is equivalent to asserting $\alpha\psi = 0$ at the physical point — that the emergent-gauge construction supplies the $1/\mathrm{sinc}^2$ enhancement, whose required form is exactly the fundamental identity (eq. 54).

What it delivers: $\kappa = G \Rightarrow$ Eflip = j $\Rightarrow$ Lemma I.1′ $\Rightarrow \varepsilon\psi = 25/169$ (T1 conditional); the further step to ηB is withdrawn (§13.3).

Falsifiability: any explicit construction of the §2.3 $\rightarrow$ k-ladder emergent-gauge map yielding $\kappa \neq G$ refutes it; conversely, a construction delivering the eq. 54 factor set proves it.

K Gravitational Sector: Condensate Response

If GN is the tensorial susceptibility of the Z6 condensate, the vacuum is a medium with characteristic frequencies, resonances, and penetration depths.

K.1 Convention CN-1 (Planck Convention)

The susceptibility relation $GN = K(\xi)\,\Lambda meta^{-2}\exp(-S/g^2)$ is written in the standard (non-reduced) convention, $GN = MPl^{-2}$ with $GN = 6.70883 \times 10^{-39}$ GeV^{-2} (CODATA 2018 in natural units) and $MPl = 1.22089 \times 10^{19}$ GeV. Any appearance of 8πGN must cite this note. Self-consistency closes exactly:

$$S/g^2 = 2\ln(MPl/\Lambda meta) + \ln K(\xi) = 32.323 \text{ (identically } 6Ne/(be\ g^2)). \qquad (55)$$

K.2 Lemma J.1 — Kinematic Gravitational Josephson Frequency ([T2+])

Premise P1 (open): the order parameter relevant to GN is bilinear in the lightest composite, pair rest energy $2mec^2$.

Premise P2 ([T1], established): in the static weak field, a state of rest energy E accumulates phase at $\dot\theta = -E(1 + \varphi/c^2)/\hbar$ (gravitational redshift of the Compton clock; Chou et al., Science 329, 1630, 2010).

$$fJ = (2mec^2/h)\cdot(\Delta\varphi/c^2) = 2.6946 \times 10^4\ \mathrm{Hz/m} \times L \text{ (terrestrial)}. \qquad (56)$$

The factor 2 is the exact analogue of the Cooper-pair charge 2e. Prediction [G1] ([T2+]): parameter-free.

K.3 Proposition J.2 — Topological Clock Identity (conditional on P24)

$$\text{ftop} = \text{mZ6c}^2/(6\text{h}) = \text{mec}^2/\text{h} = \text{fe} \text{ [if mZ6} = 6\text{me]}. \qquad (57)$$

Arithmetic identity given P24; if P24 closes at T1, the vortex tick is the electron Compton clock. Cross-check: $\text{m}\mu/6 = 17.6097$ MeV differs from mX17 = 17.62 MeV by 0.06% — under the general-level reading $\text{m}\mu = \text{Nc}\cdot\text{m}(9)$, mX17 = m(9), this is an identity of the cascade level N = 9 (numerical observation; repository [2]).

K.4 Conjecture CEC and Proposition J.3

Conjecture CEC ([T4]; [T3] with pipeline): the Z6 vacuum energy density equals the Compton-scale density of its lightest composite:

$$\text{ucond} = (\text{mec}^2)^4/(\hbar\text{c})^3 = 1.42 \times 10^{24}\ \text{J/m}^3. \qquad (58)$$

Evidence: inverting fplasma = 18 mHz gives 6.6×10^{-14} GeV^4 vs $\text{me}^4 = 6.82 \times 10^{-14}$ (3%); no other framework scale reproduces an 18 mHz plasma frequency. A single ratio at 3% is weak evidence in isolation; the conjecture is internally falsifiable by any derivation of ρcond.

Proposition J.3 ([T2] conditional on CEC): the gravitational plasma frequency is parameter-free:

$$\text{fplasma} = (1/2\pi)\ \sqrt{(\ 4\pi\text{GN}\ (\text{mec}^2)^4\ /\ ((\hbar\text{c})^3\ \text{c}^2)\ )} = 18.3\ \text{mHz}, \qquad (59)$$

placing the [G2] cutoff in the LISA band; the interpretation as collective screening remains [T3] pending PC2.

K.5 Consistency Problem PC2 — Static vs. Dynamic Screening

[Open] A static reading of the penetration depth λgrav = c/(2πfplasma) $\approx 2.6 \times 10^9$ m is falsified by planetary orbits; only dynamical perturbations below fplasma may experience collective response, and this distinction is not yet established in the PH. Until PC2 resolves, [G2] and [G3] are not rigorously falsifiable.

K.6 Predictions [G1]–[G3]

[G1] ([T2+]): fJ = $(2.6946 \times 10^4$ Hz/m) $\times$ L (Lemma J.1, eq. 56); parameter-free given Premise P1.

[G2] ([T2] cond. CEC; interpretation [T3]): suppression of stochastic GW power below 18.3 mHz (LISA band); PTA detections of the nHz background require a quantitative amplitude prediction before confrontation.

[G3] ([T3]): resonant δGN/GN $\sim 10^{-10}$ at fdrive $\in$ [0.38, 0.43] GHz, via the pipeline effective Lagrangian $\rightarrow$ per-cell coupling (Hypothesis A2, [T4]: the driving field couples photon-per-coherence-cell, depositing Edrive per composite pair; under A2 the identification Edrive = h·fdrive is legitimate — without it, Edrive is an energy density and the frequency conversion does not follow) $\rightarrow \chi(\omega) \rightarrow \delta\text{GN}(\omega)$.

K.7 Open Problem Registry

Full registry in repository [2], `docs/PROGRESS_AND_ROADMAP.md`.

ID	Problem	Status	Blocks
P23	κ = G (Lemma I.1′) — §2.3 emergent-gauge construction	[T2+]	$\varepsilon\psi \rightarrow$ T1, Neff $\rightarrow$ T1 (ηB link withdrawn)

P14-B	Conserved baryon (rung) label; sign and size of the bifurcation bias (§13.4, §13.7)	open	ηB, nucleon stability, Theorem KS
P14-C	Baryon binding scale ($(\hbar c)/a \approx 163$ MeV) and internal structure consistent with $J{\cdot}\Sigma$ mixing and with the Gottfried sum (valence $-1/9$; sea term $+0.346$ needed, not derived); hexagonal radii are phenomenological (App. L.4.7); dynamic-ring series closed: negative on radius–mass, hybrid description demoted, per-composite geometry unsupported, minimal spin extension fails on magnetic moments (App. L.4.8)	open	hadron radii, $m_n - m_p$, sea model
P26	ν-gap quantitative closure (ceff = 11/6 vs 2G)	[T2]	P23 (locking mechanism)
P25	Z6 composite gauge vertex	[T2]	P23, [G3], PC1-sub
P-4π/3	Prefactor 4π/3 in ΩDM/Ωb direct formula	[T2+]	ΩDM $\rightarrow$ T1
P1	be = 325/1152 from 2PI kernel	[T2]	mass ratios Class B
P2/P12	$\Phi(\Lambda) = \Lambda$ fixed point; closure $\rightarrow$ adimensional ratios internal; 1 dimensional datum (Remark 5.1)	[T2]	GN, me $\rightarrow$ predictions
P8b	Passo A.3 integration ($\Sigma 2$ = be $\times$ 15/8)	[T2]	lepton mass ratios precision
P24	mZ6 = 6me vs $6\sqrt{(3/\pi)}$me (fπ branch, §J.11)	[T2]	Prop. J.2, CEC
C-Δm21	Derive $\Delta m^2{}_{21}$ = (18/25) m(15)^2 from the ν-channel (P26-C)	[T4]	second datum $\rightarrow$ prediction
PC1	Gravity: emergent vs. ontological (hAB vs. graviton)	[T2]	[G3] pipeline
PC2	Static vs. dynamic screening of λgrav	open	[G2], [G3] falsifiability
P-PMNS	Sector-bridging PMNS texture; CC bridge = proto-W (I = 3); texture open; flavour basis fixed, eq. (1) alone gives $\sin^2\theta_{23} \approx 0.21$ (§17.3)	[T3]	mixing sector; F3
P-|O|	Overlap sector for absolute rates; $\|O\|^2$(CF) ≤ 0.370 (2σ)	[T2]	rates $\rightarrow$ T1
P-εconf	Formation-overlap democracy (εconf = 4/9, n/nγ)	[T2]	withdrawn with eq. (33) (§13.3)
P-Pauli	Pauli structure (restated): auto-singlets from N $\geq$ 7; spatial shells from N $\geq$ 13, nuclei at N = 15+; per-level structure open; 39-count floor	[T2]	DM spin sector
C-Q-rat	Charge rationality / integer quantization as Z6 selection principle	[T3]	why Nc = 6
P-Lorentz	Local Lorentz invariance: D1 + R1 resolved (scalar channel; conserved-charge theorem); open: loop-level E/Λmeta vs SME; induced-metric universality	[T2+]	emergent-metric T-status
P-walls	ZKO relic walls: resolved-conditional (IR tension + exact chiral-bias spectrum 5:8:9); open: κCP·V_0 normalization	[T2/T3]	cosmological safety
F-fork	RESOLVED $\rightarrow$ twelve-state bookkeeping via U-count (U3: (2, 2π/3)); residuals reviewed and held	[T2]	closed
C-web	Closed-form master web (§5.1) RESOLVED: exact rearrangements of registered structure; O(1/S) two-loop slot	[T2+]	master invariant; F17

Table 15: Open problems and cascading dependencies. P23 is the critical path; P2/P12 is the deepest.

K.8 The Chiral-Flip Channel and the e^+e^- Sector

The Z6 composite structure predicts the Chiral-Flip Charged Current, a zero-parameter structural channel ([T1]: selection rules, existence) with no analogue in three-neutrino QFT:

$$\nu\mu(0,4,2) + \mathrm{n}(0,1,5) \rightarrow \mathrm{e}^-(0,1,2) + \mathrm{p}(0,4,5),\ \mathrm{RCF} = \mathrm{NDirac}\ \mathrm{g}^2\mathrm{meta}\ (\mathrm{me}/\mathrm{m}\mu)^2 \times |O|^2 = 6.16 \times 10^{-4}\ |O|^2. \tag{60}$$

Here $|O|^2 \in [0, 1]$ is the BSE formation overlap (open sector P-|O|).

Theorem K.8.1 (flatness, [T1]). *Z6 vertex conservation forces any t-channel mediator to carry (I, Q) = (3, +1). Exhaustive enumeration of the 36 bilinears shows the only such states are the six proto-W members at MW; their propagator is flat to $< 2 \times 10^{-4}$ over accelerator kinematics, and hAB (I = 0, Q = 0) cannot mediate. RCF is therefore energy- and baseline-independent at every accessible order.*

Experimental status. A flat ratio cannot reproduce a spectrally localized excess for any $|O|^2$; MicroBooNE's exclusive 1e1p channel [10, 14] bounds $|O|^2 \leq 0.244$ (90% CL) and ≤ 0.370 (2σ), i.e. RCF $\leq 2.28 \times 10^{-4}$ — below the MiniBooNE anomaly and decoupled from it. Confrontation protocol: a genuine CF-CC signal appears as a 1e1p excess identical (after flux scaling) at SBND (110 m) and ICARUS (600 m), incompatible with any L/E fit, at RCF $\leq 2.28 \times 10^{-4}$; SBND's percent-level νe-like rate measurement tightens the $|O|^2$ ceiling directly.

Theorem K.8.2 (e^+e^- uniqueness, MB-U, [T1]). *A composite can decay to e^+e^- only if its net psion number vanishes — the mixed $\psi\bar{\psi}$ sector; every pure-psion composite is excluded by conservation. Enumeration of the neutral $\psi\bar{\psi}$ spectrum leaves a single massive candidate below 0.6 GeV: the proto-colour singlet of the X17 family at cascade level 9 (17.62 MeV).*

The next neutral $\psi\bar{\psi}$ states — the level-8 family member (0.610 GeV) and hAB (0.613 GeV) — lie above the short-baseline excess window and decay to two-shower $\gamma\gamma$ topologies, outside the single-ring e^+e^- class. Any dark-sector e^+e^- interpretation of short-baseline data is therefore confined, within the PH, to this one state. The corresponding conjecture (C-MB-X17) is registered in the repository under E4 quarantine with ex-ante kill criteria and is not carried in the prediction registry (Table 4).

K.9 Local Lorentz Invariance of the Emergent Metric

Every condensate-based emergent gravity owes an account of why local Lorentz invariance is exact or suppressed below experimental bounds: a condensate generically selects a frame. The PH clears this at every currently computable level. (i) [T1] The contact interaction of eq. (1) carries no Dirac structure — J, Σ act on the internal space and commute with every γ^μ — so the available condensation channels $\langle\bar{\psi}\psi\rangle$, $\langle\bar{\psi}J\Sigma\psi\rangle$ are Lorentz scalars; a scalar condensate preserves Lorentz exactly, as the QCD chiral condensate does. (ii) [T2] The Fierz-induced vector and axial channels cannot condense: they couple to conserved (axial-)charge densities, which vanish in the $\mu = 0$ vacuum, and the mean-field solution is uniquely trivial; the C-invariance of the interaction independently forbids the C-odd vector order parameter, and the tensor projection is subcritical (relative Fierz coefficient $-1/8$). (iii) Consequences: c_gw = c_γ by structure — the photon bilinear and the metric response propagate on one Lorentz-invariant substrate (GW170817 satisfied without tuning) — and the equivalence principle as single-substrate universality (argument grade; the induced-metric computation is the open deliverable). Residual: loop-level operators in powers of E/Λmeta ($\approx 8 \times 10^{-10}$ at 1 TeV) against the SME tables; if the suppression scale is Λmeta rather than MPl, dimension-six coefficients sit near future sensitivity — a testable window.

K.10 The Fate of the Z6 Domain Walls

Spontaneous breaking of the discrete Z6 at the first-order transition T = Λmeta Kibble-produces a relic wall network — the Zel'dovich–Kobzarev–Okun problem, fatal for a naive tension Λmeta^3 (fifteen orders above the CMB bound $\sigma^{1/3} \lesssim 1$ MeV). The resolution has two layers.

Layer 1 (infrared walls, [T2]). The walls interpolate the condensate phase between adjacent vacua — pseudo-Goldstone territory, stiffness $f\pi^2$, gap mZ6, both MeV scale ($f\pi = \sqrt{(3/\pi)}\cdot$me = 0.4994 MeV, ratified: the §3 normalization is derived by the 2π condition, $f\pi^2 = Nc\cdot me^2/(2\pi)$; mZ6 = $6f\pi$ = 2.996 MeV, with the

P18 route differing by 1.1% — audit Q2). The registered virial tension σ = fπ^2mZ6/π^2 = 0.0757 MeV^3 gives $\sigma^{1/3}$ = 0.423 MeV: forty-six orders below the naive estimate and below the Zel'dovich bound. Even a fully persistent scaling network is invisible at current sensitivity: ρwall/ρcrit $\approx 3 \times 10^{-6}$ today, induced δT/T $\approx 3.5 \times 10^{-7}$, BBN-era impact $\approx 3 \times 10^{-22}$. The normalization is convention-sensitive (the sine-Gordon form gives 1.82 MeV, straddling the bound), so robustness rests on Layer 2; the residual mZ6 provenance audit (Q2) is thereby promoted to a cosmological-safety input.

Layer 2 (chiral bias, [T2] shape / [T3] normalization). The six vacua are not protected-degenerate: mJ(k) = k − 5/2 is injective, so no nontrivial Z6 shift preserves the chiral structure. The splitting spectrum inherits the exact chiral autocorrelation ΔC(s) = C(0) − Σk mJ(k)mJ(k+s) = {0, 15, 24, 27, 24, 15}: three wall classes in ratio 5 : 8 : 9, all positively biased, with the s = 3 walls (the pure L↔R exchange) maximally split and dying first. The normalization follows by spurion analysis: any vacuum-splitting operator crosses chirality twice, forcing κCP = (me/Λmeta)^2 — the same CP-odd scale that powers baryogenesis. The estimated annihilation time, t_ann ~ $\sigma/\varepsilon \approx 4$ yr, lands after BBN (where the walls are harmless) and well before recombination: the asymmetry that makes the matter kills the walls. Falsifiable branch: should the bias fail, the persistent network predicts δT/T ~ 4×10^{-7} ISW-class signatures, within reach of near-future CMB experiments.

Status note (v86, open tension). K.10 and the CP source of §13 are not consistent as written. §13 states that $U(1)_M$ is exact in eq. (1) and that the vacuum phase of the winding condensate can be rotated away in every vacuum. K.10 treats the six vacua as discrete, with a wall gap $m_{Z_6} = 6f_\pi$ and a vacuum splitting $\kappa_{\rm CP} = (m_e/\Lambda_{\rm meta})^2$, which is an explicit breaking of $U(1)_M$ down to Z_6 and then beyond. If the symmetry is exact and spontaneously broken, the vacuum manifold is a circle and the associated mode is a massless Goldstone boson, not a pseudo-Goldstone with a gap, and there are no discrete walls. The explicit breaking that K.10 needs is not derived from eq. (1). Until it is, the wall network, the 4-year annihilation time and the ISW branch of this subsection are conditional on an additional input; the CMB bound on $\sigma^{1/3}$ is not affected by this note. A finite-model test (WP-VAC, pre-registered, three-psion sector) supports the tension: the ground energy is independent of the condensate phase to 3×10^{-14} over 96 phases, it is equal at the six phases while $\Delta C(s) = \{0, 15, 24, 27, 24, 15\}$, and the explicit $\Delta M = 1$ bond gives a single dominant harmonic ($\propto \delta^2 \cos\theta$) with the sixth harmonic at rounding level (ratio $\leq 10^{-9}$), so it cannot produce six equal minima. The test does not include spatial gradients and so does not measure a Goldstone mass; a $\Delta M = 6$ operator would be an additional postulate. A second test (WP-DM6) adds fixed-phase explicit breakings X^j + h.c. ($\Delta M = 2j$): each gives its own harmonic j with small admixtures, the sixth harmonic arises only at 1.4×10^{-3} of the third from X^3, and X^j vanishes for $j \geq 4$ with three psions, so six equal minima need a $\Delta M = 12$ operator that this sector cannot test. The bias $\Delta C(s)$ decomposes as $17.5 - 12\cos(\pi s/3) - 4\cos(2\pi s/3) - 1.5\cos \pi s$; no single operator reproduces it (best shape deviation 0.30), and three free amplitudes would fit it without deriving it. The wall structure and the bias are therefore inputs, not consequences of eq. (1).

L Method Audit Record: Eliminated Routes and Negative Findings

This appendix records the adversarial audit trail of the derivation programme: eliminated routes, self-caught circularity audits, and negative findings. A route killed by its own audit is a result, and is registered with the same care as a theorem; the complete session-level record, including all intermediate numerics, is preserved in the repository [2].

L.1 The $\varepsilon\psi$ Campaign: Audits and Kill Tables

The $\varepsilon\psi$ = 25/169 closure campaign produced, beyond the final results carried in Appendix J, an audit trail whose density is itself information: every naive shortcut to T1 was tried and killed. Eight self-caught circularity audits are on record. C-I8.1: the 'one hop, one quantum' route killed — a single hop transports 2j = 5 flux quanta, not one. C-I8.2: flux-energetics linearity gap — Schemes A (linear) and B (min-rule) registered as anchor-degenerate. C-I8.3: the holonomy definition $g^2 \equiv 2\pi/N$ presupposes self-duality — circular, killed before registration; the same audit corrected an overstatement of V.6 (equivalence, not support). C-I8.4: Scheme B's spectral identification was reverse-solved from the target — killed as derivation, retained as exact equivalence. C-I8.5: architecture finding — the §2 Lagrangian carries Z6 as a GLOBAL symmetry (domain walls, not gauge strings); the gauged-ladder line was reclassified as framework-ungrounded model study. C-I8.6: the framework JΣ operator is not the Pauli $J\cdot\sigma$; the locked diagonal values are not eigenvalues of the full operator. C-I8.7: the combination 2f3 + f2 = j is target-forced — registered as equivalence, killed as closure. C-I8.8: the self-duality route reduces to J = t, a restatement of Lemma I.1′.

Kill tables (exact). JR-anchor competitor table: $\langle Q\Sigma\rangle/(2j)^2$ and $\langle Q\Sigma\rangle/(2j)$ killed at j = 1/2 ($\varepsilon\psi$ = 16/25 $\neq$ 1); the linear form $Nc\langle Q\Sigma\rangle/(2j)$ killed at j = 5/2 ($\varepsilon\psi > 1$); the survivor is $Nc\langle Q\Sigma\rangle/(2j)^2 \equiv Ind/(2j)$. Charge-channel survey for Eflip (flip (0,1,2) $\to$ (0,1,3)): pair channel ΣQaQb requires g = −27/8 (not derived, killed); self channel ΣQa^2 requires g = 27/10 (reverse-solved, killed as derivation); total-charge and charge–chirality channels require |g| = 9/2, opposite signs (not derived, killed). Verdict: the charge sector is forced as the splitter (Theorem L.1) but not computable from the derived structures.

Theorem L.1 (flip degeneracy, [T1]). The kink bond exchanges k = 2 $\leftrightarrow$ 3 (m = −1/2 $\to$ +1/2 on one constituent); since |m| and m^2 are flip-invariant, both Σ|mi| and $C2 = \Sigma mi^2$ are exactly degenerate between any Q = −1 composite and its flip partner — e.g. (0,1,2) $\leftrightarrow$ (0,1,3), both at Σ|m| = 9/2, C2 = 35/4. The unique derived quantity split by the flip is the Z6 charge, ΔQ = 5/3: the splitter is forced to be the charge sector.

Exact JΣ spectrum (exact result). The framework operator $Jz\Sigma z + \lambda(J+\Sigma- + J-\Sigma+)$ with unit-element Σ± has exact spectrum {5/2 (×2), $-1/2 \pm \sqrt{3}$ (×2 each), $-1/2 \pm \sqrt{21}/2$ (×2 each), 1, −2}; the locked diagonal values 1/2 and 3/2 are NOT eigenvalues of the full operator — the deep-gap projection is essential, and its physical origin is exactly the P26 locking mechanism. The naive continuum embedding Hint = M·(J·Σ) was killed by direct 24-component Dirac diagnostics in the Z6 kink (bulk thresholds at M·{1, 2, 2.5, 3.5}, no zero mode).

S7 prototype and the three-body negative finding. The gauged k-ladder electrostatics scan (twelve configurations: string orientation × Z6 background) produced exactly ONE pass: $Eflip = (1/2)(2\cdot d(4)^2 - 3\cdot d(1)^2) = 5/2 = j$, with the Z6 conjugacy min-rule realized dynamically. Sensitivity ledger: $\kappa = 1$, background c = 0, and string geometry were NOT derived — three undetermined choices. The three-body extension (μ = (0,1,2) $\to$ (0,1,3), every simple prescription) FAILED to reproduce 5/2 (results ranged −23/2 to 19/2): the prototype hit is possibly a single-particle truncation artifact, and numerics were frozen pending the kink-as-gauge-object derivation. This stop condition — no prescription-shopping — is the campaign's discipline in action.

Operational definition of $\varepsilon\psi(G)$ and correction constants. $\varepsilon\psi(G) = \sin^2(2\theta\chi(G))$ with tan $\theta\chi(G)$ =

$Nc\langle\psi 0(G)|Q\Sigma z|\psi 0(G)\rangle/(2j)^2$, evaluated on the dynamical zero mode with unlocked penalty 2G. First-order symbolic PT reproduces the leading term $-480/2197$ fraction-for-fraction. Registered constants: $\kappa 6 = 0.228(1)$, with a reconstruction flag — the re-derived value is 0.2208, a 3% discrepancy attributed to subleading extraction, candidate corrected value $\kappa 6 \approx 0.221$; $\kappa 8 = 0.500(1)$. These numerics establish dynamical robustness of the formula's ingredients; they do not, and cannot, prove the Lemma.

Route eliminations. Ward identity: no canonical normalization produces $\kappa = 1$ ($\chi QQ \to 0$ in the deep gap; charge incompressible) — κ is a construction datum, not a symmetry consequence. Phase-stiffness discretization: $\kappa/G \sim 10^{-5}$, five orders short; the required enhancement carries the exact factor $Nc^6/(4\pi^2)$ — the first structural lead pointing to the vortex/winding mechanism, which then delivered $\kappa/G = Nc^2/(4\pi^2) = \mathrm{sinc}^2(\pi/Nc)$ (§J.10). Kink energy (virial, $Ekink = f\pi^2 mZ6/\pi^2$, a T2 by-product of the campaign): $\kappa/G \in \{0.203, 2\}$ depending on identification — no route gives 1. P26 Round 2: the eight natural multiplicity assignments yield $ce \in \{11/6, 9/4, 31/12, 43/12, 2, 13/4, 3, 4\}$; the single assignment hitting 2 contradicts the registered $w3 = 2$ — the R1/R2 fork of §J.9 in its original data-first form.

Failed-route lesson. The leading-order field-theoretic estimate $\delta\chi \sim f\pi^2/(\Lambda meta^2\, Ne^2) \sim 10^{-32}$ fails because it computes an energy ratio between microscopic scales, exponentially suppressed by $S \approx 213$; the correct quantity is a geometric ratio of representation, dimensionless and Λmeta-independent — the spectral width $2j = 5$ provides exactly this. Wilson coincidence (registered, not promoted): for Z6 and only Z6, the unit-coupling Wilson plaquette cost equals the derived bond, $1 - \cos(2\pi/6) = 1/2 = t$; Z6 is the largest ZN with rational Wilson cost (crystallographic N), consistent with Lemma V.3 singling out the physical group.

L.2 XL-1 Execution Record (the fπ fork, four routes)

Four-route comparison (data-first): §3.2 explicit form $\sqrt{(3/\pi)}\cdot me = 0.4994$ MeV; XL-1/P24 exact form $f\pi = me = 0.5110$ MeV; P18 table $mZ6/6 = 0.5046$ MeV (provenance unverified); CEC inversion 0.5069 MeV (provenance unverified). The two numerical routes sit between the two exact branches, ~1% from each — inconclusive at current precision. CEC granularity: the inversion evidence equals $0.968\cdot me^4$ vs branch predictions $1.000\cdot me^4$ (A) and $0.912\cdot me^4$ (B) — a slight lean toward branch A; the P18 route sits dead-centre. Decisive audits registered for the owner: Q1 — provenance of the §3.2 normalisation (decides the branch); Q2 — provenance of mZ6 in the P18 table; Q3 — provenance of fplasma = 18 mHz and ρcond. The branch bookkeeping and the ηB quarantine are carried in §J.11.

L.3 Registered Observations

12α expansion. $1 - \mathrm{sinc}^2(\pi/Nc) \approx 0.08811$ vs $12\alpha \approx 0.08757$ (0.6%); if exact, $\alpha = (4\pi^2 - 36)/(48\pi^2) \approx 1/136.19$ — better than Eddington's 1/136, not a derivation (Wyler-class caution registered); using $\kappa/G = 1 - 12\alpha$ in the pass criterion worsens ηB from percent-level to 8.8% — the coincidence is an observation, not a route.

Chord/arc/sagitta geometry. $\sqrt{(\kappa/G)}$ = chord/arc of a Z6 sector (half-angle 30°); the circle's own $\sin^2 + \cos^2 = 1$ supplies the Pythagorean relation (semi-chord, R − sagitta, R); κ lives in the flat lattice metric, G in the curved phase-circle metric. Nyquist reading: sinc is the universal sampling form factor — the Z6 condensate literally samples the massless psion's phase at Nc = 6 points, and the P23 deficit is its sampling error; the geometry explains the gap completely and closes it only as $Nc \to \infty$, which Nc = 6 [T1] forbids.

L.4 Charge-Conjugation Audit and the Lepton-Generation Mechanism

L.4.1 Composite identification (proton, neutron, exotic). Proton is LRR $(0,4,5)$, $Q = 1/9+4/9+4/9 = +1$; neutron is LLR $(0,1,5)$, $Q = 1/9-5/9+4/9 = 0$; RRR $(3,4,5)$ is a separate $Q = +2$ exotic BSM state. Both proton and neutron charges hold exactly under Theorem V.1 alone, no auxiliary scheme required. (The companion repository's `CHARGE_CONVENTIONS.md` is kept consistent with this identification; see its §3a.)

L.4.2 State-wise charge conjugation and Theorem V.1 (T1, label-independent). Conjugation is the index map $k \to (6-k)$ mod 6, fixing $k = 0,3$. Requiring $Q_k = -Q_{\mathrm{conj}(k)}$ state-wise (not merely $\Sigma_k Q_k = 0$) forces $Q_0 = Q_3 = 0$ and $Q_1 = -Q_5$, $Q_2 = -Q_4$. Theorem V.1's values ($Q_0 = 1/9$, $Q_1 = -5/9$, $Q_2 = -5/9$, $Q_3 = 10/9$, $Q_4 = 4/9$, $Q_5 = 4/9$) satisfy none of these four constraints; the $(3,4,5)$ exotic state makes this concrete: V.1 gives $Q = +2$, while any conjugation-consistent assignment would give $Q = +1$ for the same triple — a full unit apart. This is a property of the six Q_k values themselves, independent of which composite is called proton or neutron.

L.4.3 Adopted reading of C/CP. Three repairs were evaluated: (i) retune the six Q_k to satisfy state-wise conjugation exactly — compatible with proton $(+1)$/neutron (0) above, but changes the $(3,4,5)$ exotic charge from $+2$ to $+1$, a value not adopted in this revision; (ii) extend the conjugation map to include proto-colour — any involution on 3 proto-colours has an odd fixed-point count, so this cannot remove both fixed points, and the one full involution tested trades the lepton-generation charge count for an unreachable neutron neutrality; (iii) retain conjugation only as an aggregate identity over the composite spectrum ($\Sigma_k Q_k$ tracked as an ensemble quantity), not as an exact per-state operator. Option (iii) is adopted: it requires no change to any Q_k value, and §13.1(2)'s $\varepsilon\psi$-based C/CP mechanism (footnote, §13.1) is unaffected by it, since that mechanism never invokes state-wise Q_k-conjugation.

L.4.4 Dual-sector labelling and the μ^-/τ^- mechanism. §6 and Appendix H §H.3 both label LLR $(0,1,5)$, $(0,2,4)$, and $(1,2,3)$ as dual-sector (neutrino-like or baryon-like: n/ν_τ, ν_μ/Σ^0-like, ν_e/Λ^0-like), with generation ordering for the third explicitly open. Theorem 6.2 ([T1]: exactly one $N = 3$ configuration carries $Q = -1$) fixes the scope of this ambiguity precisely: it is neutrino-vs-baryon-like only, never charged-lepton-vs-baryon-like, since no $N = 3$ triple other than $(0,1,2)$ can carry $Q = -1$.

The combinatorial identity of μ^-, τ^- is therefore not an $N = 3$ tuple at all, and is instead addressed by the Leptonic Transfer Matrix (P8b, Appendix A.2.5): $t_{12} = -8$ (exact, [T1]); $\langle 3|H_{dd}|1\rangle = t_{12}^2 = 8$ (three-body symmetry cancellation, exact, [T1]); diagonal set $C_2 = \{10.75, 8.75, 6.75\}$ (exhaustive $Q = -1$-sector enumeration, [T1]) — eigenvalues of a mixing operator on the lepton sector, mapped to $m_e : m_\mu : m_\tau$, rather than three separate psion composites. $N_{\mathrm{gen}} = 3$ ([T1], §9, kinematic bound $J_R = 7/2 > j_{\max} = 5/2$) is consistent with this construction and bounds it from above; $m_\mu = 105.66$ MeV additionally appears once (§K.8, RCF formula) as an external CODATA reference, not a P8b-derived value.

L.4.6 Dual sector and nucleon stability (v86). If the same triple is a baryon or a neutrino according to its cascade level, and the level carries no internal quantum number, then $n \to \nu_\tau\gamma$ and its analogues are allowed by every symmetry of eq. (1); stability requires an amplitude suppression $\lesssim 5 \times 10^{-30}$. The tension recorded in L.4.4 is therefore a stability problem, resolved only by a conserved baryon (rung) label (§13.4, P14-B).

L.4.7 Hadron size: the adopted ring geometry and its limits (v86, P14-C, [T4]). *Adopted description.* For the charge radii of baryon-type composites the paper adopts the planar Z6 ring as a phenomenological description: psion k at angle $60°k$ on a circle, equal constituent masses, and $\langle r^2\rangle = \sum_k Q_k|\mathbf{x}_k - \mathbf{c}|^2/\sum_k Q_k$ about the centroid (not normalised when $\sum_k Q_k = 0$), with one length fixed by the proton ($r_p = 0.841$ fm, ring radius $a = 1.212$ fm). It is a description, not a derivation from eq. (1), and it was chosen after a pre-registered series of alternatives (companion repository, WP-GEO to WP-GEO12, WP-HUB, WP-PHS, WP-REL).

What it encodes. For three consecutive labels $(c-1, c, c+1)$ on the ring the mean-square charge radius is the

exact identity $\langle r^2\rangle = a^2\left[\frac{7}{9}Q_{\text{tot}} - \frac{2}{3}Q_c\right]$. The radii of arc-type triples therefore depend only on cyclic adjacency in Z6 and on the charge of the central label; the star, petal and 12-gon arrangements give identical ratios, and of the 720 orderings of the labels on the ring only the natural cyclic order passes. A neutral triple of mutually non-adjacent labels is an equilateral triangle and has $\langle r^2\rangle = 0$ exactly. The neutron radius is negative because the only positive label sits at the centre and the net charge at the rim is negative.

State	Triple	Ring $\langle r^2\rangle/\langle r^2\rangle_p$	Reference	Status
n	$(0,1,5)$	-0.154	-0.164 (measured)	pass (pre-registered)
Σ^-	$(0,1,2)$	$+0.846$	0.86 ± 0.21 (measured)	see below
Σ^0	$(0,2,4)$	0 (exact)	-0.04 to $+0.10$ (theory)	consistent with zero
Λ	$(1,2,3)$	$+0.77$	$+0.06$ to $+0.15$ (theory)	fails
Σ^+	$(1,3,5)$ or $(2,3,4)$	$+2.08$ or $+0.08$	$+0.84$ to $+1.11$ (theory)	fails
exotic	$(3,4,5)$	$+1.31$	none observed	open prediction

No radius of Σ^+, Σ^0 or Λ has been measured; the references are chiral-perturbation-theory and lattice results [22,23].

Identification caveat. The $J\cdot\Sigma$ term couples left to right psions only, so a triple of one chirality has no $J\cdot\Sigma$ bond. The triple $(0,1,2)$ is then inert, which is the exact statement used for the electron (§6, L.4.4). The Σ^- row treats $(0,1,2)$ as a hadron and is meaningful only if the electron triple and the Σ^- are distinct states (different cascade levels, P14-B); it is reported but not counted as an independent success. A model in which the $J\cdot\Sigma$ bond graph sets the size (a V with the unique-chirality psion as hub, one opening angle fitted to the neutron) makes $(0,1,2)$ pointlike and fails Σ^0, Λ and Σ^+.

Alternatives tested and not adopted. Line, centred hexagon, blind triangle, prism, spin cone, hourglass, three-triangle star, and a 12-site ring in which $k = 3$ is displaced by 30° (it fits Λ and Σ^+ by one discrete choice, two of seven positions, with no dynamical origin); the arc and circular-mean metrics of the internal phase circle (neutron -0.200). Magnetic moments fail in every point-charge-with-spin rule tested. Neutrino radii cannot be made to vanish together by any placement of the six labels on 12 sites (best $\max|R| = 0.056$), so the smallness of the neutrino and of the charged leptons must come from scale, not shape.

Limits that remain. (a) *Mixing.* $J\cdot\Sigma$ conserves $\sum k$ and mixes n, Σ^0, Λ (and p with the two Σ^+ candidates); the ring assumes no mixing. With adiabatic labels the neutron radius turns positive for $g \gtrsim 0.06$ and, taking p and n as the lowest states of their sectors, $m_n < m_p$ at every coupling tested. (b) *Scale.* The number $\hbar c/a \approx 163$ MeV is not an energy of the theory: it is r_p converted through the premises that $p = (0,4,5)$, the constituents are classical points of equal mass on one ring radius, and $E = \hbar c/a$ (that is, $\Delta t = a/c$). A massless-fermion confinement correction ($E = x\hbar c/R$, $x = 2.04$) raises the energy per psion to 333–349 MeV and gives $M_p = 3E$ within $+6\%$ and $+12\%$, but this holds for any three massless constituents in about 1.2 fm, and a common orbital gives $R_n = 0$, so the correction can apply only to the length–energy relation. No route available in the paper derives the scale: the $J\cdot\Sigma$ contact term needs tuning of one part in 10^{24}; proto-colour running needs $\alpha_{\text{pc}}(\Lambda_{\text{meta}}) \approx 1/50$ while every reading in the paper gives $\alpha_{\text{pc}} \geq 0.09$ (the induced value $3\pi/(S/2g^2) = 0.58$ agrees with $\alpha_{\text{meta}} = \pi/6$ to 11%, placing proto-colour confinement near Λ_{meta}); four pre-registered exponent forms and the cascade rungs miss by factors 3.7 and larger ($\ell = 8$ gives $r_p = 0.22$ fm, $\ell = 8.5$ gives 1.32 fm); a core bound at Λ_{meta} is excluded by the proton form factor by 24 orders of magnitude. (c) *Gottfried sum.* The valence part with the charges Q_k is $-1/9$ (quarks $+1/3$), so the sea must supply $+0.346$ to reach the measured $+0.235 \pm 0.026$, 3.5 times the QCD sea term; a closed list of cloud channels reaches it only for a cloud probability between 0.09 and 0.89, which the PH does not fix, so the proton identification is not excluded.

This is open problem **P14-C**: a derived baryon scale, a sea model, and an internal structure consistent with the $J\cdot\Sigma$ dynamics. The ring is adopted as the working description meanwhile; its failures for Λ and Σ^+ are

the first falsifiers, and nothing in the main text depends on this appendix.

L.4.8 Dynamic ring and hybrid description: demoted to a negative result (v86, P14-C). *What was tested.* The static ring of L.4.7 was replaced by a quantum ring with N_s sites ($N_s = 6$, 8, 12; also a $\mathbb{Z}_3 \times \mathbb{Z}_2$ prism and a twisted prism), the three psions of a triple occupying distinct sites, with $H = A + \kappa B + \lambda C - tT - sS$: A is a clock term (alignment of each psion with its natural site), $B = \sum q_a q_b / d_{ab}$ the Coulomb sum, $C = -\sum 1/d_{ab}$ over left–right pairs the $J\cdot\Sigma$ term, T hops of $\pm 30^\circ, \pm 60^\circ$ (12 sites) and S chirality-exchanging swaps and antipodal jumps. Radii are ground-state expectations normalised to the proton; masses are linear functionals $a\langle A\rangle + b\langle B\rangle + g\langle C\rangle$ with one common direction of a single sign, the scale fixed by $\Lambda - n$ with $\varepsilon \leq 349$ MeV and weights ≤ 1.5. All tests were pre-registered (companion repository, WP-DYN12 to WP-CND); each registry records the anticipations that proved wrong.

Negative results (they stand). (i) With $\kappa, \lambda \geq 0$ the ground-state energies give $m_n < m_p$ and $m_\Lambda < m_n$ at all of 12000 points; the sign follows from the charge assignment (the Coulomb energy of p is repulsive, that of n attractive), not from an exclusion or uncertainty principle. (ii) On the 12-site ring, radii and a natural mass scale compete: the best radius loss is 0.79 at mass reach $q \approx 0.02$ (scale 30–65 times too large), ≥ 1.08 at $q = 0.24$ and ≥ 1.5 at $q = 0.34$ (loss below 1 is viable). (iii) The prism is degenerate in its angular part, because each of the eight unconfined triples contains exactly one psion of each $\mathbb{Z}_3$ class; the 8-site ring and the twisted prism do not improve on the 12-site ring. (iv) Free weights make the mass criterion nearly vacuous: random mass targets pass at 36–64% of points.

A hybrid description (demoted). If p and n are static and every triple containing psion 2 or 3 (the bonded pair of J.6.3) is dynamic, the mass is essentially the clock energy: delocalisation costs energy, so Λ, Σ^0 and Σ^+ lie above n and p with the right sign. With Σ^- left out of the optimisation, 3 of 10 robust searches satisfy all radii (n is fixed at -0.154 by the arc identity), the Σ^0 and Σ^+ mass ratios and the natural scale, and survive $\pm 10\%$ parameter perturbations. Predicted masses relative to n are 231 MeV for Σ^0 (observed 253) and 179–190 MeV for Σ^+ (observed 250).

Out-of-sample test and its limits. The first test (three optima) gave one pass of three for the Σ^- (radius and mass) and the charge-2 triple $(3,4,5)$. A rerun with 10 independent searches (static proton, frozen windows) found 8 viable optima, of which only 1 passes (fraction 0.12 against a registered criterion of 0.6): the earlier 1/3 was a small-sample overestimate. Within the region already satisfying radii and mass ratios, passing both predictions occurs at 0.9% of points. The weaknesses were never removed: about 30 models were tried before and during this series and the look-elsewhere effect is not corrected, the rule was chosen after the simpler one failed, two mass ratios sit at the edge of their windows, the Σ^- is the inert electron triple (L.4.4), the comparison of $(3,4,5)$ with the Δ^{++} is unverified and a spinless model cannot give spin $3/2$, and the Σ^0–Λ splitting is a spin–flavour effect. A coarse $\Sigma^- - \Sigma^0$ window (0–50 MeV against 4.8 MeV observed) was met by the best optimum, but it is not independent of the Σ^- criterion and does not count as support.

Different geometries for different composites. Taking each composite's simplest geometry that fits its radius gives the hexagon (static) for n and Σ^0, the octagon (static) for $\Sigma^+(1,3,5)$ and a delocalised state for Λ and $\Sigma^+(2,3,4)$; in the dynamic geometries a single window is reached trivially because the parameters are free per composite. This is a description of possibilities, not an explanation. No simple rule on the triple predicts the class (leave-one-out 2/5, permutation $p = 0.47$). Adding the mass criterion does not choose between the mixed assignment and the all-dynamic one: both remain viable (8–10 of 10 searches), and the independent Σ^- and $(3,4,5)$ predictions pass in 0 of 18 mixed-geometry optima and 1 of 8 hybrid optima. Only three charge radii are measured (p, n and Σ^-, the last at 0.61 ± 0.21 fm^2 by SELEX), so radii cannot test the question further. *Spin extension.* A minimal extension gives each psion spin $1/2$, couples all left–right pairs by $J\,\mathbf{s}_i\cdot\mathbf{s}_j$ and takes the moment proportional to the psion charge Q_9, with no free parameter (WP-SPIN, pre-registered). The ratios to the proton are $\mu_n/\mu_p = -0.80$, $\mu_\Lambda/\mu_p = -2.0$, $\mu_{\Sigma^+}/\mu_p = +2.2$ for $J > 0$ (observed -0.685, -0.219, $+0.880$) and 0, 0, $+1.0$ for $J < 0$; the criterion (deviation ≤ 0.2 in all three) fails for both signs, and

12 of 720 permutations of Q_9 over the labels pass. The Σ^- and $(3,4,5)$ have no left–right pair, so their spin is undetermined and the Δ^{++} spin $3/2$ is not derived.

Status. Closure of the series: negative result for P14-C, not derived. The hybrid description is no longer proposed as a candidate; it is kept only to document what was tried, and no prediction of the main text depends on it. The ring model with the postulates tried (radii, mass, geometry per composite, minimal spin) does not derive the baryon scale or internal structure; a further test would need a new postulate chosen before looking at the data.

L.4.5 Precision, Table 5 vs. §A.3.1. Table 5 (§12) reports the leading-order transfer-matrix ratios, $m_\mu/m_e = 218$ (5.4% vs. 206.8) and $m_\tau/m_e = 3795$ (9.1% vs. 3477), explicitly captioned as targeting further refinement via P8b. §A.3.1 reports the ratios after the P8b correction ($\Sigma_2 = b_e \times 15/8$) at $+0.46\%$ and $+0.56\%$ respectively. The two are sequential stages of the same calculation — leading order, then P8b-corrected — not independent estimates; the P8b formal-integration step itself remains at [T2] (open-problems registry, App. A), so the 0.46%/0.56% figures carry that status pending closure of Passo A.3, while the 5.4%/9.1% leading-order figures are unconditional.

L.5 v85 Revision Record (October 2026)

This revision is an audit, not a new physical result. Every change below is traceable to a commit in the companion repository [2]; negative findings are recorded with the same weight as positive ones.

L.5.1 Dark-matter census proof. The module `proofs/P_dark_matter.py` tested stability by single-psion removal, a test that can never fire because no psion is neutral, and failed its own assertion. It now uses the fragmentation criterion (no proper sub-multiset is a neutral singlet) with two independent algorithms; Theorem DM.1 is reproduced unchanged (39 states, $N \in \{9, 12, 15, 18, 21, 27\}$), and N = 6 is confirmed fully unstable.

L.5.2 Dark-matter mass and cascade level. Three incompatible dark-matter masses coexisted (Table 3, $m(N) \leq 17.6$ MeV; §14.2, 7.2×10^{10} GeV; a repository body-number law, 14.3 GeV). Because h_{AB} and X17 share N = 2 but sit at cascade levels 8 and 9, the cascade level is not a function of body number; the Landau-pole mass of §14.2 is canonical, the 14.3 GeV result is retracted, and Table 3 is relabelled by cascade level (§8.1).

L.5.3 Theorem KS. Inequality (11) holds as arithmetic on the mass function (verified on all 1,295,920 partitions of every level up to 50), but physical fragmentation partitions body number, not level. Its physical application and Corollary 8.2 are suspended until a rule fixes the levels of fragments (§8.2). Dark-matter stability rests on DM.1 and psion-number conservation instead. Three k = 3 entries of Table 3 were wrong in v84 ($\ell = 2$: no three-part partition exists; $\ell = 8$ and 10 overstated); they are corrected.

L.5.4 F5 (direct detection). The bound $\sigma < 10^{-50}$ cm^2 had no derivation and is retired. F5a (no dark-matter state outside the 39 singlets at m_{DM}; [T2], E2) carries the falsification power; F5b (rate $\sim 10^{-54}$ cm^2, conditional on zero proto-colour coupling; [T4], E1) records the estimate. A kill criterion was registered ex ante (§9.1); the LZ event of September 2026 is below it.

L.5.5 F3 and the PMNS programme. F3 is a function of the PMNS matrix alone, which the PH does not derive, and the 7% figure has no derivation in this release; F3 is carried at [T4], conditional on P-PMNS (§17.3). The companion repository opens a PMNS programme with pre-registered models: the flavour basis is fixed exactly (generation = proto-colour of the right-handed constituent; colour-diagonal charged current), and with charged leptons diagonal both terms of eq. (1) and the locking gap give $\sin^2\theta_{12}$ near the measured value but $\sin^2\theta_{23} \approx 0.21$. The charged-lepton side is blocked: the App. A.2.5 entries as printed do not reproduce the P8b mass ratios claimed in §A.3.1 (best $m_\mu/m_e = 244$ when m_τ/m_e is matched), so the full P8b matrix is required.

L.5.6 Unchanged. No central numerical prediction of Classes A or B changes in this revision.

L.6 v86 Revision Record: the P14 Audit (October 2026)

All work packages were pre-registered, with SHA-256 and a UTC timestamp, before execution. Failures and corrections are recorded with the same weight as results; the full log is in `p14d_psion_number_audit/README.md` of repository [2].

L.6.1 Changes. §13 rewritten (13.1–13.7); η_B derivation withdrawn; F12 suspended; App. J.1 and J.4 annotated; App. A.4 antimatter statement corrected (antipsions); §8.2 status note: the missing level rule is a conserved baryon label (P14-B); §14.3 direct route marked unsupported; open-problem tables gain P14-B and P14-C; App. L.4.7 added (hadron-size geometry, phenomenological); App. L.4.8 added (dynamic-ring series: negative results, including the demotion of a hybrid description); abstract, §4.1, §12 and §15–19 summaries updated accordingly. No other numerical prediction changes.

L.6.2 Errors made during the audit and corrected. (i) Sphalerons were first treated as external to the PH; they are psion composites and conserve N_ψ. (ii) Antimatter was first modelled as antipsions, then switched to index-inverted psion triples on a reading of A.5, and restored to antipsions by the reaction test of 13.1; the second step was the wrong one. (iii) In the domain test, unitarity was first imposed jointly across processes with different density scaling, which created asymmetry in equilibrium. A control caught this, the test was corrected, and the faulty version is retained. (iv) In the calibration test, the dark-matter block first patched a copy of the module globals and was redone in an addendum. (v) The anticipated count of neutral triples was wrong (it is five). (vi) In the bifurcation test, the registered endpoint bookkeeping had a sign error, which was corrected (two antineutrinos per baryon), and the anticipated efficiency was wrong by about three orders of magnitude. (vii) The bifurcation networks first used free psions below Λ_{meta}, which App. I forbids; the mechanism was reformulated as pair production of unconfined composites. (viii) Rung assignments in two tests contradicted the stability requirement for charged leptonic copies; corrected, they make the sign of η_B law-dependent. (ix) A test of the vacuum phase counted the condensate-generated bond twice, as an independent real term; without the double counting the phase is removable by $U(1)_M$, and the claimed CP source does not exist. (x) The Gottfried sum was first compared in its valence part alone with the measured total, which includes the sea, and reported as a conflict with deep-inelastic data; the sea was then enumerated (WP-GOT), the registered anticipation (failure) was wrong for the best channel, and the statement was reduced to an open requirement on the sea. (xi) In the dynamic-ring series (L.4.8) several registered anticipations were wrong: the energy-selection mechanisms were expected to have a feasible region; the dodecagon was expected to fail on radii alone, which it does not; the scale was expected to be the culprit of the radius–mass tension where the Σ^0 ratio was; and the out-of-sample baseline was expected to be common where it was rare. The first out-of-sample result (1 of 3 optima) was a small-sample overestimate: 1 of 8 after ten searches; the hybrid was demoted accordingly. A mixed-geometry model was expected to fail on mass and did not, and a radii-normalisation error (dynamic instead of static proton) was found and corrected in WP-MIX; in WP-SPIN the null fraction was expected at $\geq 2\%$ and came out at 1.7%. All are recorded in the companion registries. (xiii) The Sec. 8.1 ladder had never been compared with the hadron ground states; the registered test (WP-KILL) found no alignment, and the anticipation that the Σ^- radius would miss the 1σ band was wrong (it sits at 0.05σ). (xii) K.10 was left unannotated against the statement that $U(1)_M$ is exact; the inconsistency is now noted in K.10 and remains open.

L.6.3 Registered control failures. All are at or near the floating-point floor or due to integrator limits: assembly-network controls (2×10^{-13} against 10^{-15}); temperature-programme step convergence (1–2%); domain test C3 (2.7×10^{-6}) and C4 (1.3×10^{-12}); phase-in-$J\cdot\Sigma$ test C3 (1.8×10^{-6}) and one sensitivity run not completed; bifurcation tests, $B + L = 0$ at the numerical floor ($\leq 1.1 \times 10^{-8}$).

References

[1] Planck Collaboration, Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641, A6 (2020). arXiv:1807.06209.

[2] K. M. Maia, psion-proofs: Verification repository for the Psion Hypothesis, GitHub, `https://github.com/Maia-KarlosMarden/psion-proofs` (2026).

[3] A. D. Sakharov, Vacuum quantum fluctuations in curved space, Sov. Phys. Dokl. 12, 1040 (1967).

[4] Y. Nambu and G. Jona-Lasinio, Dynamical model of elementary particles based on an analogy with superconductivity, Phys. Rev. 122, 345 (1961).

[5] S. Weinberg and E. Witten, Limits on massless particles, Phys. Lett. B 96, 59 (1980).

[6] I. M. Gel'fand and A. M. Yaglom, Integration in functional spaces and its applications in quantum physics, J. Math. Phys. 1, 48 (1960).

[7] R. Jackiw and C. Rebbi, Solitons with fermion number 1/2, Phys. Rev. D 13, 3398 (1976).

[8] A. J. Krasznahorkay et al., Observation of anomalous internal pair creation in ^{8}Be, Phys. Rev. Lett. 116, 042501 (2016).

[9] MiniBooNE Collaboration, Significant excess of electronlike events in the MiniBooNE short-baseline neutrino experiment, Phys. Rev. Lett. 121, 221801 (2018).

[10] MicroBooNE Collaboration, Search for an excess of electron neutrino interactions in MicroBooNE using multiple final-state topologies, Phys. Rev. Lett. 128, 241801 (2022).

[11] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, CODATA recommended values of the fundamental physical constants: 2018, Rev. Mod. Phys. 93, 025010 (2021).

[12] K. M. Maia, Derivation of the RGD Kernel from the PH Lagrangian, v2 (2026). [companion paper]

[13] K. M. Maia, Dark Matter: Existence Without Necessity (2026). [companion paper]

[14] MicroBooNE Collaboration, Search for an anomalous excess of charged-current quasielastic νe interactions with the MicroBooNE experiment using Deep-Learning-based reconstruction, Phys. Rev. D 105, 112003 (2022).

[15] MicroBooNE Collaboration, Search for an anomalous excess of charged-current νe interactions without pions in the final state with the MicroBooNE experiment, Phys. Rev. D 105, 112004 (2022).

[16] A. D. Sakharov, Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe, JETP Lett. 5, 24 (1967).

[17] Super-Kamiokande Collaboration, Search for nucleon decay via $n \to \bar{\nu}\pi^0$ and $p \to \bar{\nu}\pi^+$, arXiv:1305.4391.

[18] KamLAND Collaboration, Search for the invisible decay of neutrons with KamLAND, Phys. Rev. Lett. 96, 101802 (2006).

[19] A. G. Cohen, A. De Rújula, and S. L. Glashow, A matter–antimatter universe?, Astrophys. J. 495, 539 (1998).

[20] E. W. Kolb and S. Wolfram, Baryon number generation in the early universe, Nucl. Phys. B 172, 224 (1980).

[21] E. Witten, Current algebra, baryons, and quark confinement, Nucl. Phys. B 223, 433 (1983).

[22] B. Kubis and U.-G. Meißner, Baryon form factors in chiral perturbation theory, arXiv:hep-ph/0010283.

[23] Boinepalli et al., Precision electromagnetic structure of octet baryons in the chiral regime, arXiv:hep-lat/0604022.