Erasmus School of Economics

MOOC Econometrics


Lecture 4.2 on Endogeneity: Consequences

Dennis Fok

Erasmus University Rotterdam

Ezafus

Simulated example, $y = 1 + 2x^* + u$



Endogeneity

- Common problem in economics
 - Omitted variables
 - Strategic behavior
 - Measurement errors
 - $\rightarrow X$ is correlated with ε
- Endogeneity violates the basic assumptions
- \rightarrow How bad is this?

Ezafus,

Lecture 4.2, Slide 2 of 12, Erasmus School of Economics

Measurement error example

Under measurement error (and endogeneity in general):

• we obtain the wrong coefficients!

Test

Can we say anything about the direction of the bias?

Erafus

Direction of bias in the measurement error case

OLS is "biased towards zero"

ightarrow OLS underestimates true effect

Intuitively:

- x-values on the *left* likely have negative measurement errors
- x-values on the *right* likely have positive measurement errors

Measurement errors "stretch" the scatter in the horizontal direction \rightarrow a flatter regression line

(Zafing

Lecture 4.2, Slide 5 of 12, Erasmus School of Economics

Consistency: formal argumentation

If X is endogenous:

- If *n* grows the OLS estimator converges to the wrong value.
 - \rightarrow OLS is inconsistent

Consider the standard model $y = X\beta + \varepsilon$ and the OLS estimator


$$b = (X'X)^{-1}X'y = (X'X)^{-1}X'(X\beta + \varepsilon)$$
$$= (X'X)^{-1}X'X\beta + (X'X)^{-1}X'\varepsilon$$
$$= \beta + (X'X)^{-1}X'\varepsilon$$

So, b can be split into

- **1** True parameter value β
- 2 Random deviation $(X'X)^{-1}X'\varepsilon$

Lahm

Distribution of estimator for different n, true value= 2



Lecture 4.2, Slide 6 of 12, Erasmus School of Economics

Asymptotic properties

What happens to *b* as $n \to \infty$?

Recall: $b = \beta + (X'X)^{-1}X'\varepsilon$

- ullet β is constant
- Elements of (X'X) and $X'\varepsilon$ are sums over observations:

$$X'X = \begin{pmatrix} \sum_{i=1}^{n} x_{1i}^{2} & \sum_{i=1}^{n} x_{1i}x_{2i} & \dots & \sum_{i=1}^{n} x_{1i}x_{ki} \\ \sum_{i=1}^{n} x_{1i}x_{2i} & \sum_{i=1}^{n} x_{2i}^{2} & \dots & \sum_{i=1}^{n} x_{2i}x_{ki} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^{n} x_{ki}x_{1i} & \sum_{i=1}^{n} x_{ki}x_{2i} & \dots & \sum_{i=1}^{n} x_{ki}^{2} \end{pmatrix}, X'\varepsilon = \begin{pmatrix} \sum_{i=1}^{n} x_{1i}\varepsilon_{i} \\ \sum_{i=1}^{n} x_{2i}\varepsilon_{i} \\ \vdots \\ \sum_{i=1}^{n} x_{ki}\varepsilon_{i} \end{pmatrix}$$

 \rightarrow these diverge as $n \rightarrow \infty$

Asymptotic properties

Rewrite $b = \beta + (\frac{1}{n}X'X)^{-1}(\frac{1}{n}X'\varepsilon)$

- $(\frac{1}{n}X'X)$ is an average \rightarrow in general converges to, say, Q
- $(\frac{1}{n}X'\varepsilon)$ also converges in general

Consistency result:

b converges to β as $n \to \infty$ if

- \bullet $\frac{1}{n}X'X$ converges to Q, and
- Q^{-1} exists, and
- **3** $\frac{1}{n}X'\varepsilon$ converges to 0
 - ▶ No correlation between X and ε (for large n)
 - ► *X* is exogenous

X endogenous: b does not converge to β !

- Eafins

Lecture 4.2, Slide 9 of 12, Erasmus School of Economics

OLS in presence of endogeneity

If X endogenous

- X correlated with ε
- ullet OLS estimator for eta is not consistent
- Even with in infinite amount of data: OLS does not give useful estimates

Small sample properties

So far we discussed what happens for $n \to \infty$

Test

Why can't we derive the bias?

To obtain the bias

need to evaluate

$$E[b] = E[(X'X)^{-1}X'y] = E[(X'X)^{-1}X'(X\beta + \varepsilon)]$$
$$= E[\beta + (X'X)^{-1}X'\varepsilon] = \beta + \underbrace{E[(X'X)^{-1}X'\varepsilon]}_{=?}.$$

- X is stochastic
- cannot simplify final expectation (without further assumptions)

(Zafus

Lecture 4.2, Slide 10 of 12, Erasmus School of Economics

TRAINING EXERCISE 4.2

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

(zafus