

CS499 Homework 3 (First Draft)

Interstellar

Exercise 3.1

1.

For $T_{10} = n$, there are $n - 1$ situations for the first $n - 2$ digits of the string:

$$\begin{cases} 0, 0, 0, \dots, 0, 0 \\ 0, 0, 0, \dots, 0, 1 \\ \vdots \\ 1, 1, 1, \dots, 1, 1 \end{cases}$$

Based on the probability calculation formula of classical probabilities, we have

$$Pr [T_{10} = n] = \frac{n - 1}{2^{n-2}} \cdot \frac{1}{4} = \frac{n - 1}{2^n}$$

2.

Based on the knowledge about A_n , we have

$$Pr [T_{11} = n] = \frac{|A_{n-2}|}{2^{n-2}} \cdot \frac{1}{4} = \frac{|F_n|}{2^n}$$

3.

Based on method 3, we have

$$\begin{cases} E(T_{10}) = 1 + \frac{1}{2} \cdot E(T_{10}) + \frac{1}{2} \cdot E(T_0) \\ E(T_0) = 1 + \frac{1}{2} E(T_0) + \frac{1}{2} \cdot 0 \end{cases}$$

$$\Rightarrow E(T_{10}) = 4.$$

$$E(T_{11}) = 1 + \frac{1}{2} \cdot E(T_{11}) + \frac{1}{2} \cdot [1 + \frac{1}{2} \cdot E(T_{11})]$$

$$\Rightarrow E(T_{11}) = 6.$$

Exercise 3.2

There are 2 states of the coin, which are 0 and 1. Assuming the probability of the event under each state is P_1 and P_2 respectively, we have

$$\Rightarrow \begin{cases} P_1 = \frac{1}{2} P_1 + \frac{1}{2} P_2 \\ P_2 = \frac{1}{2} \times 0 + \frac{1}{2} \times 1 \end{cases}$$

$$\Rightarrow \begin{cases} P_1 = \frac{1}{2} \\ P_2 = \frac{1}{2} \end{cases}$$

$$\Rightarrow Pr[\varepsilon] = \frac{1}{2} (P_1 + P_2) = \frac{1}{2}$$

Exercise 3.3

1. There are 4 states of last bit pair of $\{x\}$ and $\{y\}$ sequence, which are $\{0, 0\}, \{1, 0\}, \{0, 1\}$ and $\{1, 1\}$. Assuming the expectation of T under each state is E_1, E_2, E_3, E_4 respectively, we have

$$\begin{cases} E_1 = 1 + \frac{1}{4}E_1 + \frac{1}{4}E_2 + \frac{1}{4}E_3 + \frac{1}{4}E_4 \\ E_2 = 1 + \frac{1}{4}E_2 + \frac{1}{4}E_4 \\ E_3 = 1 + \frac{1}{4}E_1 + \frac{1}{4}E_2 \\ E_4 = 1 + \frac{1}{4}E_2 \end{cases}$$

$$\Rightarrow \begin{cases} E_1 = \frac{384}{121} \\ E_2 = \frac{20}{11} \\ E_3 = \frac{272}{121} \\ E_4 = \frac{16}{11} \end{cases}$$

$$\Rightarrow E[T] = \frac{1}{4}(E_1 + E_2 + E_3 + E_4) = \frac{384}{121}$$

2.

(a) Similar to the solution above, assuming the probability of the event "10 appears in x before 11 appears in y " under each state is $Pra_1, Pra_2, Pra_3, Pra_4$, we have




Figure 1: The state machine

$$\begin{cases} Pra_1 = \frac{1}{4}Pra_1 + \frac{1}{4}Pra_2 + \frac{1}{4}Pra_3 + \frac{1}{4}Pra_4 \\ Pra_2 = \frac{1}{2} + \frac{1}{4}Pra_2 + \frac{1}{4}Pra_4 \\ Pra_3 = \frac{1}{4}Pra_1 + \frac{1}{4}Pra_2 \\ Pra_4 = \frac{1}{4} + \frac{1}{4}Pra_2 \end{cases}$$

$$\Rightarrow \begin{cases} Pra_1 = \frac{65}{121} \\ Pra_2 = \frac{41}{121} \\ Pra_3 = \frac{9}{11} \\ Pra_4 = \frac{5}{11} \end{cases}$$

$$\Rightarrow Pra = \frac{1}{4}(Pra_1 + Pra_2 + Pra_3 + Pra_4) = \frac{65}{121}$$

(b) Similar to the solution above, assuming the probability of the event "10 appears in x at the same time with 11 appears in y" under each state is $Prb_1, Prb_2, Prb_3, Prb_4$, we have

$$\begin{cases} Prb_1 = \frac{1}{4}Prb_1 + \frac{1}{4}Prb_2 + \frac{1}{4}Prb_3 + \frac{1}{4}Prb_4 \\ Prb_2 = \frac{1}{4}Prb_2 + \frac{1}{4}Prb_4 \\ Prb_3 = \frac{1}{4}Prb_1 + \frac{1}{4}Prb_2 \\ Prb_4 = \frac{1}{4} + \frac{1}{4}Prb_2 \end{cases}$$

$$\Rightarrow \begin{cases} Prb_1 = \frac{17}{121} \\ Prb_2 = \frac{7}{121} \\ Prb_3 = \frac{1}{11} \\ Prb_4 = \frac{3}{11} \end{cases}$$

$$\Rightarrow Prb = \frac{1}{4}(Prb_1 + Prb_2 + Prb_3 + Prb_4) = \frac{17}{121}$$

(c) Similar to the solution above, assuming the probability of the event "10 appears in x after 11 appears in y" under each state is $Prc_1, Prc_2, Prc_3, Prc_4$, we have

$$\begin{cases} Prc_1 = \frac{1}{4}Prc_1 + \frac{1}{4}Prc_2 + \frac{1}{4}Prc_3 + \frac{1}{4}Prc_4 \\ Prc_2 = \frac{1}{4}Prc_2 + \frac{1}{4}Prc_4 \\ Prc_3 = \frac{1}{2} + \frac{1}{4}Prc_1 + \frac{1}{4}Prc_2 \\ Prc_4 = \frac{1}{4} + \frac{1}{4}Prc_2 \end{cases}$$

$$\Rightarrow \begin{cases} Prc_1 = \frac{39}{121} \\ Prc_2 = \frac{73}{121} \\ Prc_3 = \frac{1}{11} \\ Prc_4 = \frac{3}{11} \end{cases}$$

$$\Rightarrow Prc = \frac{1}{4}(Prc_1 + Prc_2 + Prc_3 + Prc_4) = \frac{39}{121}$$

Exercise 3.4

1.

We have $Pr(T = n) = p(1 - p)^{n-1}, n = 1, 2, \dots$. Thus, T obeys geometric distribution.

Then

$$E(T^2) = \sum_{n=1}^{\infty} n^2 p(1 - p)^{n-1}$$

Here, we have

$$\sum_{n=1}^{\infty} nx^{n-1} = \left(\sum_{n=1}^{\infty} x^n \right)' = \left(\frac{x}{1-x} \right)' = \frac{1}{(1-x)^2}$$

and

$$\sum_{n=1}^{\infty} n^2 x^{n-1} = \left(\sum_{n=1}^{\infty} nx^n \right)' = \left(x \sum_{n=1}^{\infty} nx^{n-1} \right)' = \left(\frac{x}{(1-x)^2} \right)' = \frac{1+x}{(1-x)^3}$$

Then we order $x = 1 - p$. We have

$$E(T^2) = \sum_{n=1}^{\infty} n^2 p(1 - p)^{n-1} = p \sum_{n=1}^{\infty} n^2 (1 - p)^{n-1} = p \times \frac{2-p}{p^3} = \frac{2-p}{p^2}$$

2.

$$\begin{aligned}
E\left(\frac{1}{T}\right) &= \sum_{n=1}^{\infty} \frac{1}{n} p(1-p)^{n-1} \\
&= \frac{p}{1-p} \sum_{n=1}^{\infty} \frac{(1-p)^n}{n} \\
&= \frac{p}{1-p} \sum_{n=1}^{\infty} \int_1^p -(1-p)^{n-1} dp \\
&= \frac{p}{1-p} \int_1^p \left\{ \sum_{n=1}^{\infty} -(1-p)^{n-1} \right\} dp \\
&= \frac{p}{1-p} \int_1^p - \lim_{n \rightarrow \infty} \frac{1 - (1-p)^n}{p} dp \\
&= \frac{p}{1-p} \int_1^p -\frac{1}{p} dp \\
&= -\frac{p \ln p}{1-p}
\end{aligned}$$