

CS499 Homework 8

Interstellar

Exercise 8.1

1.




Figure 1:

2. The size of the largest chain is 4.

3. The size of the largest antichain is 7.

Exercise 8.2

1. $(0, 0, 0, 0, 0, \dots)$ are minimal. There is not a maximum.

2. There is a minimum, but there is not a maximum.

3. Yes. For example,

$$(1, 1, 1, 1, \dots)$$

$$(2, 1, 1, 1, \dots)$$

$$(3, 1, 1, 1, \dots)$$

054
055
056
057
058
059
060
061
062
063
064
065
066
067
068
069
070
071
072
073
074
075
076
077
078
079
080
081
082
083
084
085
086
087
088
089
090
091
092
093
094
095
096
097
098
099
100
101
102
103
104
105
106
107

$$(4, 1, 1, 1, \dots)$$

\vdots

4.

When $n = 1$, we can't find an antichain.

When $n \geq 2$, we can find an antichain. For example,

$$(0, k, 0, \dots, 0)$$

$$(1, k - 1, 0, \dots, 0)$$

$$(2, k - 2, 0, \dots, 0)$$

\vdots

$$(k - 1, 1, 0, \dots, 0)$$

Exercise 8.3

Yes. We prove it by mathematical induction.

(1) When $n = 1$, obviously, we can sort elements from small to large to get an infinite chain.

(2) We suppose every infinite subset $S \subseteq N_0^n$ contain an infinite chain, then when $n = N + 1$, we consider the last element of the previous chain. There are two situations:

1. The sequence of the last element is bounded.

Obviously, at least one value (we call it x) appears infinite times in the sequence of the last element. We select those set with last element that equals to x , obviously it forms a infinite chain of N_0^{n+1} .

2. The sequence of the last element is unbounded.

Since each element is non-negative integer and the sequence is unbounded, we can find a sub-sequence of this sequence which is monotonically increasing. We select those set with last element in this sub-sequence, then it forms a infinite chain of N_0^{n+1} .

So every infinite subset $S \subseteq N_0^n$ contain an infinite chain.

Exercise 8.4

We assume that (\mathbb{N}_0^n, \leq) has an infinite antichain T . Obviously, $T \subseteq (\mathbb{N}_0^n, \leq)$. According to **exercise 8.3**, we get T contains an infinite chain, which contradicts T is an infinite antichain. Thus (\mathbb{N}_0^n, \leq) has no infinite antichain.

Exercise 8.5

$n = 2$




Figure 2:

$n = 3$




Figure 3:

Exercise 8.6

Maximum is $(\underbrace{1, 1, 1, \dots, 1}_n)$

Minimum is $(\underbrace{0, 0, 0, \dots, 0}_n)$

Maximal is $(\underbrace{1, 1, 1, \dots, 1}_n)$

Minimal is $(\underbrace{0, 0, 0, \dots, 0}_n)$

Exercise 8.7

We construct the Hasse diagrams like **Exercise 8.5**. The first layer is an element with n one. The second layer is n elements with $n - 1$ one. The third layer is $\binom{n}{2}$ elements with $n - 2$ one.....The k layer is $\binom{n}{k-1}$ elements with $n + 1 - k$ one, where $1 \leq k \leq n + 1$.

Obviously elements in the same layer is antichain, since elements in chain must have different numbers of one and in the same layer the number of one is equivalent. Thus it has $n + 1$ antichain partition.

According to Mirsky's Theorem, max size of chain=min size of antichain partition. It means max size of chain must less than or equal to the size of each antichain partition. Thus the longest chain of $\{0, 1\}^n \leq n + 1$. Since we get $n + 1$ antichain partition in above Hasse diagrams, then the longest chain of $\{0, 1\}^n$ is $n + 1$. For example, $(0, 0, \dots, 0), (0, 0, \dots, 1), (0, 0, \dots, 0, 1), \dots, (1, 1, \dots, 1)$ is a chain with $n + 1$ elements.

Exercise 8.8

The largest antichain of $\{0, 1\}^n$ is $\binom{n}{\lfloor n/2 \rfloor}$. According to the Dilworth Theorem, the largest antichain equals to the minimum size of chain partition.

We define a layer as a set of strings containing same number of '1' and is sorted by how many '1' a string in this layer contains.

1. There are $\binom{n}{\lfloor n/2 \rfloor}$ strings in the middle layer, which has the most strings. Since any two strings from the same layer are not comparable, there are at least $\binom{n}{\lfloor n/2 \rfloor}$ chain partitions.

2. All strings in any layer except the middle one can form chains with unique strings in its adjacent layer with the following method:

Assuming there are more '1' than '0' in this layer, we calculate a strings score by the following rules: scan the string from the beginning and the initial score is 0, add one if current digit is 1, minus one

162 otherwise. Find the digit where the first highest score appears (which must be a '1'),change it to 0.
 163 Then we get a string belongs to its adjacent layer and these two strings can form a chain(they are
 164 comparable).Now we prove that this string is unique:

165 Assume that there are two different strings that transform into a same string. Assume that the first
 166 string changes the i -th digit, and the other changes the j -th digit (with no loss of generality, assume
 167 $i < j$). Then the i -th digit of the second string and the j -th digit of the first string are 0, whereas
 168 other digits are the same. Assume that the score of the $(i - 1)$ -th digit is k .Then the score of the i -th
 169 digit is $k + 1$ for the first string and $(k - 1)$ for the second. Assume that the score of the $(j - 1)$ -th
 170 digit for the first string is $k + 1 + p$,then the score of the $(j - 1)$ -th digit for the second string is
 171 $k - 1 + p$.The score of the j -th digit for the second string is $k + p$. Since the changing digit is where
 172 the first largest score occurs, we have

$$173 \quad k + 1 \geq k + 1 + p$$

$$174 \quad k < k + p$$

176 where we get $p \geq 0$ and $p > 0$ which contradict each other. So different strings cannot transform
 177 into a same string by the method. Similarly, if there are more '0' than '1' in the layer,we can use a
 178 similar method to form chains with unique strings in its adjacent layer.

179 **Question**

181 **1**How to prove the proposition that the set of integer has the smallest cardinality among all the
 182 infinite sets?

183 **2**How to handle the Russell's paradox?
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215