

CS499 Homework 5 (First Draft)

Interstellar

Exercise 5.1

(1) The degree of a vertex is defined as the number of edges linked to this vertex. And the score of a graph is a sequence ranking degree of all vertices from small to big.

(2) Graph score theorem states that, if we can find a graph for graph score $(d_1, \dots, d_{n-1}, d_n)$, then we can find a graph for graph score $(d_1, \dots, d_{n-d_n-1}, d_{n-d_n} - 1, \dots, d_{n-1} - 1)$, and vice versa. If we finally get graph score (ϕ) , the graph exists.

(3) Graph score algorithm:

First, we get a graph score $(d_1, \dots, d_{n-1}, d_n)$.

If $d_n > n - 1$, we cannot find a graph. Otherwise, we add an edge from d_n to $d_{n-d_n}, \dots, d_{n-1}$, and check the graph score $(d_1, \dots, d_{n-d_n-1}, d_{n-d_n} - 1, \dots, d_{n-1} - 1)$ after it is sorted.

We repeat the previous step. If the graph score finally comes to (ϕ) , the graph exists.

(4) The most difficult part is to prove if we can find a graph for graph score $(d_1, \dots, d_{n-1}, d_n)$, then we can find a graph for graph score $(d_1, \dots, d_{n-d_n-1}, d_{n-d_n} - 1, \dots, d_{n-1} - 1)$. We can suppose there is a solution without edge between n and k ($n - d_n \leq k \leq n - 1$), so n must have another link with j ($j \leq n - d_{n-1} < k$). As $j < k$, we know $d_j \leq d_k$, so k must have edge with some point l and $l \neq k$. We change the edges (n, j) (k, l) to (n, k) (j, l) , and we add an edge between n and k without changing the score. In this way, we can transform the answer to make sure there is an edge from d_n to $d_{n-d_n}, \dots, d_{n-1}$. Then we delete these edges, we get a graph for score $(d_1, \dots, d_{n-d_n-1}, d_{n-d_n} - 1, \dots, d_{n-1} - 1)$.

Exercise 5.2

Let's define an operation of a sequence: subtract 1 from the two largest number.

Theorem 5.3 can be written as: Let $(a_1, \dots, a_n) \in N_0^n$. There is a multigraph with this score if and only if after $\frac{\sum_{n=1}^N a_n}{2}$ operations, all the numbers in the sequence are 0 (we name a sequence consists of 0 a "zero sequence").

Exercise 5.4

proof

(1) If there is a multigraph with score (a_1, \dots, a_n) satisfying $a_1 \leq a_2 \leq \dots \leq a_n$:

(i) if $a_{n-1} = 1$, there must be an even number of 1s, it's obvious that we can change its score sequence to a zero sequence through several operations.

(ii) if $a_{n-1} = 1$ and there is an edge (v_{n-1}, v_n) , delete this edge and we get a multigraph with score $(a_1, \dots, a_{n-1} - 1, a_n - 1)$. Repeat this operation and we would get a zero sequence eventually.

(iii) else there must be an edge (v_k, v_n) and an edge (v_l, v_{n-1}) such that $k \neq l$. Delete these two edges and add an edge (v_k, v_l) and we get a multigraph with score $(a_1, \dots, a_{n-1} - 1, a_n - 1)$. Repeat this operation and we would get a zero sequence eventually.

(2)

If a sequence can transform into a zero sequence, we can get a multigraph with the score sequence by undoing an operation and add an edge accordingly $\frac{\sum_{n=1}^N a_n}{2}$ times.

Theorem 5.3 proves ture based on **1** and **2**.

Exercise 5.5&5.6&5.7

Let $(a_1, \dots, a_n) \in \mathbb{R}_0^n$. There is a weighted graph with this score if and only if suppose a_1, \dots, a_n are arranged from small to large, $(a_n) \geq \sum_{i=1}^{n-1} a_i$

proof

(1) When $n = 2$, it is obvious that there is a weighted graph if and only if $(a_1) = (a_2)$.

(2) When $n = 3$, suppose $a = a_3 = wdge(A_3)$, $b = a_2 = wdge(A_2)$, $c = a_1 = wdge(A_1)$,

$x = W_{\{A_3, A_2\}}$, $y = W_{\{A_3, A_1\}}$, $z = W_{\{A_1, A_2\}}$, $a \geq b \geq c$, then we have

$$\begin{cases} x + y = a \\ x + z = b \\ y + z = c \end{cases} \quad (1)$$

$$\Rightarrow$$

$$\begin{cases} x = \frac{a+b-c}{2} \\ y = \frac{a+c-b}{2} \\ z = \frac{b+c-a}{2} \end{cases} \quad (2)$$

So

if there is a weighted graph, then

$$x \geq 0 \Rightarrow a + b - c \geq 0$$

$$y \geq 0 \Rightarrow a + c - b \geq 0$$

$$z \geq 0 \Rightarrow b + c - a \geq 0 \Rightarrow a \leq b + c \Rightarrow a_3 \leq a_1 + a_2.$$

if $a \geq b \geq c$ and $a \leq b + c$, then

$x \geq 0, y \geq 0, z \geq 0 \Rightarrow$ there is a weighted graph.

(3) Suppose when $|V| = n$ the theorem is right, we talk about the situation of $|V| = n + 1$.

If $a_{n+1} \leq \sum_{i=1}^n a_i$, then we have

if $a_{n+1} \geq a_n + a_1$, we have $(a_2, \dots, a_n, a_{n+1} - a_1)$ and $a_{n+1} - a_1 \leq \sum_{i=2}^n a_i$, so there is a weighted graph G whose score is $(a_2, \dots, a_n, a_{n+1} - a_1)$. Now we add vertex u and $edge(u, v)$ whose weight is a_1 ($wdge(v) = a_{n+1} - a_1$ in G) to get G' . Obviously the score of G' is $(a_1, \dots, a_n, a_{n+1})$.

if $a_{n+1} < a_n + a_1$, we have $(a_2, \dots, a_n - \frac{a_1}{2}, a_{n+1} - \frac{a_1}{2})$, we have two situations:

1.

$a_{n+1} - \frac{a_1}{2}$ is the $\max\{a_2, \dots, a_n - \frac{a_1}{2}, a_{n+1} - \frac{a_1}{2}\}$.

Obviously $a_{n+1} - \frac{a_1}{2} < a_n - \frac{a_1}{2} < a_2 + \dots + a_n - \frac{a_1}{2}$

So we have a weighted graph G whose score is $(a_2, \dots, a_n - \frac{a_1}{2}, a_{n+1} - \frac{a_1}{2})$.

Now we add vertex u and $edge(u, v)$ whose weight is $\frac{a_1}{2}$ ($wdge(v) = a_{n+1} - \frac{a_1}{2}$ in G) and $edge(u, v')$ ($wdge(v') = a_{n+1} - \frac{a_1}{2}$ in G) to get G' .

Obviously the score of G' is $(a_1, \dots, a_n, a_{n+1})$.

2.

a_{n-1} is the $\max\{a_2, \dots, a_n - \frac{a_1}{2}, a_{n+1} - \frac{a_1}{2}\}$.

Obviously $a_{n-1} < a_n < a_n + a_{n+1} - a_1 = a_n - \frac{a_1}{2} + a_{n+1} - \frac{a_1}{2} \leq a_2 + \dots + a_n - \frac{a_1}{2} + a_{n+1} - \frac{a_1}{2}$.

So we have a weighted graph G whose score is $(a_2, \dots, a_n - \frac{a_1}{2}, a_{n+1} - \frac{a_1}{2})$.

In a similar way we get a weighted graph G' whose score is $(a_1, \dots, a_n, a_{n+1})$.

If there is a weighted graph whose score is (a_1, \dots, a_n) , then we have

$$a_n \leq \sum_{i=1}^n i = 1n - 1a_1. \text{ Otherwise this graph is unable to satisfy } a_n.$$

Exercise 5.8&5.9

(Score Theorem for Graphs with Real Edge Weights). Let $(a_1, \dots, a_n) \in \mathbb{R}^n$. There is a graph with real edge weights with this score if and only if $n = 2$ and $a_1 = a_2$ or $n \geq 3$.

Exercise 5.10

Proof: If $n = 2$ and $a_1 = a_2$, it is obviously true.

Consider $n \geq 3$, we let the graph be a n polygon. For each edge between adjacent vertexes, the edge weight is x_i . Thus we have $(x_1, \dots, x_n) \in \mathbb{R}^n$ and n equations:

$$x_n + x_1 = a_1$$

$$x_1 + x_2 = a_2$$

$$x_2 + x_3 = a_3$$

$$\dots$$

$$x_{n-1} + x_n = a_n$$

Obviously there are n variables and n equations, so there must exist real solutions for x_i .

Thus, for any $(a_1, \dots, a_n) \in \mathbb{R}^n$ and $n \geq 3$, there must exist a graph with real edge weights with this score.

Exercise 5.11

(For convenience, we ignore the 0 (a dot) in the following)

ID:517030910250

- (1)(2) It is neither a graph score nor a multigraph score because the sum of the ID is an odd number.
- (3) It is a weighted graph score, as is shown in the following figure.




Figure 1:

162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215

4Since it is a weighted graph score, it is the score of a graph with real edge weights.

ID:517030910258

(1)(2) It is neither a graph score nor a multigraph score because the sum of the ID is an odd number.

(3) It is a weighted graph score, as is shown in the following figure.




Figure 2:

4Since it is a weighted graph score, it is the score of a graph with real edge weights.

ID:517030910029

(1)(2) It is neither a graph score nor a multigraph score because the sum of the ID is an odd number.

(3) It is a weighted graph score, as is shown in the following figure.




Figure 3:

4Since it is a weighted graph score, it is the score of a graph with real edge weights.

ID:517030910227

(1)(2) It is neither a graph score nor a multigraph score because the sum of the ID is an odd number.

(3) It is a weighted graph score, as is shown in the following figure.




Figure 4:

4Since it is a weighted graph score, it is the score of a graph with real edge weights.

ID:517030910263

(1)(2) It is neither a graph score nor a multigraph score because the sum of the ID is an odd number.

(3) It is a weighted graph score, as is shown in the following figure.




Figure 5:

4Since it is a weighted graph score, it is the score of a graph with real edge weights.

Questions